



## PHARMACOGNOSTIC EVALUATION OF *OSYRIS QUADRIPARTITA* SALZ. EX DECNE

Balasaheb Shantilal Kale<sup>1\*</sup>, Mangesh Shankar Bhale<sup>2</sup>, Aditya Bajirao Bhagat<sup>1</sup>, Sanjay Appaji Khairnar<sup>1</sup>

1. Department of Botany, S.V. K. T. Arts, Science & Commerce College, Deolali Camp, Nashik (M.S.) India.
2. Department of Chemistry, Arts, Commerce and Science College, Jawhar, Palghar (M.S.) India.

### ARTICLE INFO

#### Received:

03 April 2022

#### Received in revised form:

10 Jun 2022

#### Accepted:

12 Jun 2022

#### Available online:

28 Jun 2022

**Keywords:** *Osyris quadripartita*, Authentication, Medicinal plant, Anatomy, Pharmacognosy, Phytochemistry

### ABSTRACT

The comprehend assess the pharmacognostic characteristics of an important ethnomedicinal plant, *O. quadripartita* Salz. ex Decne. The pharmacognostic evaluation was executed in terms of taxonomic, macroscopic, microscopic, fluorescence analysis, and phytochemical analysis. The *Osyris* members heartwood majorly used to make furniture and perfumes. The *Osyris* plant members are known for fragrance and volatile oil constituents. The microscopic features of stem bark were observed as cork, cortex, phloem fibers, phloem medullary rays, and sclerides in the transverse section (T.S.) and longitudinal section (L.S.). The microscopy of stem bark powder observed the presence of cortex cells, sieve tubes, calcium oxalate crystals, and lignified fibers. The initial screening of phytochemical constituents of stem bark crude plant extracts indicate the presence of carbohydrates, proteins, alkaloids, glycosides, flavonoids, saponins, tannins & phenolic in different polar and non-polar solvent systems such as petroleum ether, chloroform, acetone, methanol, and water. Many pharmacognostic characters observed during the study, it's help in the standardization of *O. quadripartite*.

This is an **open-access** article distributed under the terms of the [Creative Commons Attribution-Non Commercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, and build upon the work non commercially.

**To Cite This Article:** Kale BS, Bhale MS, Bhagat AB, Khairnar SA. Pharmacognostic Evaluation of *Osyris quadripartita* Salz. ex Decne. Pharmacophore. 2022;13(3):50-6. <https://doi.org/10.51847/nzKoIHDQUz>

### Introduction

In the world about 3000 to 3150 plant species used as medicinal purpose from ancient times [1]. Various different types secondary metabolites recorded in plant kingdom such as alkaloids, tannins, phenols saponins flavonoids etc., these phytochemicals used against many diseases [2, 3]. Crude drug discovery from natural medicinal plant resources these natural products very useful for the treatment of various disorders or diseases [4, 5]. *O. quadripartita* is a dry deciduous forest of family Santalaceae family and is commonly known as a wild tea plant and African sandalwood. Making the strong red-colored dye from the stem bark. The bark and heartwood of *O. quadripartite* can be easily adulterated with low-grade materials such as heartwood (African sandalwood) and strong red-colored dye [6, 7]. *O. quadripartita* comes under the least concern IUCN category [8]. The heartwood of this plant is faintly fragrant so, this plant is used as adulterating sandalwood in the commercial market. The stem bark is important for tanning the leather. The bark is used as a tonic, aromatic oils used in different products for manufacturing expensive perfumes, rare soaps, quality lotions, and sweet-smelling candles [9, 10]. In Folk, Indian, and other traditional medicine systems, bark and leaves are used by tribal peoples this plant leaves uses as making herbal tea [11, 12]. It is used as curing various diseases such as antihelmintic, leishmaniasis like leprosy diseases, abnormalities, kidney disease, forage, hygienic/perfuming, eye pain, cold, anaphylactic shock, epilepsy, circumcision wound, toothache, tonsillitis, abdominal pain, and urine problem [13, 14]. It's also used for the treatment of cancer or tumor [15, 16], breast cancer [15, 17], anthrax and pasturolosis [18], Jaundice [19], wild edible fruit [20, 21], stomachache, malaria, cough, swelling and anti-corona virus disease (COVID-19) [22, 23], reduce inflammation and powerful emetic [24, 25], diarrhea [26], rabies [27], wound healing plants [28, 29], anti-ulcer activity [30, 31], mental problems [32], mastitis and poor mothering and antiviral activity [33]. This adulteration can be avoided by means of various evaluation variables such as microscopic as well as macroscopic evaluation. Macroscopic, microscopic, and chemical profiling evaluation techniques are

**Corresponding Author:** Balasaheb Shantilal Kale; Department of Botany, S.V. K. T. Arts, Science & Commerce College, Deolali Camp, Nashik (M.S.) India. E-mail: [kaleunipune@gmail.com](mailto:kaleunipune@gmail.com).

important tools for the authentication of plant based crude drugs and the study of powdered drugs. Standardize of this plant part as per Indian Pharmacopoeia [34, 35]. It is significantly important characters such as morphological and anatomical characters, which is useful for drugs and their adulterants of commercial significance. The significant setup of the macroscopically, microscopically, and Pharmacognostic parameters of stem bark of the medicinal plant, will help in regularized, which can help quality control, identification, and purity of the sample [36].

## Materials and Methods

### Chemicals

The analytical grade chemical used for staining of the microscopic parts of the studied medicinal plant of E. Merck Limited India and Hi-Media Laboratories, Mumbai, India used for analysis of sample.

### Botanical Parameter

*Taxonomy & Morphology:* Identification and classification of plants using different Floras and other available literature [37].

### Collection Plant Materials

The fresh stem bark part of *O. quadripartite* medicinal plant, were collected from different regions of Junar taluka, Pune districts of Maharashtra (India). The plant materials were taxonomically identification, authenticated by BSI, Pune reference numberis BSIS000001612. The plant materials dried in shade until all the water get evaporated and plant part became well dried for grinding. After drying, the plant materials were crush using mechanical blender (Mixer) into fine powder. This dried sample make fine powder and use for phytochemical and Pharmacognostic analysis.

### Pharmacognostic Studies- Microscopic Characters

Plant anatomy Microscopical characters were studied by hand section of the stem *Osyris quadripartite* specimen. Wherever sections stained with safranin, fastgreen and iodine to identify and detection of the living tissues, dead tissues and starch granuels [38, 39].

### Microscopic Examination of Powder Sample by Using Maceration Technique

Study of stem bark elements such as vessels, cortex cells, sieve tubes etc. Use a maceration technique by Jeffery's fluid [38, 39].

### Sensory Evaluation

Various sensory variable done of the plant sample such as odour, taste, colour, size and shape.

### Physico- Chemical Parameters- Extractives Values

Shade dried stem bark powder of *O. quadripartite* was subjected to constantly extraction with cold decoction for water only and hot extraction.

### Ash Values

Water soluble ash, acid insoluble ash value and total ash value calculated by using plant sample **Table 2**.

### Powder Microscopy

Take crude powder sample dissolved in water, then takes drop of this sample on slide, then put a coverslip and observed under light microscope.

### Fluorescence Analysis

Take crude powder sample treated various chemicals then observe under visible light and U.V light **Table 4** [40].

### Method of Extraction

The powder of shade dried stem bark of *O. quadripartita* was subjected to continuous extraction with cold decoction for water only and maceration hot extraction used differnet organic solvents such as petroleum ether, chloroform, acetone & methanol [40].

### Phytochemicals Qualitative Tests

Followings phytochemicals qualitative tests used for analysis of *O. quadripartita* plant crude extracts, there are test for carbohydrates, amino acids, proteins, alkaloids, tannins glycosides, phenolic compounds, flavonoids, saponins & steroids [40].

## Results and Discussion

### Sensory Evaluation


The Sensory evaluation of the stem bark showed dark brown color on the upper surface and a light brown color on the lower surface. The shade dried stem bark was dark brown in color, with a bitter taste, unpleasant in odor, and coarse texture [38].

### Macroscopic Evaluation

The morphological and taxonomy characters are much branched shrubs; young branches sharply triangular. Leaves subsessile, coriaceous, elliptic-oblong, apex mucronate. Flowers yellowish-green, axillary; male flowers in long pedunculate, shortly pedicellate, 5–10-flowered umbels, the peduncles of bisexual flowers carrying 1–3 flowers and elongating in fruit, perianth of 3 triangular lobes. The dried stem bark thickness was about 3.37 mm [37].


### Microscopic Evaluation

A superficial and strongly thickened cork layer was observed in the species. The cortex cells are arranged in compact form. The pericycle is a part of ground tissue. Ground tissue's primary function is mechanical support. Pericyclic fiber is sclerenchyma in nature. The phloem is stratified into soft and hard portions. In which hard portions are nothing but primary phloem fibers, proto-phloem fibers, and secondary phloem fibers. The various cluster of crystalline structures deposition was present in the phloem cells **Figure 1** [39, 41].



**Figure 1.** a and b) Stem bark anatomy of *O. quadripartita*

Bl= Barklayer, Cl=Cortex layer, Ph = phloem, Pm= Phloem medullary layer, Sc= Sclerides, Cr= Crystalliferous cells, Tnd= Tannin deposition, Sg= Starch granules



**Figure 2.** Microscopic observation of stem bark *O. quadripartita* Coc= Calcium oxalate crystals, Tn= Tannin, Bc= Bark cells

### Physical Evaluation

The physical standards, such as ash values, and extractive values will be helpful to identify the standardness of the superior quality drugs, secondary drug, and drug adulteration from the powdered plant materials. It will maintain significant quality control of the drug obtained from medicinal plant sources. The facts collected from the crude bark powder ash values and extractive values are helpful for the quality control of drugs. Physical evaluation results are water-soluble ash 44.33%; acid

insoluble ash 21.66%; total ash 32.00 %; alcohol soluble extractives 97.6 % and water-soluble extractives 89.2 % w/w values were observed in dried stem bark of *O. quadripartita*. The results were mentioned in **Table 1** [40].

**Table 1.** Physical parameters of stem bark of *O. quadripartita*

Sr. No.	Parameters	Determined value* (% w/w)
<b>Ash values</b>		
1	Total ash	32.00
	Acid insoluble ash	21.66
	Water soluble ash	44.33
<b>Extractive values</b>		
2	Alcohol soluble extractives	6.2
	Water soluble extractives	11.8

\* Mean value of three readings

**Table 2.** Stem bark average extractive values of *O. quadripartita*

Sr No.	Solvent used	Colour appeared	Average percentage yield (% w/w on dry weight basis)
1.	Petroleum ether	Light brown	2.4
2.	Chloroform	Coffee brown	3.6
3.	Acetone	Dark brown	9.4
4.	Methanol	Dark brown	7.2

#### Average Extractive Values (% w/w)

During the experimental work of consecutively solvent extraction, the overall percentage yields obtained, were purposeful as petroleum ether (94.6 %), chloroform (92.2 %), acetone (85.4 %), and methanol (97.0 %) w/w. The crude dried stem bark powder color developed during the successive extraction process as noted in **Table 2**.

#### Fluorescence Analysis

The fluorescence analysis results obtained were mentioned in **Table 3**.

**Table 3.** Fluorescence analysis of *O. quadripartita* stems bark powder.

Sr. No.	Test	Visible Light	UV light
1.	Powder as such	Cinnamon brown	Pecan brown
2.	Powder + 1N NaOH (aqueous)	Brown	Greenish brown
3.	Powder + 1N NaOH (alcoholic)	Brown	Greenish black
4.	Powder +50% HNO <sub>3</sub>	Yellowish brown	Yellowish green
5.	Powder + 50 % H <sub>2</sub> SO <sub>4</sub>	Dark brown	Mehndi green
6.	Powder + 1N HCl	Brown	Coffee brown
7.	Powder + 5% KOH	Dark brown	Greenish black
8.	Powder +Methanol	Brown	Greenish brown
9.	Powder + Petroleum ether	Light brown	Ceramic brown
10.	Powder + Chloroform	Brown	Coffee brown

#### Powder Microscopy

Powdered stem bark of *O. quadripartita* under compound light microscopic observation showed tannin and cortex cells, calcium oxalate crystals. The calcium oxalate crystals of cubical type were observed in **Figure 2**.

#### Preliminary Phytochemical Screening of *O. Quadripartita* Stem bark

The ethnomedicinal therapeutic properties of medicinal plants presence of several secondary metabolites and chemical components like alkaloids, glycosides, flavonoids, saponins, tannins & phenolics. This preliminary analysis of phytochemical tests helpful to the detection of active principles of chemical components and may accelerate to novel drug discovery in the pharmacognosy and pharmacology field. The preliminary crude extract of phytochemical screening leads to

revealing the existence of carbohydrates, proteins, alkaloids, glycosides, flavonoids, saponins, tannins & phenolics present in the dried stem bark crude extracts of *O. quadripartita*. The results noted in the **Table 4** [40].

**Table 4.** Qualitative tests for preliminary phytochemical extracts of *O. quadripartita*

Phytochemical Constituents	Tests	Results				
		Petroleum ether	Chloroform	Acetone	Methanol	Water
Carbohydrates	Benedict's test	+	+	+	+	+
	Barfoed's test	-	-	+	+	+
Starch	Iodine test	-	-	-	-	-
Proteins & amino acids	Xanthoproteic test	+	+	-	-	+
	Millon's test	+	+	-	-	-
Alkaloids	Mayer's test	+	+	-	-	-
	Wagner's test	-	+	+	+	+
Glycosides	Legal's test	-	-	+	+	+
	Borntrager's test	-	-	+	+	+
Tannins & phenolic compounds	Ferric chloride test	-	-	+	+	+
	Potassium dichromate test	+	+	-	-	-
Flavonoids	Concentrated H <sub>2</sub> SO <sub>4</sub> test	-	+	-	-	+
	10% sodium chloride test	-	+	-	-	-
Saponins	Foam test	+	+	+	+	+

**Legend:** (-) absent, (+) present

## Conclusion

A study on morphological and anatomical characters of *O. quadripartita*, significant anatomical features helpful to maintain its quality control. The morphological, and anatomical characters of this plant, are the primary steps to establishing its botanical excellence. According to WHO rules, botanical standards set a proposed for the determination of the plant based herbal drugs. The pharmacognostic parameters is supportive for the future recognition, identification, and authentication of in the medicinal plant-based industrial companies such as herbal, cosmetics, and pharmaceutical. The physical quality analysis of stem bark plant material, like ash values (water soluble ash, acid insoluble ash, and total ash), and extractive values will be useful for recognition, identification, and standardization of the dried stem bark powdered of medicinal plant material. The scientific study developed standard quality data used for the quality control of the various medicinal preparations, were helpful in the future for drug development. The ash values as well as extractive values are very much helpful for the standardization of crude drugs. The physicochemical standards of this plant can be separated from other related species. The dried stem bark constituents can be incorporate as macroscopic and microscopically standards as per Indian Pharmacopoeia. This medicinal plant in dried stem bark has some secondary metabolites reported like alkaloids, glycosides, saponins, flavonoids, tannins and phenolic. The calcium oxalate crystals of cubical type observed in the cortex region of stem bark. Thus this preliminary analysis of phytochemical tests will use in the find out of the active essential chemical components and later may guide to novel drug finding. These novel pharmacologically active principal chemical components will be used for curing various diseases and abnormalities.

**Acknowledgments:** None

**Conflict of interest:** None

**Financial support:** None

**Ethics statement:** None

## References

1. Chandra S, Meel RK. Pharmacognostic, phytochemical and pharmacological assessment of *Morinda tinctoria* as anti-ulcerogenic potential. *Pharmacophore*. 2020;11(6):1-12.

2. Yasin G, Ahmad M, Hussain M. Pharmacological potential of plants from Himalyan region of Pakistan-assay for antioxidants indices. *Pharmacophore*. 2020;11(3):82-8.
3. Akshita C, Vijay BV, Praveen D. Evaluation of phytochemical screening and antimicrobial efficacy of *Mesua Ferrea* and *Piper cubeba* fruit extracts against multidrug resistant bacteria. *Pharmacophore*. 2020;11(2):15-20.
4. Deodhar KA, Shinde N. Pharmacognostic evaluation of root of *Celastrus paniculatus* Willd. *J Pharmacogn Phytochem*. 2014;3(3):134-6.
5. Shehu S, Abubakar MS, Ibrahim G, Iliyasu U. Phytochemical and Antibacterial Studies on Aqueous Ethanol Extract of *Thesium viride* (Santalaceae). *Br J Pharm Res*. 2016;11(2):1-6. doi:10.9734/BJPR/2016/23046
6. Silva JAT Da, Kher MM, Soner D, Nataraj M. African Sandalwood or Nepalese Sandalwood: a Brief Synthesis. *Not Sci Biol*. 2016;8(1):57-61. doi:10.15835/nsb.8.1.9714
7. Kumar S, Chaitanya MK, Andrew JS, Krishna V. Indigenous knowledge of medicinal plants used by ethnic communities of South India. *Ethnobot Res Appl*. 2019;18(4):1-112. doi:10.32859/era.18.4.1-112
8. Wilson B. *Osyris lanceolata*. IUCN Red List Threat Species. 2018;8235:1-8.
9. Khare CP. *Indian Medicinal Plants*. Springer New York, NY; 2007.
10. Shyaula SL. A review on genus *Osyris*: Phytochemical constituents and traditional uses. *J Nat Pharm*. 2012;3(2):61-70. doi:10.4103/2229-5119.102747
11. Herrera CM. The Annual Cycle of *Osyris Quadripartita*, A Hemiparasitic Dioecious Shrub of Mediterranean Scrublands. *J Ecol*. 1984;72:1065-78.
12. Kefalew A, Asfaw Z, Kelbessa E. Ethnobotany of medicinal plants in Ada'a District, East Shewa Zone of Oromia Regional State, Ethiopia. *J Ethnobiol Ethnomed*. 2015;11(25):1-28. doi:10.1186/s13002-015-0014-6
13. Mouhajir F, Pedersen JA, Rejdali M, Towers GHN. Phenolics in Moroccan Medicinal Plant Species as Studied by Electron Spin Resonance Spectroscopy. *Pharm Biol*. 2001;39(5):391-8. doi:10.1076/phbi.39.5.391.5893
14. Yohannis SW, Asfaw Z, Kelbessa E. Ethnobotanical study of medicinal plants used by local people in Menz Gera Midir District, North Shewa Zone, Amhara Regional State, Ethiopia. *J Med Plants Res*. 2018;12(21):296-314. doi:10.5897/JMPR2018.6616
15. Graham JG, Quinn ML, Fabricant DS, Farnsworth NR. Plants used against cancer – an extension of the work of Jonathan Hartwell. *J Ethnopharmacol*. 2000;73(3):347-77.
16. Esubalew ST, Belete A, Lulekal E, Gabriel T, Engidawork E, Asres K. Review of Ethnobotanical and Ethnopharmacological Evidences of some Ethiopian Medicinal Plants traditionally used for the Treatment of Cancer. *Ethiop J Heal Dev Resour*. 2017;31(3):161-87.
17. Tesfaye S, Belete A, Engidawork E, Gedif T, Asres K. Ethnobotanical Study of Medicinal Plants Used by Traditional Healers to Treat Cancer-Like Symptoms in Eleven Districts, Ethiopia. *Evid Based Complement Altern Med*. 2020;1(1):1-23.
18. Advisor KT. Ethnobotanical Study of Medicinal Plants Used by the Local People of Akaki District, East Shewa Zone, Oromia Regional State, Ethiopia. Published online 2017.
19. Maryo M, Nemomissa S, Bekele T. An ethnobotanical study of medicinal plants of the Kembatta ethnic group in Enset-based agricultural landscape of Kembatta Tembaro (KT) Zone, Southern Ethiopia. *Asian J Plant Sci Res*. 2015;5(7):42-61.
20. Arinathan V, Mohan VR, Britto AJ De, Murugan C. Wild edibles used by Palliyars of the western Ghats, Tamil Nadu. *Indian J Tradit Knowl*. 2007;6(1):163-8.
21. Meragiaw M, Asfaw Z, Argaw M. Indigenous Knowledge (IK) of Wild Edible Plants (WEPs) and Impacts of Resettlement in Delanta, Northern Ethiopia. *Res Rev J Herb Sci*. 2015;4(3):8-26.
22. Belayneh A, Bussa NF. Ethnomedicinal plants used to treat human ailments in the prehistoric place of Harla and Dengego valleys, eastern Ethiopia. *J Ethnobiol Ethnomed*. 2014;10(18):1-17. doi:10.1186/1746-4269-10-18
23. Gonfa T, Kiros T. Potential use of Eastern Hararghe Ethiopia medicinal plants for COVID-19- like symptoms: A review Teshome. *Arab J Med Aromat Plants*. 2020;6(3):76-115.
24. Kunwar RM, Duwadee NPS. Ethnobotanical notes on flora of Khaptad national Park (KNP), far-western Nepal. *Him J Sci*. 2003;1(1):25-30.
25. Neill ARO, Rana SK. An ethnobotanical analysis of parasitic plants (Parijibi) in the Nepal Himalaya. *J Ethnobiol Ethnomed*. 2016;12(14):1-15. doi:10.1186/s13002-016-0086-y
26. Teferi MY, Abdulwuhab M, Yesuf JS. Evaluation of in Vivo Antidiarrheal Activity of 80% Methanolic Leaf Extract of *Osyris quadripartita* Decne (Santalaceae) in Swiss Albino Mice. *J Evidence-Based Integr Med*. 2019;24:1-9. doi:10.1177/2515690X19833340
27. Chekole G. Ethnobotanical study of medicinal plants used against human ailments in Gubalafto District, Northern Ethiopia. *J Ethnobiol Ethnomed*. 2017;13(55):1-29. doi:10.1186/s13002-017-0182-7
28. Parrotta JA. *Healing Plants of Peninsular India*. CABI publishing; 2001.
29. Kassa Z, Asfaw Z, Demissew S. Ethnobotanical study of medicinal plants used by the local people in Tulu Korma and its Surrounding Areas of Ejere District, Western Shewa Zone of Oromia Regional State, Ethiopia. *J Med Plants Stud*. 2016;4(2):24-47.

30. Abebaw M, Mishra B, Asmelashe D. Evaluation of anti-ulcer activity of the leaf extract of *Osyris quadripartita* Decne. (Santalaceae) in rats. *J Exp Pharmacol.* 2017;9:1.
31. Njoku UO, Umeh CG, Ogugofor MO. Anti-ulcerogenic Activity of Methanol Fraction of *Hibiscus asper* Leaves in Albino Rats. *African J Biomed Res.* 2020;23(2):267-72.
32. Wubetu M, Sintayehu M, Aeta MA, Reta H, Derebe D. Ethnobotany of Medicinal Plants used to Treat Various Mental illnesses in Ethiopia: A Systematic Review. *Asian J Plant Sci Res.* 2018;8(1):9-33.
33. Eshetu GR, Dejene TA, Telila LB, Bekele DF. Ethnoveterinary medicinal plants: Preparation and application methods by traditional healers in selected districts of southern Ethiopia. *Vet World.* 2015;8(5):674-84. doi:10.14202/vetworld.2015.674-684
34. Govt. of India M of H and FW. *Indian Pharmacopoeia Vol.- II.* Indian Pharmacopoeia Commission, Ghaziabad; 2010.
35. Govt. of India M of H and FW. *Indian Pharmacopoeia Vol.- III.* Vol III.; 2010. doi:10.1017/CBO9781107415324.004
36. Sarvananda L, Shafras M, Premarathna AD. Adulteration methods and current trends in authentic identification of botanical materials used for the pharmaceuticals. *Int J Tradit Complement Med.* 2019;4(17):1-20. Available from: <https://www.academia.edu/download/62223523/IJTCM-2019-05-150620200228-116380-1gg05sn.pdf>
37. Singh NP, Lakshminarasimhan P, Karthikeyan S, Prasanna PV. *Flora of Maharashtra State: Dicotyledones.* Flora India Ser 2. 2001;2(1):1-1096. Available from: <https://books.google.co.in/books?id=76G3mgEACAAJ>
38. Wallis TE. *Textbook of Pharmacognosy.* J & A Churchill ltd; 1939.
39. Metcalfe CR, Chalk L. *Anatomy of the Dicotyledons- Vol. II.* Oxford University Press, USA; 1965.
40. Khandelwal KR, Sethi V. *Practical Pharmacognosy Techniques and Experiments.* 3rd ed. Nirali Prakashan; 2019.
41. Metcalfe CR, Chalk L. *Anatomy of The Dicotyledons- Vol. I.* Oxford at The Clarendon Press; 1957.