



## MAGNETIC RESONANCE IMAGING OF CHRONIC LOW BACK PAIN: CORRELATION BETWEEN PAIN, DISABILITY, AND DISC HERNIATION

Qurain Turki Alshammari<sup>1\*</sup>, Arwa Nasser Alsubaie<sup>2</sup>, Alaa Ahmed Alkayyat<sup>3</sup>, Ala' Mansour Qazaq<sup>4</sup>, Atheer Ahmed Alnukhali<sup>5</sup>, Mosaab Ghazi Alrefaei<sup>6</sup>, Rakan Nuwayhi Alrashdi<sup>7</sup>, Reem Hamad Alqasem<sup>5</sup>, Rahaf Sulaiman Aljohani<sup>8</sup>, Samirah Abdullah Alameer<sup>9</sup>, Salman Abdullah Alomaireeni<sup>10</sup>, Ahmed Mohamed Husieny<sup>11</sup>

1. *Department of Diagnostic Radiology, College of Applied Medical Sciences, University of Hail, Hail, Saudi Arabia.*
2. *Department of Diagnostic Radiology, King Salman Medical City, Medina Health Cluster, Medina, Saudi Arabia.*
3. *Department of Radiological Science, College of Applied Medical Sciences, King Saud bin Abdulaziz University for Health Sciences, Jeddah, Saudi Arabia.*
4. *Department of Radiology and Nuclear Medicine, Jordan University Hospital, Amman, Jordan.*
5. *Department of Diagnostic Radiology, College of Applied Medical Sciences, Taif University, Taif, Saudi Arabia.*
6. *Department of Radiology, King Abdullah Medical City, Makkah Health Cluster, Makkah, Saudi Arabia.*
7. *Department of Medical Imaging, King Abdulaziz Hospital for National Guard, National Guard Health Affair, Alhasa, Saudi Arabia.*
8. *Department of Radiology, Alhada Armed Forces, Taif, Saudi Arabia.*
9. *Department of Radiology, Sabya General Hospital, Jizan, Jizan Health Cluster, Saudi Arabia.*
10. *Department of Medical Imaging, King Fahad Specialist Hospital, Qassim Cluster Health, Qassim, Saudi Arabia.*
11. *Department of Radiology, Sohar Hospital, Ministry of Health, Sohar, Sultanate of Oman.*

### ARTICLE INFO

#### Received:

27 Mar 2023

#### Received in revised form:

17 Jun 2023

#### Accepted:

19 Jun 2023

#### Available online:

28 Jun 2023

**Keywords:** Disc herniation, Chronic low back pain, IVDP, MRI, Disability

### ABSTRACT

Chronic Low Back Pain (cLBP) manifests as compressed, radiating pain from the back to the nerve root's dermatome. Based on spinal imaging, patients with cLBP are diagnosed and treated. The best method of screening the spine is Magnetic Resonance Imaging [MRI]. The study involved 70 patients. A visual analog scale was employed in assessing the intensity of back and leg pain of patients using a visual analog scale [VAS]. A Roland Morris Disability Questionnaire (RMDQ-Arabic version) was used to assess the disability of patients. The patients were examined clinically, followed by an MRI of the spine and lower extremities. Using the Michigan State University [MSU] disc herniation classification, Disc displacement and the degree of nerve root compression were assessed. This study uses SPSS 23.0 version to analyze curated data in the documentation using Pearson's correlations. Based on the data, the mean and standard deviation for each descriptive variable were calculated. A low grade disc herniation and a VAS rating of 3 were not correlated based on the MRI results. The Pearson correlation coefficients for disc herniation with grade three ( $r = -0.212$ ) and functional disability with grade three ( $r = 0.17$ ) were weak. There was a weak correlation between the intensity of pain and functional disability of these patients ( $r = 0.159$ ). Clinical symptoms and MRI results of the patients who are suffering from cLBP should be correlated to decide whether to perform a therapeutic intervention as the results showed that there was a weak correlation between pain and functional disability based on the degree of disc herniation.

*This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, and build upon the work non commercially.*

**To Cite This Article:** Alshammari QT, Alsubaie AN, Alkayyat AA, Qazaq AM, Alnukhali AA, Alrefaei MG, et al. Magnetic Resonance Imaging of Chronic Low Back Pain: Correlation between Pain, Disability, and Disc Herniation. *Pharmacophore*. 2023;14(3):41-6. <https://doi.org/10.51847/FEgODPJLYS>

**Corresponding Author:** Qurain Turki Alshammari; Department of Diagnostic Radiology, College of Applied Medical Sciences, University of Hail, Hail, Saudi Arabia. E-mail: [g.algrain@uoh.edu.sa](mailto:g.algrain@uoh.edu.sa)

## Introduction

VAS rating of 3 and low grade disc herniation were not correlated based on MRI findings. Disc herniation had a weak Pearson correlation coefficient ( $r = -0.212$ ) while functional disability had a weak Pearson correlation coefficient ( $r = 0.17$ ) with grade three. In this study, the intensity of pain was not strongly correlated with functional disability. Chronic low back pain (cLBP) is a condition that affects roughly eighty percent of human being at some point in their lives. In approximately eighty percent of cases, this pain is caused by structural changes resulting from disc degeneration [1]. Patients with disc herniation (or intervertebral disc prolapse) ( $r = 0.159$ ) suffer from a variety of mechanical factors, including disc degeneration. As a result of weak posture or a spinal injury that occurred as a result of spinal flexion or rotation, particularly during spinal exercise and back extension [2]. Disc degeneration is the most common cause of back pain [3]. In addition to supporting the spine's stability, the neuromuscular system is crucial to its biomechanics [4-6]. Pain and disability may be exacerbated by back weakness and fatigue, as well as a reduced level of spine mobility in chronic low back pain [7, 8]. In order to maintain a healthy lumbar spine, trunk muscles must contract adequately, and balance must be upheld amongst agonist and antagonist muscles, and balance of the buildings surrounding it [8, 9]. Pain can delay the contraction of back muscles, increasing the risk of further spinal injury and lumbar spine dysfunction [9, 10].

It is imperative that trunk muscles contract adequately and that balance is maintained in order to maintain a healthy lumbar spine [11-13]. Furthermore, MRI allows us to see how the intervertebral disc is connected to the peripheral soft tissues and neural anatomical buildings [14, 15]. So, it is used to assess soft tissue pathologies around the spine, as well as spinal cord and spinal canal pathologies [16, 17]. It is possible to explore the size, extent, and location of the disc herniation through magnetic resonance imaging (MRI). Prolapsed discs within the spinal canal determine an IVDP patient's outcome primarily by their size and location [18]. In some patients, a disc herniation of the same size may be symptomatic, whereas in others, it may be asymptomatic [19]. Thus, clinical correlation is extremely important before treatment intervention in cLBP patients in order to minimize misinterpretation of abnormal MRI findings [20]. Therefore this study is aimed at examining the relationship between pain and disability with disc herniation in patients with cLBP with severe disc prolapse (grade 3-disc herniation).

## Materials and Methods

The study studied 70 individuals with pain in the lower back referred to the Radiology Department of King Khalid Hospital, Hail for lower back MRI scans between January and December 2022 after receiving ethical approval. Based on the Michigan State University grading, only grade 3 disc herniations were considered [21]. In the study, both sexes were eligible to participate, provided they were between 20 and 60 years old without cognitive impairment. Prior spinal surgery, trauma, rheumatologic disorders, spine infections, established spine exercise training within three months of the study commencement, spinal fracture, spinal abnormalities [scoliosis, kyphosis], spondylolisthesis, pregnancy, malignancy, congenital abnormalities, ankylosing spondylitis, a hernia, a visceral problem, fibromyalgia, and myofascial pain were considered exclusion criteria. Aortic aneurysms, cerebral and carotid clips, or heart pacemakers were also excluded from the study if they had absolute contraindications to MRI. After providing the patients with an overview of the Visual Analogue Scale, Rolland Morris Disability Questionnaire, and MRI scanning, their consent was obtained voluntarily. A descriptive statistical analysis of the demographic data was performed based on age, sex, occupation, height, and weight. An information booklet containing the instruction about the study, demographic information like name, age, gender, height, weight, duration of the symptom, and the measurement tools were given to the patients and asked them to mark the pain intensity in VAS and the functional disability in RMDQ.

### *Pain*

Visual analogical scales (VAS) are used in this study to assess pain intensity. A pain level of 0 on the left extremity corresponds to a pain level of 10 on the right extremity. It was asked that patients use this scale to indicate the level of pain they are experiencing at the moment. The higher the number, the stronger the pain.


### *Functional Disability*

RMDQ, a functional measure of LBP's impact on daily activities, was used to assess functional disability. The assessment and quantification of LBP can be done using a variety of functional questionnaires. A functional activity was used to determine which statement best described participants' symptoms. By adding up all the checked boxes, we calculated the final score. A score of 0 indicates no disability, an 11 indicates mild disability, an 18 indicates moderate disability, and 24 indicates severe disability [21].

### *Magnetic Resonance Imaging (MRI)*

MRIs were performed on all participants supine using a 1.5 T Siemens Avanto scanner with a 24-element body spine surface coil.

T1 and T2 weighted sagittal and axial MRI sequences were performed with turbo spin echo. The procedure used as follows: T1-sagittal weighted sequences, T1- axial multi stack and angle, T2- sagittal weighted sequences, T2-axial weighted sequences, T2- STIR sagittal, T2-axial multi stack and angle (**Figure 1**).



**Figure 1.** a) An MRI showing L4-L5 lumbar disc herniation as Grade 3 and, b) an MRI showing axial T2 weighted image and, c) sagittal T2

Two musculoskeletal radiologists used standardized evaluation protocols to evaluate and interpret the MRI findings (L1 to S1). MRI is used to assess lumbar disc herniation according to the MSU Classification (**Figure 1**) [16].

#### *Statistical Analysis*

Statistical analysis was performed using SPSS version 23.0 (SPSS Inc., Chicago, IL, USA). Age, duration, height, weight, mean and standard deviation of VAS scores for all participants are displayed. The MRI variables pain intensity [VAS], functional impairment [RMDQ], and grade 3 disc herniation were correlated using the Pearson correlation coefficient.

#### **Results and Discussion**

Based on demographic data such as age, height, weight, and duration of the symptoms, the Mean was calculated (**Table 1**). A total of 70 individuals who have grade-3 disc herniation were included, 35 were males (50%) and 35 were females (50%) The mean age of the patients was  $39 \pm 11.7$  (range: 19-62 years old). There were 37 patients in our study who had level three disc

herniations of the AB type, followed by 22 patients who had level three disc herniations of the B type. Among the 11 subjects [15.7%] with type A herniation, only 11 had it (**Table 2**).

**Table 1.** Demographic data

Variables	Mean $\pm$ SD
Age (years)	39 $\pm$ 11.7
Weight (kg)	78 $\pm$ 7.6
Height (cms)	169.8 $\pm$ 8.17
Duration (days)	92 $\pm$ 7.73

**Table 2.** Distribution of the subjects

MSU [Grade 3]	
A	11 [15.7%]
B	22 [31.4%]
AB	37 [52.8%]

\* MSU – Michigan State University Classification.

It was found that there was no considerable relationship existing between the level of lumbar disc herniation and the sex of the patients ( $p = 0.487$ ). This group scored an average of 8 out of 10 on the VAS, while the average of 16 out of 23 on the RMDQ was determined. Pain intensity calculated with VAS score in the participants with L4-L5 level and L5-S1 level disc herniation was recorded as  $7.2 \pm 1.3$  and  $7.9 \pm 1.19$ , respectively. Pain intensity and functional disability were weakly correlated with grade-3 disc herniation ( $r = -0.212$ ;  $P = 0.17$ ;  $r = 0.197$ ;  $P = 0.49$ ). In participants with severe LBP, pain intensity was also weakly correlated with functional disability ( $r = 0.159$ ;  $P = 0.51$ ) (**Table 3**).

**Table 3.** Correlation between pain, disability, and level of disc prolapse

	'r' Value	Interpretation
<b>Pain &amp; Disability</b>	0.159	Weak Correlation
<b>Pain &amp; MSU</b>	-0.212	Weak Correlation
<b>Disability &amp; MSU</b>	0.197	Weak Correlation

\* MSU – Michigan State University Classification.

Assessment of clinical significance of anatomical abnormalities detected by MRI technology, present study examined the clinical symptoms of grade-3 disc herniation individuals and clinically correlated them with MRI.

The most common musculoskeletal problem is chronic low back pain, and disc herniations are a serious health problem [22]. Low back pain is caused by various causes, but lumbar disc degeneration is the most common, and disc herniation is caused by degeneration of the disc. In lumbar disc degeneration, A nerve root compromise can be graded most accurately with MRI. The current research comprised both males and females, unlike previous studies that only included males [23-25].

The cause of chronic low back pain can be various, including modification in relation to age physical activity, and history [26]. Obese patients had a 5.7-fold greater chance of affection by LBP than normal subjects when compared to those with a normal body mass index [27]. Additionally, lifestyle contributes to the emergence of cLBP. One study reached a conclusion that those who train lightly have a higher risk of developing severe back pain than those who exercise vigorously [27].

We found that demographic and clinical variables, as well as MRI factors, are weakly related to pain intensity and disability in both males and females. The findings of our study were in line with an autopsy study of 647 lumbar spines that revealed disc degeneration is frequent at levels L4-L5 [28]. Pathology usually occurs at L4-L5 but not to the same degree as previously reported [29]. There is a relation amongst pain intensity, functional disability, and cLBP, but the correlation is weak in this study [30]. Possibly because of the young age of the patients, in our study, the mean age was 39 years and 11.7 years. According to literature, pain intensity is a major variable in predicting disability among chronic low back pain patients in a multivariate analysis [31]. A disability score is the most reliable method of measuring the intensity of back pain. Moreover, it helps us understand how acute pain transforms into chronic pain [32] as well as acts as a factor of function like resuming to work [33]. A herniated disc may cause more clinical manifestations and a higher incidence of disability [34]. It was seen that in our study that pain level had a weak correlation with disability, contrary to previous studies which found that disability was correlated with pain level as well as with depression, fatigue, psychosocial factors, financial status, and unemployment [35].

Patients suffering from chronic low back pain can benefit from MRI imaging of the lumbar spine, which is done in an upright position in order to detect disc degeneration [36]. Around 30% of disc abnormalities were found in MRI studies of asymptomatic subjects [37] and the incidence of LBP in these subjects are high [38]. Degenerative findings in spine images and their clinical significance have been the subject of many studies in the literature [39]. The long-term relationship between degenerative changes in the spine and disability has also been studied but in limited numbers [40]. It has been found that

protrusion of the discs and bulges are highest in the 5th and 6th decade [41], as we also found in our study. A major limitation of the current research was the little sample size and also it was done in one base. In addition, this study did not assess BMI and psychosocial factors like depressed state, panic, economical level, smoking habits, etc. This study didn't include subjects who were asymptomatic (control group).

### Conclusion

The current study indicates that grade-3 disc herniations in the lumbar spine do not correlate with pain intensity or functional disability among subjects with severe chronic low back pain. As a result, we encourage clinicians to extend the correlation between disc herniation on MRI and clinical symptoms in individuals who suffer from pain from the lower back.

**Acknowledgments:** The authors take the opportunity to thank all the participants of this study and the MRI Technicians and the nurse who helped in taking the MRI for the patients.

**Conflict of interest:** None

**Financial support:** None

**Ethics statement:** None

### References

1. Abdullah IB, Al-Mutairi MM, Alghubayan MA, Alamir AA, Bu-Jubarah AY, Almajed N, et al. Literature review on prevalence, risk factors and, evaluation of acute lower back pain. *Int J Pharm Res Allied Sci.* 2020;9(1):35-40.
2. Adams MA, Freeman BJ, Morrison HP, Nelson IW, Dolan P. Mechanical initiation of intervertebral disc degeneration. *Spine.* 2000;25(13):1625-36.
3. Ammar A, Alwadei A, Al Hayek A, Alabbas FM, Almatrafi FR, Elshawarby M. The correlation between histopathology of herniated lumbar intervertebral disc and clinical findings. *Asian J Neurosurg.* 2020;15(3):545-53.
4. Joyce C, Roseen EJ, Keysor JJ, Gross KD, Culpepper L, Saper RB. Can Yoga or Physical Therapy for Chronic Low Back Pain Improve Depression and Anxiety Among Adults From a Racially Diverse, Low-Income Community? A Secondary Analysis of a Randomized Controlled Trial. *Arch Phys Med Rehabil.* 2021;102(6):1049-58.
5. Lamichhane B, Jayasekera D, Jakes R, Ray WZ, Leuthardt EC, Hawasli AH. Functional Disruptions of the Brain in Low Back Pain: A Potential Imaging Biomarker of Functional Disability. *Front Neurol.* 2021;12:669076.
6. Cerillo JL, Becsey AN, Sanghadia CP, Root KT, Lucke-Wold B. Spine Bracing: When to Utilize-A Narrative Review. *Biomechanics (Basel).* 2023;3(1):136-54.
7. Mandroukas A, Michailidis Y, Kyranoudis AE, Christoulas K, Metaxas T. Surface Electromyographic Activity of the Rectus Abdominis and External Oblique during Isometric and Dynamic Exercises. *J Funct Morphol Kinesiol.* 2022;7(3):67.
8. Horii O, Sasaki M. Influences of trunk stability on exercise performance of closed kinetic chain of upper and lower limbs. *J Phys Ther Sci.* 2023;35(5):379-83.
9. Lee K. Motion Analysis of Core Stabilization Exercise in Women: Kinematics and Electromyographic Analysis. *Sports (Basel).* 2023;11(3):66.
10. Vasilievich PV. Determination of Stresses in the Apparatus of External Fixation of the Human Spine. *Int J Pharm Res Allied Sci.* 2021;10(2):62-9.
11. Othman M, Menon VK. The prevalence of Schmorl's nodes in osteoporotic vs normal patients: a Middle Eastern population study. *Osteoporos Int.* 2022;33(7):1493-9.
12. Ohnmeiss DD, Gatchel RJ. North American Spine Society compendium of outcome instruments for assessment and research of spinal disorders. North American Spine Society, La Grange; 2001.
13. Byvaltsev VA, Kalinin AA, Shepelev VV, Pestryakov YY, Aliyev MA, Hozeev DV, et al. Prevalence of lumbosacral transitional vertebra among 4816 consecutive patients with low back pain: A computed tomography, magnetic resonance imaging, and plain radiographic study with novel classification schema. *J Craniovertebr Junction Spine.* 2023;14(1):35-43.
14. Li Y, Fredrickson V, Resnick DK. How should we grade lumbar disc herniation and nerve root compression? a systematic review. *Clin Orthop Relat Res.* 2015;473(6):1896-902.
15. Shevchenko YS, Plohova DP, Bulakhova IN, Mishvelov AE, Kubalova ME, Badriev GB, et al. Experience of carrying out magnetic resonance imaging with the use of specialized protocols and programs computer post-processing. *Pharmacophore.* 2020;11(2):77-81.
16. Lund T, Schlenzka D, Lohman M, Ristolainen L, Kautiainen H, Klemetti E, et al. The intervertebral disc during growth: Signal intensity changes on magnetic resonance imaging and their relevance to low back pain. *PLoS One.* 2022;17(10):e0275315.

17. Waseem S, Kyriakides J, Amiri AR, Shetty R, Shetty N, Chammaa R. Management strategies for the painless foot drop: a systematic review of the literature. *Eur Spine J.* 2023;32(4):1099-105.
18. Thapa SS, Lakhey RB, Sharma P, Pokhrel RK. Correlation between Clinical Features and Magnetic Resonance Imaging Findings in Lumbar Disc Prolapse. *J Nepal Health Res Counc.* 2016;14(33):85-8.
19. Hu S, Li Y, Hou B, Zhang Y, Liu WV, Wu G, et al. Multi-echo in steady-state acquisition improves MRI image quality and lumbosacral radiculopathy diagnosis efficacy compared with T2 fast spin-echo sequence. *Neuroradiology.* 2023;65(5):969-77.
20. Hamaguchi H, Kitagawa M, Sakamoto D, Katscher U, Sudo H, Yamada K, et al. Quantitative Assessment of Intervertebral Disc Composition by MRI: Sensitivity to Diurnal Variation. *Tomography.* 2023;9(3):1029-40.
21. Mysliwiec LW, Cholewicki J, Winkelpleck MD, Eis GP. MSU classification for herniated lumbar discs on MRI: toward developing objective criteria for surgical selection. *Eur Spine J.* 2010;19:1087-93.
22. Zahra N, Sheha EA, Elsayed H. Low back pain, disability and quality of life among health care workers. *Int J Pharm Res Allied Sci.* 2020;9(2):34-44.
23. Skovlund SV, Bláfoss R, Skals S, Jakobsen MD, Andersen LL. Technical field measurements of muscular workload during stocking activities in supermarkets: cross-sectional study. *Sci Rep.* 2022;12(1):934.
24. Buruck G, Tomaschek A, Wendsche J, Ochsmann E, Dörfel D. Psychosocial areas of worklife and chronic low back pain: a systematic review and meta-analysis. *BMC Musculoskelet Disord.* 2019;20(1):480.
25. Chung JW, So HC, Yan VC, Kwok PS, Wong BY, Yang JY, et al. A survey of work-related pain prevalence among construction workers in Hong Kong: A case-control study. *Int J Environ Res Public Health.* 2019;16(8):1404.
26. Videman T, Levalahti E, Battie MC. The effects of anthropometrics, lifting strength, and physical activities in disc degeneration. *Spine.* 2007;32(13):1406-13.
27. Ordoñez-Hinojos A, Durán-Hernández S, Hernández-López JL, Castillejos-López M. Association between strenuous occupational activity and low back pain. *Acta Ortop Mex.* 2012;26(1):21-9.
28. Kuisma M, Karppinen J, Niinimäki J, Ojala R, Haapea M, Heliövaara M, et al. Modic changes in endplates of lumbar vertebral bodies: Prevalence and association with low back and sciatic pain among middle-aged male workers. *Spine.* 2007;32(10):1116-22.
29. Deyo RA, Loeser JD, Bigos SJ. Herniated lumbar intervertebral disk. *Ann Intern Med.* 1990;112(8):598-603.
30. Taghizadeh ME, Oraki M, Badakhshan M. The effect of cognitive-behavioral hypnotherapy on modifying the pain beliefs and pain self-efficacy in patients with Chronic Low Back Pain (CLBP). *Int J Pharm Phytopharmacol Res.* 2019;9(3):128-33.
31. Mannion AF, Junge A, Taimela S, Muntener M, Lorenzo K, Dvorak J. Active therapy for chronic low back pain. Part 3. Factors influencing self-rated disability and its change following therapy. *Spine.* 2001;26(8):920-9.
32. Pincus T, Burton AK, Vogel S, Field AP. A systemic review of psychological factors as predictors of chronicity/disability in prospective cohorts of low back pain. *Spine.* 2002;27(5):109-20.
33. Nordin M, Skovron ML, Hiebert R, Weiser S, Brisson PM, Campello M, et al. Early predictors of delayed return to work in patients with low back pain. *J Musculoskel Pain.* 1997;5(2):5-27.
34. Dora C, Schmid M, Elfering A, Zanetti M, Hodler J, Boos N. Lumbar Disk Herniation: Do MR Imaging Findings Predict Recurrence after Surgical Discectomy? *Radiology.* 2005;235(2):562-7.
35. Sterud T, Tynes T. Work-related psychosocial and mechanical risk factors for low back pain: A 3-year follow-up study of the general working population in Norway. *Occup Environ Med.* 2013;70(5):296-302.
36. Tarantino U, Fanucci E, Iundusi R, Celi M, Altobelli S, Gasbarra E, et al. Lumbar spine MRI in upright position for diagnosing acute and chronic low back pain: Statistical analysis of morphological changes. *J Orthop Traumatol.* 2013;14(1):15-22.
37. Yang J, Xiong Y, Hu Y, Huang M, Zhang L, Pu X, et al. The reliability, correlation with clinical symptoms and surgical outcomes of dural sac cross-sectional area, nerve root sedimentation sign and morphological grade for lumbar spinal stenosis. *BMC Musculoskelet Disord.* 2023;24(1):225.
38. Kim P, Ju CI, Kim HS, Kim SW. Lumbar Disc Herniation Presented with Contralateral Symptoms. *J Korean Neurosurg Soc.* 2017;60(2):220-4.
39. El Barzouhi A, Vleggeert-Lankamp CL, van der Kallen BF, à Nijeholt GJ, van den Hout WB, Koes BW, et al. Back pain's association with vertebral end-plate signal changes in sciatica. *Spine J.* 2014;14(2):225-33.
40. Romero-Munoz LM, Barriga-Martin A, Segura-Fragoso A, Martin-Gonzalez C. Are Modic changes in patients with chronic low back pain indicative of a worse clinical course? 10 years of follow-up. *Rev Esp Cir Ortop Traumatol.* 2018;62(4):274-81.
41. Omoigui S, Fadare A. A New Minimally Invasive Procedure for Muscle, Back, Neck Pain and Radiculopathy - The Myofascial Nerve Block. *Ann Clin Case Rep.* 2022;7(1):2233.