



ON OVERVIEW OF BIOACTIVE COMPOUNDS, BIOLOGICAL AND PHARMACOLOGICAL EFFECTS OF MISTLETOE (*VISCUM ALBUM L*)

Eva Kleszken^{1#}, Adrian Vasile Timar², Adriana Ramona Memete^{1#}, Florina Miere (Groza)³, Simona Ioana Vicas^{2*}

1. *Doctoral School of Biomedical Science, University of Oradea, Oradea, Romania.*
2. *Department of Food Engineering, University of Oradea, Oradea, Romania.*
3. *Faculty of Medicine and Pharmacy, University of Oradea, Oradea, Romania.*

#These Authors Contributed equally to this work

ARTICLE INFO

Received:

17 Oct 2021

Received in revised form:

12 Jan 2022

Accepted:

17 Jan 2022

Available online:

28 Feb 2022

Keywords: *Viscum album*, Host trees, Flavonoids, Phenolic acid, Antioxidant, Anticancer

ABSTRACT

The genus *Viscum* includes many species that are mainly distributed in Europe, Africa, Asia, America, and Australia. *Viscum* extracts or their various preparations are widely used as complementary and alternative medicines in the treatment of various ailments. In the present review, articles related to the phytochemical composition of mistletoe were selected, depending on the host tree on which it grows, as well as articles in which its beneficial effects were highlighted. *Viscum* contains different active ingredients, including lectins, viscotoxins along with phenolic acids, flavonoids, alkaloids, terpenoids, and polysaccharides. Based on its composition, mistletoe extract is associated with multiple bioactivities, including anticancer, anti-inflammatory, and cardiovascular disease, attenuating the side effects of chemotherapy and enhancing immunity. The purpose of this review was to highlight the link between the host tree and the bioactive components of mistletoe such as lectin and viscotoxin, with a focus on phenolic compounds such as flavonoids and phenolic acids. The potential therapeutic effects of mistletoe are summarized by subspecies and host trees. Numerous mistletoe-based patents with various applications have been developed and presented in this review. Mistletoe is a medicinal plant with great biological potential that is worth exploring for various targeted treatments.

This is an **open-access** article distributed under the terms of the [Creative Commons Attribution-Non Commercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by/4.0/), which allows others to remix, tweak, and build upon the work non commercially, as long as the author is credited and the new creations are licensed under the identical terms.

To Cite This Article: Kleszken E, Timar AV, Memete AR, Miere(Groza) F, Vicas SI. On Overview of Bioactive Compounds, Biological and Pharmacological Effects of Mistletoe (*Viscum Album L*). *Pharmacophore*. 2022;13(1):10-26. <https://doi.org/10.51847/Tmo2sXGQRs>

Introduction

Parasitic angiosperms are over 4,700 species of 277 genera, of which 87 genera and at least 1,670 species belong to the mistletoe which is considered hemiparasites that grow on the host branches of various species of shrubs and trees looking like a highly developed shrub [1, 2]. Living organisms are involved in parasitic relationships, either as parasites or as hosts, these interactions having an essential role in the proper functioning of the biosphere and the process of biological evolution [3, 4]. Mistletoe (*Viscum album L.*) is an evergreen plant that depends on the host tree for some nutrients and water, while it produces carbohydrates in a process of photosynthesis, normally found growing on a variety of trees, both deciduous and coniferous trees [2]. Mistletoe (*V. album*), is a large hemiparasitic species found in northern, central, and southern Europe, Africa, America, Australia, and Asia, accompanied by a wide variety of habits, host preferences, morphologies, and development patterns depending on the geographical distribution [5-7].

Several chemical and pharmacological studies have identified various types of compounds in mistletoe, such as lectins, viscotoxins, lignans, amines, flavonoids, and polysaccharides [8, 9]. Among them, flavonoids and phenolic acids are natural antioxidants involved in the biological activity of the plant and used in the prevention of diseases such as cancer, caused by oxidative stress induced by free radicals in the body [10-13]. Cancer therapy is one of the most important uses of mistletoe, proving to have beneficial effects on various types of cancer such as breast [14, 15], pancreatic [16], laryngeal [17] bladder [18], or leukemia [19]. Over time, other bioactive activities of mistletoe have been reported, such as those related to neurological disorders, antiviral, antibacterial, anti-inflammatory, antiepileptic, or immunostimulatory activities [2, 11, 13,

Corresponding Author: Simona Ioana Vicas; Department of Food Engineering, University of Oradea, Oradea, Romania. E-mail: svicas@uoradea.ro.

20].

Due to its bioactive compounds and its beneficial effects on the human body, mistletoe has become more and more studied, developing various pharmaceutical formulations (Iscador, Isorel, Iscucin, Lektinol, Eurixor, Helixor, Abnoba-Viscum and recombinant lectin ML -1) [13, 21].

This review aimed to highlight the link between the host tree and the bioactive components of mistletoe such as lectin and viscotoxin, with a focus on phenolic compounds such as flavonoids and phenolic acids. The potential therapeutic effects of mistletoe are summarized by subspecies and host tree. Numerous mistletoe-based patents with various applications have been developed and presented in this review.

Materials and Methods

This review was performed using the PRISMA 2020 flow chart based on the suggestion of Page *et al.*, 2021 [22]. The steps and selection criteria, followed by the number of studies used for our review, are shown in **Figure 1**. Databases such as PubMed, Scopus, Science Direct, Elsevier, Google Scholar, Google Patents have been accessed to search the literature. The keywords of the medical subject titles included in the search were: "*Viscum album*", "*V. coloratum*", "*V. articulatum*", "mistletoe viscotoxin", "mistletoe lectin", "mistletoe effects", "mistletoe patents", "mistletoe compounds", "mistletoe history", etc. All the information systematized in the tables was obtained from research articles. Studies published in languages other than English and Romanian were excluded. A total of 100 studies were selected and included in this review.




Figure 1. PRISMA 2020 flow diagram for the present review

The Historical Approach of Mistletoe

Given the co-evolutionary history between plants and the organisms that use them, over 150 years ago, the co-adaptation between mistletoe and birds caught the attention of Charles Darwin, stimulating considerable scientific research. Equally important for other researchers was the understanding of the interactions between herbivorous insects and plants, which can make the transition between plant taxa. The mistletoe is also considered the ancestral host of butterflies [23-25].

The number of host species infected with mistletoe is critical in that it influences the overall distribution of the parasite, its prevalence, and virulence. Therefore, macroecological analyzes of this feature related to the history of mistletoe life are missing for many regions, and with the evolution of mankind, geographical and environmental changes have influenced the parasitic species, through hybridization processes [1, 26, 27].

Mistletoe was later discovered for its miraculous effects, and traditional medicine spread it over time to many parts of the world, in particular, *V. album* L. and *Loranthus (Taxillus Chinensis)* (DC.) Danser [28, 29]. The latter belongs to the family *Loranthaceae*, distributed mainly in southern and southwestern China, is known as "Sang Ji Sheng", and its leaves and stems have a long history in traditional Chinese medicine, is used to treat rheumatism, arrhythmia, angina pectoris, hypertension,

stroke [28]. *V. album* L. also has a long traditional history of about 100 years, used in German-speaking countries in cancer treatment therapy [29].

In addition to the fact that mistletoe was used for medicinal purposes, there are also several legends gathered from different regions of the world, related to this plant, legends that our ancestors respected and mentioned and which are still remembered today.

Because of its beliefs to bring good luck, to heal illnesses, and to attract angels, mistletoe is known since old times. It is said to bring health and prosperity. It had a very important part in the Celtic rituals. They believed that mistletoe was created by a bolt of lightning that hit a tree giving him magical powers. The Celts believed that a very precious mistletoe was grown by the oak, also known as the „Oak Tear”, and poaching this kind of mistletoe was always a moment of ceremonial cutting. Thus the mistletoe was cut only after the winter solstice with the Golden Sickle, by a white dressed priest. Mistletoe was also considered the symbol of femininity (*Naturalis Historia*, books XVI, XXIV, XXXIII).

The Greeks considered mistletoe from the underworld, believing that a branch of mistletoe was the key Aeneas that opened the underworld. The Romans put on Goddess Diana's head a crown of mistletoe, as a symbol of fertility. In the Scandinavian legends, mistletoe was something very special symbolizing peace, love, harmony, spiritual cleansing, and throwing away bad spirits.

In European mythology, mistletoe has a unique place. Around Christmas and New Year's Eve, people put mistletoe twigs on doors and windows to bring luck and health. Other legends talk about a peculiar custom that if two enemies were met under a tree that grew mistletoe, they come to terms on behalf of the mistletoe. Later this custom was taken by the lovers, who kissing under the mistletoe felt like a love pledge, bringing happiness and fulfillment. In the town Tenbury Wells, from the UK, a mistletoe festival is held every year and a mistletoe Queen is chosen, bearing a crown of mistletoe [30].

Botanical Characterization

V. album is a hemiparasite shrub species that grows on branches and twigs of trees. It is a plant with slow development, usually being a nod to the blooming season. Every year mistletoe develops more and more branches, so its growth tells us its age. Mistletoe can live up to 70 years [31, 32] It looks like a very big bush, having a diameter of up to one meter (**Figure 2a**). The mistletoe is stuck on the host tree. It looks like a vertical, well-developed ax that enters the bark of the host tree, pervading its woody tissue. This network continues to grow, creating a root system (haustorium) that absorbs water, sugars, amino acids, and minerals from host trees, being able to biosynthesize their own primary and secondary metabolites [33]. Its stem is cylindrical, thick, dichotomic ramification, wide off 30 - 60 cm, thicker towards the nodes, where it can be torn easily, having a yellow-greenish color. Leaves are green all year round, have no stalk, full edges, being seen like a ribbon, thick, skinny, bearing no wintergreen shrubs, having a yellow to intense green color, and are set in opposite pairs. On its lower part, one can notice 4 - 5 veins (**Figure 2b**). The size of the mistletoe leaves varies according to the host tree and the harvesting time.

The leaves harvested in April are larger, especially in the case of mistletoe leaves hosted by acacia. Leaves have a specific smell and a soft bitter sour taste. The optimal period to harvest is November to April, as the flowers are small (2-3 mm in diameter), yellow-greenish color, dioice, can be seen on top of young twigs. It blooms in March- April (**Figure 2c**). The fruits are spherical, having a glassy look, with a diameter of approximately 8 mm, green at the base. Between September - October, (**Figure 2d**) the fruits are white translucent, while in November- March they are toxic (**Figure 2f**). As a rule, the fruits are gathered thrisome during September-December (**Figure 2e**), containing 1-2 seeds covered by a jelly, sticky liquid (**Figure 2g**).

Up to the 18th-century botanists confused white mistletoe (*V. album*) with oak mistletoe (*L. europaeus*). The *V. album* belongs to the Viscaceae family, while *L. europaeus* belongs to the Loranthaceae family. The link between them is of taxonomic order and both belong to the Santalales order [7, 34].

The *L. europaeus* has younger twigs, brownish color, lance-like leaves, short stark and whole edge. It loses its leaves during wintertime, while white mistletoe keeps its leaves the entire year-round, being always green. The last leaves are small, all yellow-greenish, like a spike shape, blooming around May - June. The fruits of *V. album* are small, yellowish, in the form of a small cluster shape. Its fruit is ripe in October, usually has a seed, that is spread by birds [35].



a



b



c




Figure 2. Morphology of *V. album*. a) Mistletoe (*V. album* on acacia), b) leaves of mistletoe, c) Flower of mistletoe, d) Unripe fruits of mistletoe, e) Unripe fruits with seeds, f) White fruits of mistletoe, g) White fruits with seeds

Taxonomy

The Viscaceae family has seven genera: *Viscum*, *Phoradendron*, *Notothixos*, *Korthalsella*, *Ginalloa*, *Dendrophthora*, and *Arceuthobium*. Regarding the scientific rating the mistletoe (*V. album* L) belongs to the *Viscum* genus, part of the Viscaceae family, but genetic research, new approach in APG (Angiosperm Phylogeny Group) III system, show that mistletoe belongs to the Santalaceae family. The genus *Viscum* comprises about 90 species, of which two-thirds are native to Africa and one-third to Eurasia and Africa [36].

The *Phoradendron* mistletoe is set in a part of Northern America (the North American Oak mistletoe, *Phoradendron serotinum*) and South America, Argentina. *Loranthus europaeus* belongs to the family Loranthaceae, which is widespread in America. Genera *Notothixos* mistletoe is found in the East of Australia and the Island. The *Korthalsella* kind is seen in East Asia, Malaysia, Australia, New Zealand. The genus *Ginalloa* includes only a few species that can be found in the South East of Asia. The mistletoe belonging to the genus *Dendrophthora* has a distribution similar to that of the genus *Phoradendron*, with an area in Central America. Genera *Arceuthobium* has approximately 24 species, being found in the USA and Mexico [37].

Host Trees and Geographical Area

The European mistletoe has several accepted subspecies that can be easily differentiated according to the host tree, as they are morphologically very similar. Mistletoe species differ in leaves' shape and size, their fruits color, and host tree. *V. album* species parasitize over 400 host trees [38]. **Table 1** summarizes literature data on mistletoe species/subspecies, their host trees on different continents, leaf size, and fruit color.

Table 1. Mistletoe species from different continents, their host trees, leaf size, and fruit color

Mistletoe species	Spread	Host tree	Leaves	Fruits	References
Europe					
<i>V. album ssp. abietis</i>	Central Europe	Fir (<i>Abies alba</i>)	Up to 8 cm	whites	[7]
<i>V. album ssp. album</i>	Europe, South - vest Asia	Apple(<i>Malus</i>), Linden (<i>Tilia</i>), Willow (<i>Salix vitellina</i>), Poplar (<i>Populus nigra</i>), Oak (<i>Quercus</i>) Hawthorn (<i>Crataegus monogyna</i>), Apricot (<i>Prunus armeniaca</i>), Acacia (<i>Robinia pseudoacacia</i>) It never appears on beech.	Up to 5 cm	whites	[7]
<i>V. album ssp. austriacum</i>	Central Europe, rarely in Greece	Coniferous (<i>Larix</i> , <i>Pinus</i> , <i>Picea</i>).	4-6 cm	yellow	[7]

<i>V. album ssp. Creticum</i>	Eastern Crete	Pine (<i>Pinus</i>)	small	whites	[5]
Asia					
<i>V. album ssp. meridianum</i>	Southeast Asia	Hornbeam (<i>Carpinus monbeigiana</i>), Ash (<i>Fraxinus</i>), Maple (<i>Acer</i>), Walnut (<i>Juglans regia</i>), Plum (<i>Prunus pseudocerasus</i>) Mountain ash (<i>Sorbus megalopa</i>)	3-5 cm	yellow	[39]
<i>V. album ssp. coloratum</i>	China	Oak (<i>Quercus sp.</i>)	2 – 4 cm	red	[40-42]
Africa					
The red mistletoe (<i>V. cruciate</i>).	North Africa (Morocco, Libya) end in South Africa, Rarely in South-western Spain, southern Portugal	Olive (<i>Olea europaea</i>)	small	red	[43]
<i>L. ferrugineus</i>	Africa	Oak (<i>Quercus sp.</i>), Acacia (<i>Robinia pseudoacacia</i>) Euphorbia	4-8 cm	yellow	[44]
<i>L. micranthus</i>	Nigeria	Oak (<i>Quercus</i>),	4-6 cm	yellow	[45]
Australia					
<i>Amyema maidenii ssp. maidenii</i>		Eucalyptus	flat	yellow	[46]
<i>A. gibberula ssp. gibberula</i>			Long and cylindrical	whites	[46]
<i>A. bifurcata ssp. bifurcata</i>	Australia		flat	red	[47]
<i>Lysiana exocarpi ssp. exocarpi</i>		Acacia	3-15 cm	red or black	[48]
<i>L. murrayi</i>			2.5 – 6cm	pink or red	[46]
<i>L. spathulata</i>			3-7 cm	red	[47]
<i>L. subfalcata</i>			2-12 cm	yellowish	[46]
America					
The American Mistletoe (<i>Phoradendron californicum</i>) and the dwarf mistletoe (<i>Arceuthobium minutissimum</i>)	North, America, California, South America, Argentina	Oak (<i>Quercus sp.</i>), Pine (<i>Pinus</i>)	4-8 cm	red	[49]

Mistletoe can be seen in forests, on hills, and in the mountains, on various host trees, such as birch, ash, maple, poplar, lime, willow, or on conifers: fir, pine. The mistletoe also parasitizes fruit trees such as apples, pears, plums, cherries. The genus *Viscum* included species seen in the temperate areas of Europe, in Asia, in the tropical and subtropical parts of Africa, in Madagascar, in Australia, and the north and south of America

Mistletoe Bioactive Compounds

The mistletoe species (*V. album* L) has presented and still has a high interest in researching the bioactivity of its compounds. Many types of metabolites have been isolated from European mistletoe, some of which are not synthesized by mistletoe, but are obtained from the host tree, for example, some alkaloids [50].

Jäger *et al.*, 2021 identified the specific biomarkers (arginine, pipercolic acid or lysine, dimethoxycoumarin, and sinapyl alcohol) of *V. album ssp. album* grew on three different host trees (*M. Domestica*, *Q. Robur*, and *U. carpinifolia*) [33].

The metabolite profile of *V. coloratum* harvested from three different host plants (*Ulmus pumila* L., *Salix babylonica*, and *Populus ussuriensis* Kom.) in two habitats (temperate continental climate and warm temperate humid monsoon climate, China) was investigated by Zhang *et al.*, 2021 [51]. Three main metabolites, from the flavanone class, were identified, their synthesis and accumulation in mistletoe were dependent both on the host plant and environment [51].

Polyphenols - Chemical Structure and Classification

The flavonoids are a class of polyphenol compounds, secondary metabolites of vegetal origin, with antioxidant properties, which are found in the free state or seen as esters or glycosides in mistletoe leaves. More than 4.000 types of flavonoids have

been identified in the plant kingdom [52].

Polyphenols occupy an important place in the life of plants by intervening in their metabolism. There are known about 8.000 plant-derived polyphenol compounds with antioxidant and antitumor properties [53-58]. Polyphenols are organic, water-soluble compounds that contain in their molecule one or more aromatic rings with one or more hydroxyl groups. Due to their chemical structural characteristics, polyphenols are compounds with high antioxidant power [52, 59-61]. The main polyphenols identified in mistletoe are from phenolic acids class, both hydroxybenzoic acid (gallic acid, protocatechuic acid) and hydroxycinnamic acid (caffeic, ferulic acid, synaptic acid) and flavonoids including flavanone (Naringenin, Eriodictiol), flavone (Apigenin), flavonol (3-O-Met Quercetin, Myricetin, Kampherol). The main polyphenols identified in the different subspecies of *V. album* and its host trees are presented in **Figure 3**.




Figure 3. The classification and chemical structure of the main polyphenols identified in *V. album*

The amount and composition of mistletoe flavonoids can vary significantly depending on the host tree, vegetation period, and harvest period [33, 41, 51, 62, 63].

The most important flavonoids identified so far in mistletoe leaves or stems are shown in **Table 2**.

Table 2. Polyphenols identified in different subspecies mistletoe leaves or stems depending on the host tree

Species	Host trees	The part used	Flavonoids	Phenolic acids	References
<i>V. album</i> ssp <i>album</i>	Maple (<i>Acer platanoides</i> L)	leaves	Flavone: Apigenin Flavonol: 3-O-Metil Quercetin Flavanones: Naringenin	Hydroxybenzoic acid: Gallic acid, Gentisic acid, Protocatechuic acid, p-hydrobenzoic, Salicylic acid,	[64]
				Cinnamic acid: Caffeic acid, Ferulic acid, p-cumaric acid, Rosmarinic acid, Sinapic acid	[65]
<i>V. album</i> ssp <i>album</i>	Jugastru (<i>Acer campestre</i>)	leaves	Flavonol: Kampherol Quercetin	Hydroxybenzoic acid: Salicylic acid, p-Hydrobenzoic acid,	[64]
				Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid, Ferulic acid, Sinapic acid, trans-Cinnamic acid.	[65]
				Hydroxybenzoic acid: p-Hydrobenzoic acid Salicylic acid,	[65]

				Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid, Ferulic acid, Sinapic acid, trans-Cinnamic acid,	
		stems	Flavonol: Kampherol	Hydroxycinnamic acid: Rosmarinic acid, trans-Cinnamic acid	[65]
			Flavanol: Myricetin Flavonol: 3-O-Metil Quercetin Rhamnazin		[12]
<i>V. album ssp album</i>	Silver coat(<i>Acer saccharinum</i> L)	leaves		Hydroxibenzoic acid: Gallic acid, Gentisic acid, p-Hydrobenzoic acid, Salicylic acid, Vanilic acid Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid, Sinapic acid, Rosmarinic acid,	[64]
<i>V. album ssp album</i>	Hawthorn (<i>Crataegus monogyna</i>)	leaves	Flavanones: Eriodictyol Flavonol: 3-O-Metil Quercetin	Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid Syringic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid p-Coumaric acid, Sinapic acid,	[12]
<i>V. album ssp album</i>	Ash (<i>Fraxinus excelsior</i> L)	leaves	Flavonol: Kampherol Quercetin 3-O-Metil Quercetin Flavanones: Eriodictyol	Hydroxibenzoic acid: Protocatechuic acid, p-Hydrobenzoic acid, Salicylic acid, Syringic acid Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid Ferulic acid, p-Coumaric acid Sinapic acid, trans-Cinnamic acid,	[65]
		stems	Flavonol: Kampherol	Hydroxibenzoic acid: Syringic acid Protocatechuic acid Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid, Ferulic acid, Rosmarinic acid, Sinapic acid, trans-Cinnamic acid,	[65]
<i>V. album ssp album</i>	Ash (<i>Fraxinus pennsylvanica</i> Marsh)	leaves	Flavanones: Eriodictyol Naringenin Flavonol: 3-O-Metil Quercetin Sakuratin Rhamnazin	Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid, Syringic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, Sinapic acid	[64]

			Flavanones: Eriodictyol Naringenin Flavonol: Quercetin 3-O-Metil Quercetin Sakuratin Rhamnazin	[12]
			Flavonol: Quercetin 3-O-Metil Quercetin Myricetin, Sakuratin Isorhamnetin Rhamnetin Rhamnazin Flavanones: Naringenin Eriodictyol	[64]
			Flavonol: Quercetin	[65]
			Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Sakuratin Rhamnazin Flavanones: Naringenin	[12]
<i>V. album</i> ssp <i>album</i>	Apple (<i>Malus Domestica</i> Borkh)	leaves	Hydroxybenzoic acid: Gallic acid Vanilic acid, Gentisic acid, Protocatechuic acid, Veratric acid	[64]
			Hydroxycinnamic acid: p-Coumaric acid	
			Hydroxybenzoic acid: p-hydroxybenzoic acid, Protocatechuic acid, Salicylic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, Sinapic acid, Rosmarinic acid,	[65]
<i>V. album</i> ssp <i>album</i>	Apple (<i>Malus Domestica</i> Borkh)	stems	Hydroxybenzoic acid: Syringic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, Rosmarinic acid Sinapic acid,	[65]
<i>V. album</i> ssp <i>album</i>	Poplar (<i>Populus nigra</i> L)	leaves	Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Sakuratin, Rhamnazin Flavanones: Naringenin Flavone: Apigenin	[65]
			Hydroxybenzoic acid: Gallic acid, Gentisic acid, Protocatechuic acid, Hydroxycinnamic acid: p-Coumaric acid	[64]
			Hydroxybenzoic acid: Salicylic acid, Protocatechuic acid, p-hydrobenzoic acid, Hydroxycinnamic acid: Ferulic acid,	[65]
			Flavonol: Quercetin	

				Rosmarinic acid, Sinapic acid,	
			Flavanones: Naringenin Eriodictyol Sakuranetin Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Rhamnazin	Hydroxibenzoic acid: p-hydrobenzoic acid, Salicylic acid, Syringic acid Vanillic acid	[12]
<i>V. album</i> ssp <i>album</i>	Poplar (<i>Populus nigra</i> L.)	stems		Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid Hydroxycinnamic acid: Caffeic acid, Ferulic acid,	[65]
<i>V. album</i> ssp <i>album</i>	Acacia (<i>Robinia pseudoacacia</i>)	leaves	Flavonol: Quercetin Kaempferol	Hydroxibenzoic acid: Gallic acid, Hydroxycinnamic acid: Ferulic acid, Sinapic acid,	[65]
		stems	Flavonol: Kampherol	Hydroxibenzoic acid: Protocatechuic acid,	[65]
			Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Myricetin, Sakuratin Rhamnazin Flavanones: Naringenin		[12]
<i>V. album</i> ssp <i>album</i>	Mountain ash (<i>Sorbus aucuparia</i> L.)	leaves		Hydroxibenzoic acid: Gallic acid, Gentisic acid Protocatechuic acid, p-hydroxybenzoic acid Salicylic acid, Syringic acid Vanillic acid, Veratric acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid Sinapic acid,	[64]
			Flavanones: Naringenin Eriodictyol Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Sakuratin Rhamnazin		[12]
<i>V. album</i> ssp <i>album</i>	Linden (<i>Tilia cordata</i> Mill)	leaves	Flavanones: Naringenin Eriodictyol Flavonol: 3-O-Metil Quercetin Isorhamnetin Rhamnazin Flavones: Luteolin	Hydroxibenzoic acid: Gallic acid, Protocatechuic acid, p-hydrobenzoic acid, Salicylic acid Syringic acid, Vanillic acid Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid, Sinapic acid.	[12]

<i>V. album</i> ssp <i>abietis</i>	Fir (<i>Abietis alba</i> Mill)	leaves	Flavanones: Naringenin Flavonol: 3-O-Metil Quercetin Rhamnazin Rhamnetin Flavone: Apigenin	Hydroxibenzoic acid: Salicylic acid, Protocatechuic acid, 4-hydrobenzoic acid, Vanilic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid, Sinapic acid.	[66]
<i>V. album</i> ssp <i>austriacum</i>	Pine (<i>Pinus sylvestris</i> L)	leaves	Flavanones: Naringenin Eriodictyol Sakuranetin Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Myricetin, Rhamnazin Rhamnetin, Kampherol		[12]
		leaves fruits seeds		Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid, 4-hydroxybenzoic acid, Vanilic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid, Sinapic acid	[67]
<i>V. album</i> ssp <i>austriacum</i>	Guava(<i>Psidium guajava</i>)	leaves	Flavanones: Naringenin		[68]
<i>V. album</i> ssp <i>coloratum</i>	Oak (<i>Quercus crispula</i>)	leaves stems	Flavanones: Eriodictyol		[19]
<i>V. album</i> ssp <i>coloratum</i>	Elm (<i>Ulmus pumila</i> L) Willow (<i>Salix babylonica</i> L) Poplar (<i>Populus ussuriensis</i> Kom)	leaves	Flavanones: Eriodictyol		[51]

V. album is a medicinal plant that contains a wide variety of phytochemicals. Quercetin, with high antioxidant activity and a strong antitumor effect, is one of the flavonols present in mistletoe being identified in *V. album* ssp. the *album*, *V. album* ssp. *abietis*, *V. album* ssp. *austriacum* and *V. album* ssp. *coloratum* [12, 63]. Among the flavanones, Naringenin and Eriodictyol are found both in *V. album* ssp. *album* and *V. album* ssp. *austriacum* [12, 51, 65]. Caffeic acid, a hydroxycinnamic acid, has been identified in various subspecies of *V. album* [12, 67].

Lectins and Viscotoxins

V. album is a medicinal plant that contains a wide variety of chemical compounds, from small molecules such as phenolic acids, flavonoids, to high molecular weight molecules proteins: lectins and viscotoxins [69].

Lectins are a class of proteins or glycoproteins, that can reversibly bind specific carbohydrates without altering their structure. Based on the specificity of carbohydrate-binding, lectins can be: monospecific - binds to one carbohydrate, (glucose, galactose) and polyspecific - binds to several sugars [24, 70]. In terms of chemical structure, mistletoe contains three lectins that differ in their molecular weight, about 60000Da and depending on the binding of carbohydrates [71, 72]. These lectins are coded as follows: ML 1, ML 2 and ML 3; (ML – Mistellelectin) [73]. Lectin ML 1, is a glycoprotein that has two polypeptide chains linked together by a disulfide bond [74]. The chain A of lectin ML 1 is composed of three distinct individual domains, having a molecular weight of 29 kDa, 27 kDa, and 25 kDa, possessing RNA N-glycosidase activity [75]. The lectin B chain consists of two domains with the same configuration, with a molecular weight of 32 kDa and 25 kDa, respectively [24]. It has a specific carbohydrate-binding system, it binds specifically to D-galactose [76, 77]. The specificity of lectin B chain binding to carbohydrates plays a key role in explaining the selective cytotoxicity of lectins to tumor cells when interacting with various receptors. The specific interaction of lectins with receptors on the surface of cancer cells causes agglutination, apoptosis, and the stopping of the cell cycle and thus inhibition of angiogenesis and cell proliferation [78]. Thus, both lectin chains have

cytotoxic action [24]. Lectin ML-2 binds specifically to D-galactose / N-acetyl-D-galactosamine [79]. Lectin ML-3 has a specificity for N-acetyl-D-galactosamine [72]. *V. an album* that grows on the following species: oak (*Quercus* ssp), poplar (*Populus nigra*) contains mostly lectin ML-1. *V. album* on conifers: fir (*Abies alba*), pine (*Pinus sylvestris*) contain mainly lectin ML-3 [71].

Viscotoxins are small polypeptides, rich in cysteine, and consist of about 46 amino acids, containing 3-4 disulfide bridges. They have a molecular weight of about 5KDa [80]. Depending on the amino acid sequence, the *V. album* contains 6 isomeric compounds: A1, A2, A3, B, B2, C1, 1-PS [72]. Viscotoxin has cytotoxic activity against different types of tumor cells and has immunomodulatory effects [15]. The composition and content of lectin and viscotoxin vary depending on the host tree.

In the *V. album* ssp *album* viscotoxin A2, A3 predominates, and 1-PS is missing while the *V. album* ssp *austriacum* is dominated by viscotoxin 1-PS, and viscotoxins A2, A3 have been detected in small amounts. *V. album* ssp *abietis* contains all viscotoxins, A3 predominates and viscotoxin A2 was not detected. [81]

Biological Properties and Pharmaceutical Action

Many biological effects of mistletoe, such as anticancer, apoptosis-inducing, antiviral, antibacterial, and immunomodulatory activities have been reported [2, 11, 20, 82, 83].

A series of *in vitro* and *in vivo* studies confirm a broad spectrum of the therapeutic action of mistletoe (*V. album*). **Table 3** summarizes the pharmacological activities of mistletoe (*V. album* ssp / *Viscum coloratum* / *Viscum articulatum*) depending on the host tree.

Table 3. The pharmacological activity of *V. album* ssp/ *Viscum coloratum*/ *Viscum articulatum*

Species/Host tree	Biological activity	Bioactive compounds	Sample type	Type of experiment	References
<i>Viscum album</i> / Apple (<i>Malus domestica</i>)	Anticancer	lectins	Aqueous Extract + triterpene extract	The human osteosarcoma cell lines 143B and Saos-2	[84]
<i>V. album</i> ssp. <i>coloratum</i> / Poplar (<i>Populus nigra</i>)		lectins	Aqueous extract	<i>In vitro</i> and <i>vivo</i> -on the growth of melanoma cells in mice.	[85]
<i>V. album</i> ssp. <i>coloratum</i> /ns*		viscotoxins	Aqueous extract	Randomized clinical trial in patients with locally advanced or metastatic pancreatic cancer	[86]
<i>Viscum album</i> / Apple (<i>Malus domestica</i>), Oak (<i>Quercus</i> ssp) Poplar (<i>Populus nigra</i>) Acacia (<i>Robinia pseudo-acacia</i>)	Patient with differentiated squamous cell carcinoma			[87]	
<i>V. album</i> L / Citrus	Antihypertension	Flavonoids	Aqueous extract	Male rats-salt induced hypertension	[88]
<i>V. album</i> L /ns*			Ethanollic extract	Wistar rats of both sexes	[89]
<i>V. album</i> /ns*			Ethanollic extract	Hypertensive patient	[90]
<i>V. articulatum</i> / <i>Cordia macleodi</i>			Acid oleonic	Methanollic extract	male Wistar rats- dexamethasone-induced hypertension
<i>V. album</i> / <i>Kola acuminata</i> tree	Antidiabetic		Aqueous extract	Alloxanized male Wistar rats	[92]
			Aqueous extract	STZ-diabetic male Wistar rats	[93]
<i>V. album</i>	Antivirals		Aqueous extract	Human parainfluenza virus type 2 (HPIV-2) growth in Vero cells	[83]
<i>V. album</i> ssp <i>abietis</i>	Antibacterial		n-hexane extract	<i>In vitro</i> (<i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Enterobacter cloacae</i> and <i>Proteus vulgaris</i>)	[94]
<i>V. album</i> /ns*			Extracts (1% HCl, ethanol, acetone, and 5% acetic acid)	<i>In vitro</i> (<i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> and <i>Klebsiella aerogenes</i>)	[95]
<i>V. album</i> / cocoa and cola trees				Methanollic extract	<i>In vitro</i> (<i>Aspergillus niger</i> , <i>Fusarium exosporium</i> , <i>Penicillium oxalium</i> , and <i>Microsporum canis</i>)

<i>V. album</i> L. ssp. <i>album/Armeniaca vulgaris</i> Lam. (apricot); <i>V. album</i> ssp. <i>abietis/Abies bornmülleriana</i> Mattf. (fir); ; <i>V. album</i> ssp. <i>austriacum/Pinus nigra</i> (pine)		Ethanollic extract	<i>In vitro</i> (<i>Mycobacterium tuberculosis</i> H37Ra)	[97]	
<i>V. album</i> L. ssp. <i>album/Armeniaca vulgaris</i> Lam. (apricot); <i>V. album</i> ssp. <i>abietis/Abies bornmülleriana</i> Mattf. (fir); ; <i>V. album</i> ssp. <i>austriacum/Pinus nigra</i> (pine)	Antioxidant	Ethanollic and aqueous extracts	STZ-induced diabetic rats/ (GSH and MDA in liver, kidney, and heart tissues)	[98]	
<i>V. album/Acer campestre; Fraxinus excelsio; Populus nigra; Malus domestica; Robinia pseudoacacia</i>		flavonoids	Aqueous extract	<i>In vitro</i> (DPPH, ORAC, and TEAC assay)	[65]
<i>V. coloratum</i> /ns*			Ethanollic extract	<i>In vitro</i> (Hydroxyl Radical Scavenging Assay; Superoxide Anion Radical Scavenging Assay)	[99]
<i>V. album</i> L/ Citrus	Antiepileptic	Aqueous extract	Swiss albino mice and Wistar albino rats (MES-induced seizure; INH-induced convulsions)	[100]	

*ns- not specified; MDA- malondialdehyde; MES- Maximum electroshock; INH-Isoniazid;

Therapeutic, Cosmetic, and Functional Applications of Mistletoe Extract

Table 4 shows the patents published with mistletoe extracts (*V. album*) with therapeutic, cosmetic, and functional applications in the period 2011-2021.

Table 4. List of patents published from on *V. album* (2011-2021)

Patent no.	Publication date	Title	Purpose
KR20110136539A	2011-12-21	Korean mistletoe extracts having antiobesity and hepatic steatosis protection activity	-suppresses obesity by inhibiting fat around epididymis and fatty liver
KR20140115730A	2014-10-31	Shampoo composition for preventing hair losing and promoting hair growth comprising extracts of <i>V. album</i> and <i>Chamaecyparis obtusa</i>	-prevents hair loss -possesses an improved antioxidant capacity
CN104707097A	2015-06-17	Pharmaceutical composition <i>Viscum</i> and astragali radix powder and use thereof in the preparation of medicines for blocking precancerous lesions of the liver and treating liver cancer or viral hepatitis	-reduce the GGT foci incidence of parts of precancerous lesions -hydroxyl free radical scavenging effect blocking the occurrence and development of precancerous lesions of the liver
KR20160017903A	2016-02-17	<i>V. album</i> extract containing a sweet jelly preparation method and sweet jelly prepared by the method	-enhances immunity -proposed for immunotherapy and cancer prevention
KR20170053544A	2017-05-16	Cosmetic composition for moisturizing and anti-aging-reduction of cytotoxicity caused by external stress containing extracts of <i>Viscum album</i> , <i>Chamaecyparis obtusa</i> , <i>Quercus Robur</i> , and <i>Camellia japonica</i> Linne	-skin astringent -control of the expression of inflammatory mediating factors.
CN106692909A	2017-05-24	<i>Pyrus pyrifolia</i> tree <i>Viscum coloratum</i> grease with lung-clearing and phlegm-eliminating effects	-effects on cleaning the lungs and eliminating phlegm
CN108295100A	2018-07-20	Application of the <i>V. album</i> extract in preparing drugs for rheumatoid arthritis	-inhibit the synthesis of inflammatory mediator downstream by targeting NF- κ B* signal accesses, -inhibit the inflammatory reaction of the synovial membrane, -mitigate the destruction of articular cartilage and bone.
KR102007096B1	2019-08-02	Composition for preventing or treating hearing loss comprising <i>Viscum ovalifolium</i> extracts	-suppress hearing loss

KR20200109068A	2020-09-22	A soap comprising <i>V. album</i> fermentation product and method for preparing the same	-easily absorbed into the skin, -alleviate skin inflammation. -exfoliating effect, it can remove harmful agents from the surface of the skin, -prevent the aging of the skin, due to the antioxidant components of the <i>V. album</i>
BR102020003563A2	2021-09-08	Topical pharmaceutical composition with antitumor activity containing <i>V. album</i>	-antitumor activity

GGT gamma-glutamyltranspeptidase, GPT glutamic—pyruvic transaminase, NF- κ B Nuclear factor- κ B

Conclusion and Future Prospects

Mistletoe is a plant with a high potential for the treatment of various diseases. Its phytochemical composition and at the same time its biological effects depend on the host tree. The key components responsible for anticancer activity are lectin and viscotoxin. In addition, mistletoe also contains a sufficient amount of secondary metabolites from the class of flavonoids and phenolic acids, compounds with important beneficial effects. These compounds are widely distributed in plants and there is sufficient evidence to show that their consumption is closely linked to a decrease in the incidence of cancer, diabetes, and cardiovascular disease. Studies on the synergistic effects between mistletoe polyphenols and lectins should be further developed to identify new targeted therapeutic applications of mistletoe preparations.

Acknowledgments: The authors acknowledge the support provided by the University of Oradea through the grant competition "Excellent scientific research related to the priority fields with the goal of technology transfer: INO-TRANSFER-UO", Project number 309/21.12.2021.

Conflict of interest: None

Financial support: This research was funded by the European Union, through the Horizon 2020 project "NextFood", Grant agreement No. 771738.

Ethics statement: None

References

- Amico GC, Nickrent DL, Vidal-Russell R. Macroscale analysis of mistletoe host ranges in the Andean-Patagonian forest. *Plant Biol Stuttg Ger.* 2019;21(1):150-6.
- Szurpnicka A, Zjawiony JK, Szterk A. Therapeutic potential of mistletoe in CNS-related neurological disorders and the chemical composition of *Viscum* species. *J Ethnopharmacol.* 2019;231:241-52.
- Combes C. Parasitism: the ecology and evolution of intimate interactions (interspecific interactions). Q Rev Biol. The University of Chicago Press; 2002;77:207. doi:10.1086/343967
- Krasylenko Y, Těšitel J, Ceccantini G, Oliveira-da-Silva M, Dvořák V, Steele D, et al. Parasites on parasites: hyper-, epi, and autoparasitism among flowering plants. *Am J Bot.* 2021;108(1):8-21.
- Böhling N, Greuter W, Raus T, Snogerup B, Snogerup S, Snogerup B. Notes on the Cretan mistletoe, *Viscum album* subsp. *creticum* subsp. *nova* (Loranthaceae/Viscaceae). *Isr J Plant Sci.* 2002;50(sup1):77-84.
- Costa JPL, Brito HO, Galvão-Moreira LV, Brito LGO, Costa-Paiva L, Brito LMO. Randomized double-blind placebo-controlled trial of the effect of *Morus nigra* L. (black mulberry) leaf powder on symptoms and quality of life among climacteric women. *Int J Gynaecol Obstet.* 2020;148(2):243-52.
- Zuber D. Biological flora of central Europe: *Viscum album* L. *Flora - Morphol Distrib Funct Ecol Plants.* 2004;199(3):181-203. Available from: <https://www.sciencedirect.com/science/article/pii/S0367253004701103>
- Peñaloza E, Holandino C, Scherr C, Araujo PIP de, Borges RM, Urech K, et al. Comprehensive metabolome analysis of fermented aqueous extracts of *Viscum album* L. by liquid chromatography-high resolution tandem mass spectrometry. *Mol Basel Switz.* 2020;25(17):E4006.
- Vergara-Barberán M, Lerma-García MJ, Nicoletti M, Simó-Alfonso EF, Herrero-Martínez JM, Fasoli E, et al. Proteomic fingerprinting of mistletoe (*Viscum album* L.) via combinatorial peptide ligand libraries and mass spectrometry analysis. *J Proteomics.* 2017;164:52-8.
- Bonamin LV, De Carvalho AC, Waisse S. *Viscum album* (L.) in experimental animal tumors: A meta-analysis. *Exp Ther Med.* 2017;13(6):2723-40. Available from: <https://www.spandidos-publications.com/10.3892/etm.2017.4372>
- Kwon YS, Chun SY, Kim MK, Nan HY, Lee C, Kim S. Mistletoe extract targets the STAT3-FOXO1 pathway to induce apoptosis and inhibits metastasis in breast cancer cells. *Am J Chin Med.* 2021;49(02):487-504. doi:10.1142/S0192415X21500221
- Pietrzak W, Nowak R, Gawlik-Dziki U, Lemieszek M, Rzeski W. LC-ESI-MS/MS identification of biologically active phenolic compounds in mistletoe berry extracts from different host trees. *Molecules.* 2017;22(4):624.

13. Pietrzak W, Nowak R. Impact of harvest conditions and host tree species on chemical composition and antioxidant activity of extracts from *Viscum album* L. *Mol Basel Switz.* 2021;26(12):3741.
14. Fritz P, Dippon J, Müller S, Goletz S, Trautmann C, Pappas X, et al. Is Mistletoe treatment beneficial in invasive breast cancer? a new approach to an unresolved problem. *Anticancer Res.* 2018;38(3):1585-93.
15. Rostock M. Mistletoe in the treatment of cancer patients. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 2020;63(5):535-40.
16. Wode K, Hök Nordberg J, Kienle GS, Elander NO, Bernhardson BM, Sunde B, et al. Efficacy of mistletoe extract as a complement to standard treatment in advanced pancreatic cancer: study protocol for a multicentre, parallel-group, double-blind, randomized, placebo-controlled clinical trial (MISTRAL). *Trials.* 2020;21(1):783.
17. Shakeel M, Trinidade A, Geider S, Ah-See KW. The case for mistletoe in the treatment of laryngeal cancer. *J Laryngol Otol.* 2014;128(3):302-6.
18. Rexer H, Geschäftsstelle der AUO. Study for the treatment of nonmuscle invasive bladder cancer: A phase III efficacy trial for intravesical instillation of mistletoe extract in superficial bladder cancer (TIM) - AB 40/11 of the AUO. *Urol Ausg A.* 2016;55(7):963-5.
19. Park YK, Do YR, Jang BC. Apoptosis of K562 leukemia cells by Abnobaviscum F®, a European mistletoe extract. *Oncol Rep.* 2012;28(6):2227-32. Available from: <https://www.spandidos-publications.com/10.3892/or.2012.2026>
20. Menke K, Schwermer M, Eisenbraun J, Schramm A, Zuzak TJ. Anticancer effects of *Viscum album fraxini* extract on medulloblastoma cells in vitro. *Complement Med Res.* 2021;28(1):15-22.
21. Schötterl S, Miemietz JT, Iliina EI, Wirsik NM, Ehrlich I, Gall A, et al. Mistletoe-based drugs work in synergy with radio-chemotherapy in the treatment of glioma in vitro and in vivo in glioblastoma bearing mice. *Evid-Based Complement Altern Med ECAM.* 2019;2019:1376140.
22. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71. Available from: <https://www.bmj.com/content/372/bmj.n71>
23. Halbritter DA, Willett DS, Gordon JM, Stelinski LL, Daniels JC. Behavioral evidence for host transitions in plant, plant parasite, and insect interactions. *Environ Entomol.* 2018;47(3):646-53.
24. Liu B, Bian H, Bao J. Plant lectins: Potential antineoplastic drugs from bench to clinic. *Cancer Lett.* 2010;287(1):1-12. Available from: <https://www.sciencedirect.com/science/article/pii/S0304383509003450>
25. Liu B, Le CT, Barrett RL, Nickrent DL, Chen Z, Lu L, et al. Historical biogeography of Loranthaceae (Santalales): Diversification agrees with the emergence of tropical forests and radiation of songbirds. *Mol Phylogenet Evol.* 2018;124:199-212. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1055790317308217>
26. Baena-Díaz F, Ramírez-Barahona S, Ornelas JF. Hybridization and differential introgression are associated with environmental shifts in a mistletoe species complex. *Sci Rep.* 2018;8(1):5591.
27. Maclean AE, Hertle AP, Ligas J, Bock R, Balk J, Meyer EH. Absence of complex i is associated with diminished respiratory chain function in European mistletoe. *Curr Biol.* 2018;28(10):1614-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960982218303658>
28. Fu J, Wan L, Song L, He L, Jiang N, Long H, et al. Identification of MicroRNAs in *Taxillus Chonensis* (DC.) Danser Seeds under Cold Stress. *BioMed Res Int.* 2021;2021:5585884.
29. Oei SL, Thronicke A, Schad F. Mistletoe and immunomodulation: insights and implications for anticancer therapies. *Evid-Based Complement Altern Med ECAM.* 2019;2019:5893017.
30. Jones P. The Tenbury mistletoe festival & national mistletoe day. *Holidappy.* [cited 2022 Jan 16]. Available from: <https://holidappy.com/holidays/The-Tenbury-Mistletoe-Festival>
31. Ostermann T, Appelbaum S, Poier D, Boehm K, Raak C, Büssing A. A systematic review and meta-analysis on the survival of cancer patients treated with a fermented *Viscum album* L. extract (Iscaador): an update of findings. *Complement Med Res.* 2020;27(4):260-71.
32. Ostermann T, Büssing A. Retrospective studies on the survival of cancer patients treated with mistletoe extracts: a meta-analysis. *Explore N Y N.* 2012;8(5):277-81.
33. Jäger T, Holandino C, Melo MN de O, Peñaloza EMC, Oliveira AP, Garrett R, et al. Metabolomics by UHPLC-Q-TOF reveals host tree-dependent phytochemical variation in *Viscum album* L. plants. 2021;10(8):1726. Available from: <https://www.mdpi.com/2223-7747/10/8/1726>
34. Ramm H, Urech K, Scheibler M, Grazi G. Cultivation and development of *viscum album* L. Mistletoe. CRC Press; 2000. pp. 91-110.
35. Maul K, Krug M, Nickrent DL, Müller KF, Quandt D, Wicke S. Morphology, geographic distribution, and host preferences are poor predictors of phylogenetic relatedness in the mistletoe genus *Viscum* L. *Mol Phylogenet Evol.* 2019;131:106-15.
36. Büssing A. Mistletoe: The Genus *Viscum*. CRC Press; 2000.
37. Wilson CA, Calvin CL. An origin of aerial branch parasitism in the mistletoe family, Loranthaceae. *Am J Bot.* 2006;93(5):787-96. doi:10.3732/ajb.93.5.787
38. Barney C, Hawksworth F, Geils B. Hosts of *Viscum album*. *Eur J for Pathol.* 2007;28(3):187-208.

39. Danser DG. Long. *Viscum album* subsp. *meridianum*. Plants World Online. 1982. Available from: <http://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:921668-1>
40. Long C, Fan R, Zhang Q, Zhang Z, Wang D, Xia Y, et al. Simultaneous identification and quantification of the common compounds of *Viscum coloratum* and its corresponding host plants by ultra-high performance liquid chromatography with quadrupole time-of-flight tandem mass spectrometry and triple quadrupole mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2017;1061:176-84. doi:10.1016/j.jchromb.2017.07.028
41. Lyu SY, Park SM, Choung BY, Park WB. Comparative study of Korean (*Viscum album* var. *coloratum*) and European mistletoes (*Viscum album*). *Arch Pharm Res.* 2000;23(6):592-8. doi:10.1007/BF02975247
42. Patel BP, Singh PK. *Viscum articulatum* Burm. f.: a review on its phytochemistry, pharmacology and traditional uses. *J Pharm Pharmacol.* 2018;70(2):159-77. doi:10.1111/jphp.12837
43. Kunwar R, Adhikari N, Devkota M. Indigenous use of mistletoes in tropical and temperate region of Nepal. *Banko Janakari.* 2007;15(2):38-42.
44. Deeni Y, Sadiq N. Antimicrobial properties and phytochemical constituents of the leaves of African mistletoe (*Tapinanthus dodoneifolius* (DC) Danser) (Loranthaceae): An ethnomedicinal plant of Hausaland, Northern Nigeria. *J Ethnopharmacol.* 2003;83(3):235-40.
45. Obatomi D, Bikomo EO, Temple V. Anti-diabetic properties of the African mistletoe in streptozotocin-induced diabetic rats. *J Ethnopharmacol.* 1994;43(1):13-7.
46. Clark NF, McComb JA, Taylor-Robinson AW, Clark NF, McComb JA, Taylor-Robinson AW. Host species of mistletoes (Loranthaceae and Viscaceae) in Australia. *Aust J Bot.* 2020;68(1):1-13.
47. Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A. The world checklist of vascular plants, a continuously updated resource for exploring global plant diversity. *Sci Data.* 2021;8(1):215. Available from: <https://www.nature.com/articles/s41597-021-00997-6>
48. Watson DM, Herring M. Mistletoe as a keystone resource: an experimental test. *Proc Biol Sci.* 2012;279(1743):3853-60.
49. Watson DM. Mistletoe—A keystone resource in forests and woodlands worldwide. *Annu Rev Ecol Syst.* 2001;32(1):219-49. doi:10.1146/annurev.ecolsys.32.081501.114024
50. Cordero CM, Gil Serrano AM, Ayuso Gonzalez MJ. Transfer of bipiperidyl and quinolizidine alkaloids to *Viscum cruciatum* Sieber (Loranthaceae) hemiparasitic on *Retama sphaerocarpa* boissier (Leguminosae). *J Chem Ecol.* 1993;19(10):2389-93. doi:10.1007/BF00979672
51. Zhang RZ, Zhao JT, Wang WQ, Fan RH, Rong R, Yu ZG, et al. Metabolomics-based comparative analysis of the effects of host and environment on *Viscum coloratum* metabolites and antioxidative activities. *J Pharm Anal.* 2021. Available from: <https://www.sciencedirect.com/science/article/pii/S2095177921000472>
52. Kaurinovic B, Vastag D. Flavonoids and phenolic acids as potential natural antioxidants. IntechOpen; 2019. Available from: <https://www.intechopen.com/chapters/65331>
53. Fritea L, Ganea M, Zdrinca M, Dobjanschi L, Antonescu A, Vicas SI, et al. Perspectives on the combined effects of *Ocimum basilicum* and *Trifolium pratense* extracts in terms of phytochemical profile and pharmacological effects. *Plants.* 2021;10(7):1390.
54. Vicas SI, Cavalu S, Laslo V, Tocai M, Costea TO, Moldovan L. Growth, photosynthetic pigments, phenolic, glucosinolates content and antioxidant capacity of broccoli sprouts in response to nanoselenium particles supply. *Not Bot Horti Agrobot Cluj-Napoca.* 2019;47(3):821-8.
55. Alabdallat NG. In-vivo antioxidant effects of the orally administered paracetamol, aqueous extracts of *salvia triloba*, and *origanum syriacum*. *J Biochem Technol.* 2021;12(4):19-22. Available from: <https://jbiochemtech.com/article/in-vivo-antioxidant-effects-of-the-orally-administered-paracetamol-aqueous-extracts-of-salvia-trilo-g3huxydynkdagsa>
56. Al-Harbi D, Awad N, Alsberi H, Abdein M. Apoptosis induction, cell cycle arrest and in vitro anticancer potentiality of *convolvulus spicatus* and *astragalus vogelii*. *World J Environ Biosci.* 2020;8(4):69-75. Available from: www.environmentaljournal.org
57. Darwati D, Safitri AN, Ambardhani N, Mayanti T, Nurlelasari N, Kurnia D. Effectiveness and anticancer activity of a novel phenolic compound from *garcinia porrecta* against the MCF-7 breast cancer cell line in vitro and in silico. *Drug Des Devel Ther.* 2021;15:3523-33. Available from: <https://www.dovepress.com/effectiveness-and-anticancer-activity-of-a-novel-phenolic-compound-fro-peer-reviewed-fulltext-article-DDDT>
58. Alotaibi BS, Ijaz M, Buabeid M, Kharaba ZJ, Yaseen HS, Murtaza G. Therapeutic Effects and safe uses of plant-derived polyphenolic compounds in cardiovascular diseases: a review. *Drug Des Devel Ther.* 2021;15:4713-32. Available from: <https://www.dovepress.com/therapeutic-effects-and-safe-uses-of-plant-derived-polyphenolic-compou-peer-reviewed-fulltext-article-DDDT>
59. Vicas S, Prokisch J, Rugina OD, Socaciu C. Hydrophilic and lipophilic antioxidant activities of mistletoe (*Viscum album*) as determined by FRAP method. *Not Bot Horti Agrobot Cluj-Napoca.* 2009;37(2):112-6.
60. Vicas SI, Rugin D, Socaciu C. Comparative study about antioxidant activities of *Viscum album* from different host trees, harvested in different seasons. *J Med Plants Res.* 2011;5(11):2237-44. Available from: <https://academicjournals.org/journal/JMPR/article-abstract/4B70C8121233>
61. Olanlokun JO, Ekundayo MT, Ebenezer O, Koorbanally NA, Olorunsogo OO. Antimalarial and erythrocyte membrane stability properties of *globimetula braunii* (engle van tiegh) growing on cocoa in plasmodium berghei-infected mice.

- Infect Drug Resist. 2021;14:3795-808. Available from: <https://www.dovepress.com/antimalarial-and-erythrocyte-membrane-stability-properties-of-globimet-peer-reviewed-fulltext-article-IDR>
62. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: an overview. *Sci World J.* 2013;2013:162750.
 63. Vicas SI, Rugin D, Socaciu C. Comparative study about antioxidant activities of *Viscum album* from different host trees, harvested in different seasons. *J Med Plants Res.* 2011;5(11):2237-44.
 64. Luczkiewicz M, Cisowski W, Kaiser P, Ochocka R, Piotrowski A. Comparative analysis of phenolic acids in mistletoe plants from various hosts. *Acta Pol Pharm.* 2001;58(5):373-9.
 65. Vicaş SI, Rugina OD, Leopold L, Pintea A, Socaciu C. HPLC fingerprint of bioactive compounds and antioxidant activities of *Viscum album* from different host trees. *Not Bot Horti Agrobot Cluj-Napoca.* 2011;39(1):48-57. Available from: <https://www.notulaebotanicae.ro/index.php/nbha/article/view/3455>
 66. Haas K, Bauer M, Wollenweber E. Cuticular waxes and flavonol aglycones of mistletoes. *Z Für Naturforschung C J Biosci.* 2003;58(7-8):464-70. Available from: <https://eurekamag.com/research/003/699/003699122.php>
 67. Stefanucci A, Zengin G, Llorent-Martínez E, Dimmito M, Della Valle A, Pieretti S, et al. *Viscum album* L. homogenizer-assisted and ultrasound-assisted extracts as potential sources of bioactive compounds. *J Food Biochem.* 2020;44(9):e13377.
 68. Urech K, Baumgartner S. Chemical constituents of *Viscum album* L.: Implications for the pharmaceutical preparation of mistletoe. *Mistletoe: From mythology to evidence-based medicine.* 2015;4:11-23.
 69. Elluru SR, Duong Van Huyen JP, Delignat S, Prost F, Bayry J, Kazatchkine M, et al. Molecular mechanisms underlying the immunomodulatory effects of mistletoe (*Viscum album* L.) extracts *Iscador*: Review. *Arzneimittelforschung.* 2006;56(06):461-6.
 70. Peumans WJ, Van Damme JM, Barre A, Rougé P. Classification of plant lectins in families of structurally and evolutionary related proteins. In: Wu AM, editor. *Mol Immunol Complex Carbohydr —2.* Boston, MA: Springer US; 2001. pp.27-54. doi:10.1007/978-1-4615-1267-7_3
 71. Peumans W, Verhaert P, Pfüller U, Damme EV. Isolation and partial characterization of a small chitin-binding lectin from mistletoe (*Viscum album*). *FEBS Lett.* 1996;396(2-3):261-5.
 72. Wacker R, Stoeva S, Pfüller K, Pfüller U, Voelter W. Complete structure determination of the A chain of mistletoe lectin III from *Viscum album* L. ssp. *album*. *J Pept Sci Off Publ Eur Pept Soc.* 2004;10(3):138-48.
 73. Mazalovska M, Kouokam JC. Transiently expressed mistletoe lectin II in *Nicotiana benthamiana* demonstrates anticancer activity in vitro. *Mol Basel Switz.* 2020;25(11):E2562.
 74. Kang TB, Song SK, Yoon TJ, Yoo YC, Lee KH, Her E, et al. Isolation and characterization of two Korean mistletoe lectins. *BMB Rep.* 2007;40(6):959-65. Available from: <http://www.koreascience.or.kr/article/JAKO200710103429562.page>
 75. Endo Y, Tsurugi K, Franz H. The site of action of the A-chain of mistletoe lectin I on eukaryotic ribosomes. The RNA N-glycosidase activity of the protein. *FEBS Lett.* 1988;231(2):378-80.
 76. Soler MH, Stoeva S, Schwamborn C, Wilhelm S, Stiefel T, Voelter W. Complete amino acid sequence of the A chain of mistletoe lectin I. *FEBS Lett.* 1996;399(1-2):153-7. doi:10.1016/s0014-5793(96)01309-9
 77. Wacker R, Stoeva S, Betzel C, Voelter W. Complete structure determination of N-acetyl-D-galactosamine-binding mistletoe lectin-3 from *Viscum album* L. *album*. *J Pept Sci Off Publ Eur Pept Soc.* 2005;11(6):289-302.
 78. Beztsinna N, de Matos MBC, Walther J, Heyder C, Hildebrandt E, Leneweit G, et al. Quantitative analysis of receptor-mediated uptake and pro-apoptotic activity of mistletoe lectin-1 by high content imaging. *Sci Rep.* 2018;8(1):2768.
 79. Franz H. Mistletoe lectins and their A and B chains. *Oncology.* 1986;43(Suppl 1):23-34.
 80. Bohlmann H, Apel K. Thionins. *Annu Rev Plant Biol.* 1991;42(1):227-40.
 81. Schaller G, Urech K, Grazi G, Giannattasio M. Viscotoxin composition of the three European subspecies of *Viscum album*. *Planta Med.* 1998;64(07):677-8. doi:10.1055/s-2006-957553
 82. Cavalu S, Damian G. Rotational correlation times of 3-carbamoyl-2, 2, 5, 5-tetramethyl-3-pyrrolin-1-yloxy spin label with respect to heme and nonheme proteins. *Biomacromolecules.* 2003;4(6):1630-5.
 83. Karagöz A, Önay E, Arda N, Kuru A. Antiviral potency of mistletoe (*Viscum album* ssp. *album*) extracts against human parainfluenza virus type 2 in Vero cells. *Phytother Res.* 2003;17(5):560-2.
 84. Kleinsimon S, Kauczor G, Jaeger S, Eggert A, Seifert G, Delebinski C. *Viscum*TT induces apoptosis and alters IAP expression in osteosarcoma in vitro and has synergistic action when combined with different chemotherapeutic drugs. *BMC Complement Altern Med.* 2017;17(1):1-3.
 85. Han SY, Hong CE, Kim HG, Lyu SY. Anti-cancer effects of enteric-coated polymers containing mistletoe lectin in murine melanoma cells in vitro and in vivo. *Mol Cell Biochem.* 2015;408(1):73-87.
 86. Tröger W, Galun D, Reif M, Schumann A, Stanković N, Miličević M. *Viscum album* [L.] extract therapy in patients with locally advanced or metastatic pancreatic cancer: a randomized clinical trial on overall survival. *Eur J Cancer.* 2013;49(18):3788-97. Available from: [https://www.cabdirect.org/cabdirect/abstract/20143031671?q=\(au%3A%22Reif%2C+M.%22\)](https://www.cabdirect.org/cabdirect/abstract/20143031671?q=(au%3A%22Reif%2C+M.%22))
 87. Werthmann P, Sträter G, Friesland H, Kienle G. Durable response of cutaneous squamous cell carcinoma following high-dose peri-lesional injections of *Viscum album* extracts – A case report. *Phytomedicine Int J Phytother Phytopharm.* 2013;20(3-4):324-7.

88. Ofem OE, Eno AE, Imoru J, Nkanu E, Unoh F, Ibu JO. Effect of crude aqueous leaf extract of *Viscum album* (mistletoe) in hypertensive rats. *Indian J Pharmacol.* 2007;39(1):15. Available from: <https://www.ijp-online.com/article.asp?issn=0253-7613;year=2007;volume=39;issue=1;spage=15;epage=19;aulast=Ofem;type=0>
89. Radenkovic M, Ivetic V, Popovic M, Brankovic S, Gvozdenovic L. Effects of mistletoe (*Viscum album* L., Loranthaceae) extracts on arterial blood pressure in rats treated with atropine sulfate and hexocycline. *Clin Exp Hypertens.* 2009;31(1):11-9. Available from: [https://www.cabdirect.org/cabdirect/abstract/20093082399?q=\(au%3a%22Radenkovic%2c+M.%22\)](https://www.cabdirect.org/cabdirect/abstract/20093082399?q=(au%3a%22Radenkovic%2c+M.%22))
90. Poruthukaren KJ, Palatty PL, Baliga MS, Suresh S. Clinical evaluation of *Viscum album* mother tincture as an antihypertensive: a pilot study. *J Evid-Based Complement Altern Med.* 2014;19(1):31-5.
91. Bachhav SS, Patil SD, Bhutada MS, Surana SJ. Oleanolic acid prevents glucocorticoid-induced hypertension in rats. *Phytother Res.* 2011;25(10):1435-9. doi:10.1002/ptr.3431
92. Shahabuddin M, Pouramir M, Moghadamnia A, Lakzaei M, Mirhashemi SM, Motallebi M. Antihyperglycemic and antioxidant activity of *Viscum album* extract. *Afr J Pharm Pharmacol.* 2011;5(3):433-6.
93. Adaramoye O, Amanlou M, Habibi-Rezaei M, Pasalar P, Ali MM. Methanolic extract of African mistletoe (*Viscum album*) improves carbohydrate metabolism and hyperlipidemia in streptozotocin-induced diabetic rats. *Asian Pac J Trop Med.* 2012;5(6):427-33.
94. Ertürk Ö, Kati H, Yayli N, Demirbag Z. Antimicrobial activity of *Viscum album* L. subsp. *abietis* (Viesb). *Turk J Biol.* 2003;27(4):255-8. Available from: <https://app.trdizin.gov.tr/makale/TXpBME1qYzM=/antimicrobial-activity-of-viscum-album-l-subsp-abietis-viesb->
95. Orhue PO, Edomwande EC, Igbinsosa E, Momoh ARM, Asekomhe OO. Antibacterial activity of extracts of mistletoe (*Tapinanthus dodoneifolius* (Dc) Dancer) from cocoa tree (*Theobroma cacao*). *Int J Herbs Pharmacol Res.* 2014;3(1):24-9. Available from: <https://www.ajol.info/index.php/ijhpr/article/view/104727>
96. Yusuf L, Oladunmoye MK, Ogundare AO, Mohmo MO, Daudu OA. Comparative antifungal and toxicological effects of the extract of mistletoes growing on two different host plants in Akure North, Nigeria. *Int J Biotechnol Food Sci.* 2014;2(2):31-4. Available from: <http://sciencewebpublishing.net/ijbfs/archive/2014/Feb/Abstract/Yusuf%20et%20al.htm>
97. Deliorman Orhan D. Evaluation of antimycobacterial activity of *Viscum album* subspecies. *Pharm Biol.* 2001;39(5):381-3.
98. Orhan DD, Aslan M, Sendogdu N, Ergun F, Yesilada E. Evaluation of the hypoglycemic effect and antioxidant activity of three *Viscum album* subspecies (European mistletoe) in streptozotocin-diabetic rats. *J Ethnopharmacol.* 2005;98(1-2):95-102.
99. Yao H, Liao ZX, Wu Q, Lei GQ, Liu ZJ, Chen DF, et al. Antioxidative flavanone glycosides from the branches and leaves of *Viscum coloratum*. *Chem Pharm Bull (Tokyo).* 2006;54(1):133-5. doi:10.1248/cpb.54.133
100. Gupta G, Kazmi I, Afzal M, Rahman M, Saleem S, Ashraf MS, et al. Sedative, antiepileptic and antipsychotic effects of *Viscum album* L. (Loranthaceae) in mice and rats. *J Ethnopharmacol.* 2012;141(3):810-6. Available from: <https://www.sciencedirect.com/science/article/pii/S0378874112001717>