



INTERVENTION OF MEDICINAL PLANTS FOR IMPROVING MALE FERTILITY

Kanan Gamanlal Gamit^{1*}, Manan Arunkumar Raval¹, Niraj Yashvantrai Vyas¹

1. Department of Pharmacognosy and Phytochemistry, Ramanbhai Patel College of Pharmacy, Charotar University of Science and Technology (CHARUSAT), Gujarat, India.

ARTICLE INFO

Received:

19 Apr 2022

Received in revised form:

27 Jul 2022

Accepted:

29 Jul 2022

Available online:

28 Aug 2022

Keywords: Infertility, Testosterone, Medicinal plants, Leydig cell

ABSTRACT

Sexuality is a complex interplay of multiple facets, including anatomical, physiological, psychological, developmental, cultural, and relational factors. This complex process might be divided into four phases, namely desire, arousal, orgasm, and resolution. Male infertility is a major problem worldwide which is defined as the inability of sexually active, non-contraceptive couples to have offspring in a year. Many factors affect the quality of semen including environmental, nutritional, and "lifestyle-related reasons. Male infertility problems may be contributory to 30 to 40 percent of infertile couples. There are many qualitative and quantitative parameters to check infertility in men which are quality of semen, number of sperm per ml of semen, sperm motility, abnormalities in sperm morphology, etc. Many plants are scientifically studied for their effect on male sexual dysfunction. Evidence showed that plants investigated on male infertility uplift the level of testosterone. The studies made it clear that testosterone plays an important role in sexual interest and associated sexual arousal.

This is an *open-access* article distributed under the terms of the [Creative Commons Attribution-Non Commercial-Share Alike 4.0 License](https://creativecommons.org/licenses/by-nc-sa/4.0/), which allows others to remix, and build upon the work non commercially.

To Cite This Article: Gamit KG, Raval MA, Vyas NY. Intervention of Medicinal Plants for Improving Male Fertility. Pharmacophore. 2022;13(4):72-9. <https://doi.org/10.51847/LoZXKRhy3I>

Introduction

The inability of a sexually active, non-contraceptive couple to become pregnant within a year is known as infertility [1]. According to a review by Agrawal *et al.*, the percentage of infertile men ranged from 2.5% to 12%, and the distribution of infertility caused by male factors ranged from 20% to 70%. The highest rates of infertility were found in Central/Eastern Europe and Africa [2].

A problem with the seminal fluid or the quantity and morphology of sperm could be the cause of male infertility. Men are deemed infertile when their sperm concentration falls below 20 million/mL [3]. According to WHO reports, there are 60-80 million infertile couples worldwide [1].

Around the world, 8–12% of couples experience infertility, with regional differences [4]. About 10% of American couples are classified as infertile if they are unable to conceive after 12 months of unprotected sexual activity [5, 6]. According to the National Center for Health Statistics, the total number of women with impaired fecundity increased by about 2.7 million between 1982 and 2002, from 4.56 million to 7.26 million, before slightly declining to 6.71 million between 2006 and 2010 [7]. In 2015, there were 22.3 teenagers born per 1,000 females, an 8% decrease from the previous year [8]. Prevalence of male sexual dysfunction in Asia is shown in **Table 1**.

The persistent inability to obtain and maintain an erection strong enough to allow for satisfactory sexual performance is known as erectile dysfunction (ED) [9]. Premature ejaculation is a male sexual dysfunction that is characterized by ejaculation that always, or nearly always, occurs before or within a minute of vaginal penetration, according to the International Society of Sexual Medicine [10]. As a clinical and biochemical syndrome associated with getting older and characterized by sexual dysfunction and other typical symptoms as well as a lack of serum testosterone levels, hypogonadism is the clinical and biochemical entity formerly known as andropause, androgen deficiency in the aging male, and partial androgen deficiency in the aging male [11].

Table 1. Prevalence of male sexual dysfunction in Asia [12]

Corresponding Author: Kanan Gamanlal Gamit; Department of Pharmacognosy and Phytochemistry, Ramanbhai Patel College of Pharmacy, Charotar University of Science and Technology (CHARUSAT), Gujarat, India. E-mail: kanangamit.ph@charusat.ac.in

Prevalence (%)			
Country	Erectile dysfunction	Premature ejaculation	Hypogonadism
Malaysia	26.8-69	22.3	18.5
Mainland China	38.3	19.5	-
Korea	32.2	27.5	-
Taiwan, China	9-17.7	13	24.1
Thailand	37.5	23	-
Singapore	51.3-73	12	-
Hong Kong, China	63.6	29.7	9.52

Although there are many treatments available, including phosphodiesterase type 5 inhibitors, intracavernosal prostaglandin E1, medicated urethral systems for erection, vacuum devices, and penile prostheses for erectile dysfunction; testosterone replacement therapy for hypogonadism; psychological and behavioral therapy, local anesthetic creams, tramadol, and selective serotonin reuptake inhibitors are used for premature ejaculation [12]. The use of numerous plants and Ayurvedic formulas as aphrodisiacs is documented in ancient literature.

Meiosis, spermatogenesis, and mitotic cell division all play a part in the intricate process known as spermatogenesis. Both endocrine and paracrine mechanisms are involved in regulating spermatogenesis. Follicle stimulating hormone (FSH) and luteinizing hormone (LH), the latter acting through the intermediary of testosterone, which is produced by the Leydig cells in the testis, are both involved in the endocrine stimulation of spermatogenesis [13].

The keywords, male infertility, clinical evaluation, and medicinal plants were searched on PubMed, MEDLINE, EBSCO, and Google Scholar. Around 250 articles were reviewed on the effect of medicinal plants on male infertility. Those articles which showed no or less effect of medicinal plants on infertility were excluded. Preclinical and clinical studies on effects of plants on male infertility are mentioned in **Table 2** and **Table 3** respectively.

Table 2. Preclinical studies on effects of medicinal plants on male infertility

Name of the plant	Part of the plant	Animal/Cell used	Dose level	Efficacy	Mechanism	Ref
<i>Pedaliumpurex</i>	Ethanol extract of the fruits	Albino rats	50, 100 and 150 mg/kg	Increased and improved sexual performance, increased levels of serum testosterone	Nitric oxide release	[14]
<i>Pedaliumpurex</i>	Petroleum ether extract of a plant	Albino rats	200 and 400 mg/kg	Higher increment of mating performance, increased pregnancy rate, increased levels of testosterone by antioxidants	Viable sperm of the male rats, protection of germ and Sertoli cells	[15]
<i>Citrulluslanatus</i>	Methanolic extract of the fruit rind of the fruit	Albino rats	200 mg/kg	Improvement in all the semen parameters, sperm count; increase in FSH, LH, and testosterone	High phenolic flavonoid and ascorbic acid contents act as anti-oxidants; ascorbic acid enhances testosterone synthesis and release of FSH and LH	[16]
<i>Amaranthusspinosus</i>	Methanolic extract of stem	Albino rats	250 and 500 mg/kg	Increase in the sperm count, increase in serum and testicular testosterone; the weight of testis,	-	[17]
<i>Dactyloctenaghatagirea</i>	Lyophilized aqueous extract of roots	Albino rats	200 mg/kg	Increased sexual behavior, penile erection index	Involvement of nitrous oxide (NO)	[18]

<i>Lecaniodiscus cupanioides</i>	Aqueous extract of roots	Wistar	25, 50, and 100 mg/kg	Increased sexual behavior, increased levels of testosterone, LH, FSH	Reduce the synaptic concentration of serotonin as well as increase the amount of dopamine release	[19]
<i>Fadogia agrestis</i>	Aqueous extract of stem	Albino rats	18, 50 and 100 mg/kg	Increased sexual behavior, testosterone,	Because it increases androgen, saponin, a plant component, enhances the aphrodisiac effects.	[20]
<i>Piper guineense</i>	Aqueous extract of dry fruits	Sprague Dawley	200 mg/kg	Increase in testicular weight, testosterone; normal parameter of semen analysis	Zinc is a co- essential factor for the endogenous production of testosterone by inhibiting the conversion of testosterone to dihydrotestosterone	[21]
<i>Garcinia kola</i>	Ethanollic extract of seeds	Wistar rats	200 and 400 mg/kg	Increased pre-coital sexual behavior; increase in weight of testis, testosterone,	factors that increase gonadotropins or locally through an increase in Leydig cells' number or their sensitivity to LH	[22]
<i>Fumaria parviflora</i>	Ethanollic extract of leaves	Wistar rats	100, 200 and 400 mg/kg	Increases in the testis and epididymis weight; seminiferous tubules diameter and germinal epithelium height were higher; increase in sperm density; No significant differences in serum FSH and LH; increase in testosterone	Increased androgen biosynthesis, improvement of testicular oxidative status by its extract components	[23]
<i>Nigella sativa</i>	Ethanollic extract of seeds	Wistar rats	200 and 400 mg/kg	No effect on the sperm motility and viability; increase in sperm count, testosterone, LH, fertility index	Increase in LH hormone concentration; LH stimulates the production of testosterone in Leydig cells, which stimulates spermatogenesis via testosterone, Thymoquinone	[24]
	Seed oil	Rabbits	5 ml/kg	The higher number of spermatozoa, an increase in testosterone,	Thymoquinone as antioxidant	[25]
<i>Cardiospermum halicacabum L</i>	Aqueous extract of leaves	Wistar rats	100 and 200 mg/kg	No significant change in reproductive organ weight; a significant increase in total sperm count, motile sperm, testosterone	An increase in serum testosterone levels, antioxidant activity brought on by flavonoids, and an increase in plasma levels of testosterone that may be caused by plant saponin components	[26]
<i>Basella alba</i>	Methanollic extract of leaves	Leydig cells	10 and 100 µg/ml	Stimulates testosterone, estradiol, mRNA levels of aromatase,	Regulation of protein kinases and cAMP	[27]
<i>Basella alba</i> and <i>Hibiscus macranthus</i> Hochst A ex Rich	Aqueous extract of both fresh and dried leaves	Albino rats	720 mg/kg (Fresh leaves extract), 108 mg/kg (Dry leaves extract)	An abundance of spermatozoa in the seminiferous tubule, an increase in testosterone level,	Induction of synthesis of testosterone by the Leydig cells	[28]
<i>Carpolobia alba</i>	Aqueous extract of roots	Sprague-Dawley	0.001, 0.01, 0.1, 1 and 10 mg/kg	Increase the level of testosterone <i>in vivo</i> and <i>in vitro</i>	-	[29]

<i>Anacyclus pyrethrum</i>	Alcoholic extract of roots	Albino rats	50, 100 and 150 mg/kg	Increase sperm count, motility, serum FSH, LH, and testosterone	Stimulation of the hypothalamus-adenohypophysis- gonad axis	[30]
<i>Jurenia dolomieu</i>	Methanolic extract of the root	Sprague-Dawley	200 and 400 mg/kg	Increase the level of testosterone; density of germinal cells and sperms in the seminiferous tubules	Pituitary LH release may be caused by saponins, whereas the synthesis of androgens may be mediated by flavonoids.	[31]
<i>Urtica dioica</i>	Hydro-alcoholic extract of leaves	BALB/c male mice	10, 20, and 50 mg/kg	Increase in weight of testis, sperm motility, sperm count, testosterone; normal sperm morphology	Enhancing the anti-oxidant defense, inhibition of cannabinoids' activity	[32]
<i>Allium sativum</i>	Aqueous extracts of bulb	Sprague-Dawley	200 mg/kg	Increase in testicular weight, Serum testosterone	According to reports, selenium and zinc regulate testicular activities, particularly the production of steroidogenic enzymes.	[33]
<i>Eruca sativa</i>	Aqueous extract of leaves	Albino mice	30 and 40 mg/kg	Increase in testosterone, a significant decrease in sperm mortality and abnormalities, a significant increase in diameter of seminiferous tubules and spermatid	Encourage the development of the testes and improve spermatozoa proliferation, maturation, and differentiation	[34]
<i>Ashwagandhadi Lehya (Withania somnifera)</i>	Ayurvedic formulation <i>Ashwagandhadi lehya</i>	Wistar rats	250, 500, 750 mg/kg <i>Ashwagandhadi Lehya</i> , 250 mg/kg <i>Withania somnifera</i>	The increased serum concentration of testosterone and cholesterol; spermatozoa, the diameter of seminiferous tubules	Nitric oxide synthesis	[35]
<i>Hygrophila spinosa T. Ander</i>	Unsonified fraction of seeds	Wistar rats	100, 200 and 300 mg/kg	Increased concentration of testosterone, sperm counts; increased diameter of seminiferous tubules, high numbers of spermatozoa in seminiferous tubules	Through interaction with Leydig cells, a fraction could elevate serum testosterone levels in animals receiving treatment.	[36]
<i>Argyria speciosa</i>	Alkaloidal fraction of root	Leydig cell	10, 100, and 1000 µg/ml	Increased concentration of testosterone.	Synthesis of testosterone.	[37]
<i>Phyllanthus amarus</i>	Methanolic extracts from the leaves	Guinea pigs	100, 200, 400 and 800 mg/kg	Increases in the level of testosterone but has little or no effect on the levels of FSH and LH	Increase in the level of testosterone can lead to an increase in the spermatozoa	[38]
	Aqueous and methanolic extract of leaves and stem	Albino rats	200 and 400 mg/kg	An increase in Testosterone levels increased Sperm concentration	Stimulation of the sex organs	[39]

Table 3. Clinical studies on medicinal plants which improve fertility

Plant's name	Patients & methods	Period of treatment	Efficacy	Findings	Ref
<i>Zingiber officinale</i>	75 infertile men	3 months	Increase in sperm count, sperm motility, sperm viability, normal sperm morphology; increase in ejaculate volume; increases in serum FSH, LH & testosterone	Increase the α -glycosidase enzyme in the epididymis, and fructose sugar in seminal vesicle	[40]

<i>Mucuna pruriens</i>	150 men	3 months	Infertile men's levels of testosterone, LH, dopamine, adrenaline, and noradrenaline significantly improved	Dopamine-mediated CNS action and adrenaline- and noradrenaline-mediated action on the reproductive tract	[41]
	75	3 months	increased serum levels of testosterone and LH, decreased levels of FSH and prolactin, and improved sperm motility and count.	Choline, beta-sitosterol, withanolides I–VII, withaferin A, and sitoindosides VII–X may all play a role in the various effects of <i>W. somnifera</i> on semen properties.	[42]
<i>Withania somnifera</i>	46(n=21)	90 days	Increase in sperm count, semen volume, motility; increase in serum testosterone, LH	Higher levels of testosterone;	[43]
	50	90 days	Increase in serum testosterone, sperm concentration, % motile sperm	Improved serum testosterone; testicular synthesis of peptides, which had effects on sperm motility	[44]
<i>Nigella sativa</i> L.	80(n=40)	2 months	Significant improvement in sperm count, motility and morphology and semen volume, pH, and round cells	Antioxidant properties of <i>N. sativa</i> oil	[45]
<i>Alpinia officinarum</i>	76 (n=31)	3 months	An increase in sperm count, the total number of spermatozoa with normal morphology,	Antioxidant and scavenging activity against the ROS via its phytochemical mainly including galangin	[46]
Sesame	25	3 months	Significant improvement in the sperm count and motility	Antioxidant	[47]
<i>Crocus sativus</i> L	20	10 days	Significant improvement in tip rigidity and tip tumescence; higher ILEF-15 score; increased number and duration of erectile events	Antioxidant	[48]
<i>Panax ginseng</i>	119	8 weeks	Higher ILEF-15 PEDT score	Vas deferens and seminal vesicles' increased NO synthesis, decreased sympathetic tone, and smooth muscle dilation	[49]
<i>Tribulus terrestris</i> and <i>Ecklonia bicyclis</i>	84	30 days	Improved ILEF-15 score; increase in testosterone	Activate the pathways of NO; ACE inhibitory activity	[50]

Spermatogenesis is a process by which spermatozoa (sperms) are formed in spermatogonia in the seminiferous tubule in the testes. The formation of spermatozoa from a single cell to the entire sperm takes seventy-four days to complete the process. At the age of fourteen years, the spermatogenesis process starts in males [51]. Various hormones are responsible for the process of spermatogenesis directly or indirectly. Numerous hormones that act directly or indirectly on spermatogenesis.

The hypothalamic neurosecretory cells secrete more gonadotropin-releasing hormone (GnRH), which in turn stimulates the anterior pituitary to secrete more luteinizing hormone (LH) and follicle-stimulating hormone (FSH).

Follicle Stimulating Hormone (FSH)

FSH is the hormone that triggers the beginning of spermatogenesis. FSH binds to Sertoli cells and spermatogonia and influences the growth of spermatogonia. Furthermore, FSH stimulates Sertoli cells' production and secretion of Androgen Binding Protein (ABP). The testosterone concentration in seminiferous tubules stays high because secreted ABP binds to testosterone. Testosterone is the hormone that stimulates the final step in the process of spermatogenesis. Sertoli cells release the hormone inhibin, after the process of spermatogenesis is over.

Testosterone

The male organ known as the testes produces testosterone, a steroidal hormone, from cholesterol. It is in charge of preserving spermatogenesis. Through the negative feedback mechanism of testosterone, hypothalamic neurosecretory cells suppress GnRH secretion. The key hormone for the development of the accessory sex organs, including the penis and scrotum, is testosterone (genital duct, seminal vesicle, and prostate gland). Due to testosterone, males develop a more masculine body, larger sex organs, longer bones, wider shoulders, longer pelvises, thicker skin, more evenly distributed hair, and a deeper voice.

Luteinizing Hormone

The Cholesterol Side Chain Cleavage (CSCC) reaction, which increases the conversion of cholesterol to pregnenolone, is triggered when a receptor on a particular membrane on the surface of a Leydig cell binds to LH. This reaction increases cyclic AMP and calcium ions (Ca^{2+}), protein kinase activation, phosphorylation, and protein synthesis. The cholesterol side chain cleavage (CSCC) reaction, which takes place in the mitochondria and is catalyzed by an enzyme complex, is typically regarded as the rate-limiting step in the steroid synthesis process. In Leydig cells, pregnenolone leaks from the mitochondria and is converted to a number of other steroids, including testosterone as the main end product.

Growth Hormone

For general metabolic processes in testis, growth hormone is necessary. It is also essential for the proliferation of spermatogonia.

Inhibin

Upon stimulating FSH, the Sertoli cell releases inhibin. A peptide hormone by the name of inhibin, inhibin inhibits the release of FSH through a feedback mechanism and, as a result, is crucial for spermatogenesis. There is an increase in inhibin secretion at the same time that the rate of spermatogenesis increases. A slower rate of spermatogenesis is caused by Inhibin's action on the anterior pituitary, which inhibits the release of FSH [52].

Results and Discussion

Sexual dysfunction is a major health issue nowadays. It is mostly affected by lifestyle, stress, and environmental factor or it may be genetic. This review main emphases on natural compounds with Aphrodisiac potential. Medicinal plants, extract/fraction, active constituents, or chemical moiety which enhances sexual drive or pleasure are termed Aphrodisiac. These compounds improve sexual dysfunction and behaviors. This review indicates the effectiveness of medicinal plants at the hormonal level and molecular level which improve sexual dysfunction like *Pedaliom murex*, *Citrullus lanatus*, *Amaranthus spinosus*, *Piper guineense*, etc. Furthermore, many Ayurvedic formulations have been investigated for their aphrodisiac potential. Many medicinal plants including *Withania somnifera*, *Mucuna pruriens*, *Tribulus terrestris*, and *Panax ginseng* proved their spermatogenic potential clinically.

According to the review, antioxidant effects are linked to the positive effects of medicinal plants on male reproductive function, which confirmed the ability of phytoconstituents to treat male infertility due to their presence.

The desire for sex, erection, mounting frequency, and ejaculatory frequency have all been treated with Ayurvedic formulations and medicinal plants. The accomplishment of cellular events is established by these features of male reproductive function.

In vivo studies on mammals typically last from a few days to several months, according to the review that has been presented. Additionally, the dosage of the medication varies, which may be brought on by the presence of phytoconstituents in a particular extract. Additionally, many medicinal plants continue to exist unexplored and are used by professionals in the trade. Therefore, substantiating claims about medicinal plants and defining their typical use is a huge challenge for researchers.

Conclusion

Herbal and Ayurvedic products are widely used in developing countries for the treatment of male infertility. Plants or products which have been clinically evaluated for treatment should be advised to treat the male infertility problem. Plants that are evaluated in vivo or in vitro for their claimed action should be encouraged for complete clinical evaluation of potential effects in the treatment of male infertility disorder. Also, the physicians treating male infertility must know those plants which are scientifically investigated and should combine traditional therapy with the modern one.

Acknowledgments: Authors are thankful to Ramanbhai Patel College of Pharmacy and CHARUSAT for providing necessary facilities.

Conflict of interest: None

Financial support: None

Ethics statement: None

References

1. WHO. WHO manual for the standardized investigation and diagnosis of the infertile couple. 2000.
2. Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe. *Reprod Biol Endocrinol*. 2015;13(1):1-9.
3. Shah KM, Gamit KG, Raval MA, Vyas NY. Male infertility: A scoping review of prevalence, causes, and treatments. *Asian Pacific J Reprod J*. 2021;10(5):195-202.
4. Sciarra J. Infertility: An international health problem. *Int J Gynecol Obstet*. 1994;46(2):155-63.

5. Stephen EH, Chandra A. Updated projections of infertility in the United States: 1995- 2025. *Fertil Steril.* 2000;70(1):30-4.
6. Stephen EH, Chandra A. Declining estimates of infertility in the United States: 1982 – 2002. *Fertil Steril.* 2006;86(3):516-23.
7. Chandra A, Copen CE, Stephen EH. Infertility and impaired fecundity in the United States, 1982-2010: Data from the national survey of family growth. U.S. Department of Health and Human Services, Centers for Disease Control and Prevention National Center for Health Statistics; 2013. 1-22 p.
8. Martin JA, Hamilton BE, Osterman MJK, Driscoll AK. Births: Final Data for 2015. *Natl Vital Stat Rep.* 2017;66(1):1-70.
9. Hatzimouratidis K, Amar E, Eardley I, Giuliano F, Montorsi DH, Yoram Vardi, et al. Guidelines on male sexual dysfunction: Erectile dysfunction and premature ejaculation. *Eur Urol.* 2010;57(5):804-14.
10. McMahon CG, Althof SE, Waldinger MD, Porst H, Dean J, Sharlip ID, et al. An evidence-based definition of lifelong premature ejaculation: Report of the international society for sexual medicine (ISSM) Ad hoc committee for the definition of premature ejaculation. *J Sex Med.* 2008;5(7):1590-606.
11. Nieschlag E, Swerdloff R, Behre HM, Gooren LJ, Kaufman JM, Legros JJ, et al. Investigation, treatment and monitoring of late-onset hypogonadism in males. *Aging Male.* 2005;8(2):56-8.
12. Ho CCK, Singam P, Hong GE, Zainuddin Z. Male sexual dysfunction in Asia. *Asian J Androl.* 2011;13(4):537-42.
13. De Kretser DM, Loveland KL, Meinhardt A, Simorangkir D, Wreford N. Spermatogenesis. *Hum Reprod.* 1998;13(suppl_1):1-8.
14. Sharma V, Thakur M, Dixit VK. A comparative study of ethanolic extracts of *Pedaliu murex* Linn. fruits and sildenafil citrate on sexual behaviors and serum testosterone level in male rats during and after treatment. *J Ethnopharmacol.* 2012;143(1):201-6. doi:10.1016/j.jep.2012.06.024
15. Balamurugan G, Muralidharan P, Polapala S. Aphrodisiac activity and curative effects of *Pedaliu murex* (L.) against ethanol-induced infertility in male rats. *Turkish J Biol.* 2010;34(2):153-63.
16. Kolawole T. Ameliorative Effects of the Methanolic Extract of the Rind of *Citrullus lanatus* on Lead Acetate Induced Toxicity on Semen Parameters and Reproductive Hormones of Male Albino Wistar Rats. *European J Med Plants.* 2014;4(9):1125-37.
17. Sangameswaran B, Jayakar B. Anti-diabetic, anti-hyperlipidemic and spermatogenic effects of *Amaranthus spinosus* Linn. on streptozotocin-induced diabetic rats. *J Nat Med.* 2008;62(1):79-82.
18. Thakur M, Dixit VK. Aphrodisiac activity of *Dactylorhiza hatagirea* (D.Don) Soo in male albino rats. *Evid Based Complement Alternat Med.* 2007;4(S1):29-31.
19. Ajiboye TO, Nurudeen QO, Yakubu MT. Aphrodisiac effect of aqueous root extract of *Lecaniodiscus cupanioides* in sexually impaired rats. *J Basic Clin Physiol Pharmacol.* 2014;25(2):241-8.
20. Yakubu MT, Akanji MA, Oladiji AT. Aphrodisiac potentials of the aqueous extract of *Fadogia agrestis* (Schweinf. Ex Hiern) stem in male albino rats. *Asian J Androl.* 2005;7(4):399-404.
21. Memudu A, Akinrinade I, Ogundele O, Dare B. Effects of Crude Extract of Dry Fruits of *Piper guineense* on Male Fertility Parameters of Adult Sprague Dawley Rats. *European J Med Plants.* 2015;5(3):297-303.
22. Ralebona N, Sewani-Rusike CR, Nkeh-Chungag BN. Effects of ethanolic extract of *Garcinia kola* on sexual behaviour and sperm parameters in male Wistar rats. *Afr J Pharm Pharmacol.* 2012;6(14):1077-82.
23. Dorostghoal M, Seyyednejad SM, Khajepour L, Jabari A. Effects of *Fumaria parviflora* leaves extract on reproductive parameters in adult male rats. *Iran J Reprod Med.* 2013;11(11):891-8.
24. Umar Z, Qureshi AS, Rehan S, Ijaz M, Faisal T, Umar S. Effects of oral administration of black seed (*Nigella sativa*) oil on histomorphometric dynamics of testes and testosterone profile in rabbits. *Pak J Pharm Sci.* 2017;30(2):531-6.
25. Parandin R, Yousofvand N, Ghorbani R. The enhancing effects of alcoholic extract of *Nigella sativa* seed on fertility potential, plasma gonadotropins and testosterone in male rats. *Iran J Reprod Med.* 2012;10(4):355-62.
26. Peiris LDC, Dhanushka MAT, Jayathilake TAHDG. Evaluation of aqueous leaf extract of *Cardiospermum halicacabum* (L.) on fertility of male rats. *Biomed Res Int.* 2015;2015.
27. Nantia EA, Travert C, Manfo FPT, Carreau S, Monsees TK, Moundipa PF. Effects of the methanol extract of *Basella alba* L (Basellaceae) on steroid production in Leydig cells. *Int J Mol Sci.* 2011;12(1):376-84.
28. Moundipa FP, Kamtchouing P, Koueta N, Tantchou J, Foyang NPR, Mbiapo FT. Effects of aqueous extracts of *Hibiscus macranthus* and *Basella alba* in mature rat testis function. *J Ethnopharmacol.* 1999;65(2):133-9.
29. Manfo FPT, Nantia EA, Tchana AN, Monsees TK, Moundipa PF. Evaluation of the effect of *carpolobia alba* (Polygalaceae) aqueous extract on male reproductive function in rats. *J Appl Anim Res.* 2011;39(1):80-4.
30. Shahraki MR, Dehvari J, Shahrakipoor M, Shahreki E, Sharaki AR, Dashipour AR. The effects of anacyclus pyrethrum alcoholic root extract on FSH, LH, testosterone and sperm count in diabetic male rats. *Zahedan J Res Med Sci.* 2019;21(2).
31. Shah NA, Khan MR. Increase of glutathione, testosterone and antioxidant effects of *Jurenia dolomiaea* on CCl4 induced testicular toxicity in rats. *BMC Complement Altern Med.* 2017;17(1):1-9.
32. Jalili C, Salahshoor MR, Naseri A. Protective effect of *Urtica dioica* l against nicotine-induced damage on sperm parameters, testosterone and testis tissue in mice. *Iran J Reprod Med.* 2014;12(6):401-8.

33. Memudu AE, Akinrinade ID, Ogundele OM. Retention of testicular integrity and testosterone levels upon ingestion of garlic cloves (*Allium sativum*) in the Sprague-Dawley rat. *Asian Pac J Trop Biomed.* 2015;5(4):319-23. doi:10.1016/S2221-1691(15)30351-8
34. Hussein ZF. Study the Effect of *Eruca Sativa* Leaves Extract on Male Fertility in Albino Mice. *J Al-Nahrain Univ Sci.* 2013;16(1):143-6.
35. Gamit KG, Vyas NY, Chudasama P, Raval MA. Aphrodisiac and spermatogenic potential of Ayurveda formulation-Ashwagandhadi Lehya. *J Biol Act Prod Nat.* 2020;10(4):285-302.
36. Vyas N, Gamit K, Raval M. Aphrodisiac and spermatogenic potential of unsaponifiable fraction from seeds of *Hygrophila spinosa* t. Ander in rats. *Int J Pharm Sci Res.* 2020;11(10):4902-9.
37. Vyas N, Gamit K, Raval M, Patel SG. Isolation and chemical characterization of bioactive alkaloid from *Argyrea speciosa* Linn. having action on isolated rat Leydig cells. *Asian J Pharm Clin Res.* 2019;12(10):276-80.
38. Obianime A, Uche F. The Phytochemical constituents and the effects of methanol extracts of *Phyllanthus amarus* leaves (kidney stone plant) on the hormonal parameters of Male guinea pigs. *J Appl Sci Environ Manag.* 2010;13(1).
39. Azubuike NC, Okwuosa CN, Onwukwe OS, Onyemelukwe AO, Ikele I, Achukwu PU. Effects of *Phyllanthus amarus* on epididymal sperm characteristics, testosterone levels and histology of reproductive organs of male rats. *Pharmacologyonline.* 2018;3:57-67.
40. Mares WAA, Najam WS. The effect of Ginger on semen parameters and serum FSH, LH & testosterone of infertile men. *Tikrit Med J.* 2012;18(2):322-9.
41. Shukla KK, Mahdi AA, Ahmad MK, Shankhwar SN, Rajender S, Jaiswar SP. *Mucuna pruriens* improves male fertility by its action on the hypothalamus-pituitary – gonadal axis. *Fertil Steril.* 2009;92(6):1934-40.
42. Ahmad MK, Mahdi AA, Shukla KK, Islam N, Rajender S, Madhukar D, et al. *Withania somnifera* improves semen quality by regulating reproductive hormone levels and oxidative stress in seminal plasma of infertile males. *Fertil Steril.* 2010;94(3):989-96.
43. Ambiyi VR, Langade D, Dongre S, Aptikar P, Kulkarni M, Dongre A. Clinical Evaluation of the Spermatogenic Activity of the Root Extract of *Ashwagandha* (*Withania somnifera*) in Oligospermic Males: A Pilot Study. *Evid Based Complement Altern Med.* 2013;2013:1-6.
44. Raval M, Patel M, Gamit K, Patel K, Gupta SN. A Pilot Study on Evaluation of Standardized Ayurveda formulation *Ashwagandhadi lehya* as Aphrodisiac and in treatment of Oligospermia. *Res J Pharm Tech.* 2019;12(5):2383-90.
45. Kolahdooz M, Nasri S, Modarres SZ, Kianbakht S, Huseini HF. Effects of *Nigella sativa* L. seed oil on abnormal semen quality in infertile men: A randomized, double-blind, placebo-controlled clinical trial. *Phytomedicine.* 2014;21(6):901-5.
46. Kolangi F, Shafi H, Memariani Z, Kamalinejad M, Shirafkan H, Bioos S, et al. Effect of *Alpinia officinarum* Hance rhizome extract on spermatogram factors in men with idiopathic infertility: A prospective double-blinded randomized clinical trial. *Andrologia.* 2018; 51(1):e13172.
47. Khani B, Bidgoli SR, Moattar F, Hassani H. Effect of sesame on sperm quality of infertile men. *J Res Med Sci.* 2013;18(3):184-7.
48. Shamsa A, Hosseinzadeh H, Molaei M. Evaluation of *Crocus sativus* L. (saffron) on male erectile dysfunction: A pilot study. *Phytomedicine.* 2009;16(8):690-3.
49. Choi YD, Park CW, Jang J, Kim SH, Jeon HY, Kim WG, et al. Effects of Korean ginseng berry extract on sexual function in men with erectile dysfunction: a multicenter, placebo-controlled, double-blind clinical study. *Int J Impot Res.* 2012;25(2):45-50.
50. Russo A, Maisto E, Romis L, Celentano G. Use of a Natural Compound Made of *Ecklonia bicyclis* Seaweed, *Tribulus terrestris*, and Water-Soluble Chitosan Oligosaccharide, in Male Sexual Asthenia with Mild or Mild-Moderate Erectile Dysfunction and Serum Testosterone Levels at the Lower Limit of Norm. *Health (Irvine Calif).* 2016;8(15):1668-78.
51. Tortora GJ, Derrickson BH. Principles of anatomy and physiology. John Wiley & Sons; 2008.
52. Cooke P. Overview of Spermatogenesis and Steroidogenesis: John Wiley & Son.