



COMPUTED QUANTUM CHEMICAL MODELING OF THE EFFECT OF NANOSILVER ON CORONAVIRUS COVID-19

Veronika Viktorovna Mishununa¹, Mansur Musaevich Chapanov², Kamilla Imranovna Gakaeva², Madina Beslanovna Tsoroeva³, Salima Alberdovna Kazanova⁴, Mariya Igorevna Gorlova⁵, Anastasiya Aleksandrovna Blinova⁶, Anna Aleksandrovna Remizova⁷, Alexey Alekseevich Gvozdenko⁶, Alexey Borisovich Golik⁶, Daniil Remizov⁶, Artem Evgenevich Mishvelov^{8*}, Marina Petrovna Marinicheva⁹, Sergey Nikolaevich Povetkin⁵, Egor Leonidovich Demchenkov¹⁰

1. Faculty of Medical, Saint-Petersburg Medico-Social Institute, Saint-Petersburg, Russia.
2. Department of Therapy, Faculty General Medicine, Chechen State University, Grozny, Chechen Republic, Russia.
3. Faculty of Medical, North-Western State Medical University named after I.I. Mechnikov, Saint-Petersburg, Russia.
4. Faculty of Medical, Kabardino-Balkarian State University named after H.M. Berbekov, Nalchik, Kabardino-Balkarian Republic, Russia.
5. Faculty of Medical, Stavropol State Medical University, Stavropol, Russia.
6. Department of Physics and Technology of Nanostructures and Materials, Faculty of Physics and Technology, North Caucasus Federal University, Stavropol, Russia.
7. Department of Dentistry, Faculty of Dentistry, North Ossetian State Medical Academy, Vladikavkaz, Republic of North Ossetia, Russia.
8. Laboratory of 3D technologies, Stavropol State Medical University, Stavropol, Russia.
9. Department of Animal disease and VSE, Saratov State Agrarian University named after N.I. Vavilov, Saratov, Russia.
10. FreezeM Inc, Kibbutz Nahshonim, Israel.

ARTICLE INFO

Received:

27 Dec 2020

Received in revised form:

09 Mar 2021

Accepted:

14 Mar 2021

Available online:

28 Apr 2021

Keywords: COVID-19, Nanosilver, Quantum-chemical modeling, HOMO, LUMO

ABSTRACT

This article presents an analysis of available scientific data on the morphology and nanostructure of the COVID-19 coronavirus. Possible mechanisms of influence of nanosilver particles on the coronavirus are considered. Models of nanosilver complexes with spike protein of coronavirus amino acids were constructed using computer quantum-chemical modeling. The values of electron density distribution, highest occupied molecular orbital, lowest unoccupied molecular orbital, and electron density distribution gradient for each constructed model are obtained. As a result of quantum chemical modeling, it was found that silver nanoparticles can interact with the following amino acids: Proline, glutamine, lysine, arginine, asparagine, histidine, glutamic and aspartic acids, tryptophan, and cysteine, which is due to the presence of additional $-NH_2$, $-NH$, $-SH$ and $-COOH$ groups in these amino acids that are not involved in the formation of a peptide bond. The freedom of additional groups makes it possible to interact with nanosilver. Analysis of the obtained data showed that the most energy-efficient interaction is the formation of the "tryptophan-nanosilver" complex ($E = -5856.83$ kcal/mol). Based on the findings of quantum chemical calculations, the most stable complex is the "cysteine-nanosilver" ($\Delta E = 0.16$ a.u.).

This is an *open-access* article distributed under the terms of the *Creative Commons Attribution-Non Commercial-Share Alike 4.0 License*, which allows others to remix, tweak, and build upon the work non commercially, as long as the author is credited and the new creations are licensed under the identical terms..

To Cite This Article: Mishununa VV, Chapanov MM, Gakaeva KI, Tsoroeva MB, Kazanova SA, Gorlova MI, et al. Computed Quantum Chemical Modeling of the Effect of Nanosilver on Coronavirus COVID-19. *Pharmacophore*. 2021;12(2):14-21. <https://doi.org/10.51847/LcTdy7pSqE>

Introduction

2020 was a real epic of medical feats and testing of our research and development [1-6]. However, the most important news

Corresponding Author: Artem Evgenevich Mishvelov; Laboratory of 3D technologies, Stavropol State Medical University, Stavropol, Russia. E-mail: ruslankalmykov777@yandex.ru.

of this year was the pandemic caused by the coronavirus.

Coronavirus disease (COVID-19) is a worldwide pandemic influencing more than 60 million people and has immensely affected healthcare networks globally [7-10]. On March 11, 2020, the World Health Organization (WHO) officially identified this public health problem as a global crisis [11]. Due to the massive redirection of medical care and priority towards the COVID-19 infected patients, the surgical arena has been affected. The economic requirements of the mentioned universal problem are rising, and they are being intensified by the demand for surgical proceedings [12].

The COVID-19 pandemic has forced the scientific world to expand its knowledge of this type of virus. Coronaviruses are a set of viruses covered with an unsegmented genome consisting of a single plus-chain RNA. Severe Acute Respiratory Syndrome (SARS-CoV) and Middle East Respiratory Syndrome (MERS-CoV) coronaviruses are highly pathogenic zoonotic coronaviruses that have caused zonal and universal prevalence at different times [13, 14].

Currently, coronavirus infection has become the number one target for scientific and medical research in the world because of its very fast distribution. Most authors are inclined to develop effective antiviral agents. In general, many interesting areas have been covered in recent years. Some scientists suggest using active phytopreparations in food to prevent viruses [15-17]. There are works on the study of the venom activity of migmamorphic spiders [18], which concluded that the polypeptide combinations of the studied poisons have strong antiviral activity. The effect of other biologically active substances on antiviral activity has also been studied [19-21]. Of particular importance for the formation of a significant immune response to various viral particles is also the normalization of the intake of essential nutrients into the body. The most important of which is the essential trace element zinc. The most noteworthy is the highly biologically active colloidal hellate forms presented in the following works [1, 22-25]. It is also worth noting the development of digitalization of medical wards [3, 26] of computer simulation of operations and the use of an electronic pharmacologist in postoperative therapy [2]. The development of special antiseptics and disinfectants is also an important factor in the fight against new types of viruses [27-29].

A large number of works indicates a serious study of the issue of combating COVID-19. However, there is no definitive solution yet. To understand which of the areas of the fight against coronavirus has great prospects, it is necessary to analyze its morphology and microstructure.

Based on the International Committee on Taxonomy of Viruses coronavirus is a *Nidovirales*, family *Coronaviridae*, subfamily *Coronavirinae* [30]. According to primary serological and subsequent genomic researches, representatives of *Coronavirinae* are classified into 4 genera: α -, β -, γ - and δ -coronaviruses [31]. Four types of viruses (A, B, C, and D) belong to the genus β -coronaviruses [32]. Among the 6 familiar human coronaviruses (HCoV), HCoV-229E and HCoV-NL63 are included in the genus α -coronaviruses, while HCoVOC43 and HCoV-HKU1 are included in type A, SARS-CoV in type B, and MERS-CoV in type C of the genus β -coronaviruses [33].

The illustrative diagram indicating the genome composition of 6 familiar HCoV (not at scale) shows the 5'-cap composition and 3'-end polyadenylation (AnAOH-3'). Open reading frames (ORF1a and ORF1b) are indicated as short red rectangles. Genes encoding structural spike proteins (S), envelope (E), membrane (M), nucleocapsid (N), and hemagglutinin esterase (HE) are presented as blue rectangles. Genes encoding auxiliary proteins are indicated as grey rectangles [34].

Having a diameter of 80-120 nm., coronaviruses are spherical or pleomorphic. The surface of the virion is dotted with club-shaped protrusions formed by triangular spines (S) made of glycoprotein. Shorter protrusions consisting of the dimeric hemagglutinin esterase (HE) protein are seen in a number of β -coronaviruses, including HCoV-OC43 and HCoV-HKU1 [35]. S and HE are types I transmembrane proteins with a long extracellular domain and a short intracellular domain. The coating of the virus is formed by a membrane glycoprotein (M), a structural protein that is embedded in the coating and includes 3 transmembrane areas [36]. In addition, the nucleocapsid (N) contains a small amount of a transmembrane protein with a relatively small molecular weight, known as envelope protein (E). Eventually, the nucleocapsid (N) protein binds to the RNA genome as "beads on a string", forming a helically symmetric structure [36].

The coronavirus genome is an unsegmented genome consisting of a single plus-chain RNA of an astonishingly big size (from 27 to 32 thousand base pairs). Genomic RNA that has a 5' cap and a poly (A) tail at the 3' end includes many Open Reading Frames (ORF). The genes are arranged in the following order: 5' - replicase-S-E-M-N-3' with various little ORF (encoding auxiliary proteins) scattered among the structural genes. The coronavirus replicase is encoded by two large overlapping ORF (ORF1a and ORF1b), which occupy about two-thirds of the genome and are straightly translated from genomic RNA. However, structural and auxiliary genes are translated from subgenomic RNAs (sgRNA) that are synthesized while genome transcription/replication, as explained below. The coronavirus replication cycle is classified into various stages:

- * Attaching and embedding in the cell;
- * Broadcast of viral replicase;
- Transcription and replication of the genome;
- * Translation of structural proteins;
- * Virion assembly and release.

Coronavirus duplication is started by the binding of the S-protein to the cell membrane receptor(s). The S-protein consists of two functional subunits: S1 ("bulb") for binding to the receptor and S2 ("stem") for fusing to the membrane. The special

interplay between S1 and a special receptor causes a sharp structural alteration in the S2 subunit, resulting in the merger of the viral coating with the cell membrane and the penetration of the nucleocapsid into the cytoplasm [34, 36]. Interaction with the receptor is a determining factor in the tissue tropism of the coronavirus to human target cells. Some HCoV use cell membrane enzymes as receptors: HCoV-229E-aminopeptidase N, HCoV-NL63, and SARS-CoV-angiotensin-converting enzyme 2 (ACE-2), MERS-CoV-dipeptidyl peptidase 4; HCoV-OC43 and HCoV-HKU1 use 9-O-acetylated sialic acid as a receptor [37].

The S1 and S2 subunits of the coronavirus S-protein are cleaved by one or more host cell proteases. For example, activity of the SARS-CoV S-protein needs consecutive division by the endosomal cysteine protease cathepsin L and another trypsin-like serine protease [38]. However, the MERS-CoV S-protein includes two division areas for furin, which is a ubiquitously expressed protease. Surprisingly, the S1/S2 site is cleaved within MERS-CoV S-protein synthesis, while the other site (S2') is cleaved within virus penetration. Previously, researchers discovered a similar cleavage process in the Infectious Bronchitis Virus (IBV) or avian coronavirus and the typical gamma-coronavirus that infects chickens [39]. Moreover, type II transmembrane serine proteases (TMPRSS2 and TMPRSS11D) are involved in the activation of SARS-CoV and HCoV-229E S-proteins. In addition to S-protein activity, host cell elements can also be involved in next phases of virus entrance. For instance, a valosin-comprising protein promotes the release of coronavirus from early endosomes, since a decrease in the amount of this protein leads to a decrease in the duplication of both HCoV-229E and IBV [14].

Host cell factors can also limit the addition and penetration of HCoV. For instance, Interferon-Induced Transmembrane proteins (IFITM) have broad-spectrum antiviral effects against different RNA viruses. IFITM also restricts the entrance of SARS-CoV, MERS-CoV, HCoV-229E, and HCoV-NL63 into the cell. In contrast, HCoV-OC43 uses IFITM2 or IFITM3 as an implementation element. A late research recognized many amino acid residuals in IFITM that control the intensity of HCoV introduction [40].

A detailed analysis of the morphology of the coronavirus, its structure, and microstructure of the nanocomposites RNA leads to the conclusion that the most effective tool in the fight against coronavirus may be the use of nanoscale silver particles exhibiting acute bactericidal (**Figure 1**) and antiviral activity [41, 42].




Figure 1. Mechanism of the Toxic Effect of Nanosilver on Bacterial Cells.

The mechanism of toxic action of nanosilver on bacterial cells and viruses is possible in several ways: interaction and damages to cell membranes, cellular uptake, reactive oxygen species (ROS) production, interaction with and damage to cellular proteins, binding and damages to cellular DNA and RNA repair [43]. In the case of coronavirus, there may be 3 directions of nanosilver exposure: membrane damage, RNA disintegration, and spike protein destruction [44]. The effectiveness of each type of toxic effect of nanosilver on coronavirus has yet to be determined in numerous experiments, which will be based on models of the mechanism of action.

The purpose of this work was to conduct computer quantum-chemical modeling of the mechanism of action of nanoscale silver particles on the morphology and micro- and nanostructure of coronavirus.

Materials and Methods

In the course of the work, integration with Protein Data Bank of Research Collaboratory for Structural Bioinformatics (RCSB PDB) was performed for accurate modeling of HCoV spike protein sections (**Figure 2**) and evaluation of the most affected nanosilver nodes.




Figure 2. Prefusion Structure of a Human Coronavirus Spike Protein (Source RCSB PDB <https://www.rcsb.org/3d-view/5I08>).

Computer quantum-chemical modeling of the process of silver nanoparticle's effect on coronavirus was performed in the

QChem program using the IQmol molecular editor [45, 46]. The calculation was performed on the equipment of the Data Processing Center (Schneider Electric) of the North Caucasus Federal University. Calculation characteristics: Energy, method – B3LYP; basis – 3-21G, convergence-5, force field-Ghemic [47-49].

Amino acids were considered as the target of nanosilver. According to the research of Tok and Tatar (2017) and Masters (2006), the size of a coronavirus can vary from 80 to 120 nm, so the average size of a coronavirus is assumed to be 100 nm [36, 50]. In this regard, depending on the size of nanosilver particles, 3 types of interaction with coronavirus are possible: membrane damage (25-50 nm), spike protein destruction (10-25 nm), and RNA disintegration (<10 nm). The proposed types of influence of nanosilver on the coronavirus are shown in **Figure 3**.




Figure 3. Schemes for the Effect of Nanosilver (Gray Particles) on Coronavirus at Different Particle Sizes: a) 25-50 nm, b) 10-25 nm, c) <10 nm

Since nanosilver can be very active with amino acids, it was suggested that the main target of nanosilver will be spike protein (**Figure 4**).




Figure 4. Mechanism of Nanosilver Effect on Spike Protein of HCoV.

The model of the effect of nanosilver on coronavirus presented in **Figure 4** is taken as the main one in this paper.

Results and Discussion

At the primer phase of study, quantum-chemical modeling of coronavirus's spike protein amino acids and nanosilver interaction was performed.

As a result of quantum chemical modeling, it was found that silver nanoparticles can interact with the following amino acids: Proline, glutamine, lysine, arginine, asparagine, histidine, glutamic and aspartic acids, tryptophan, and cysteine, which is due to the presence of additional $-NH_2$, $-NH$, $-SH$ and $-COOH$ groups in these amino acids that are not involved in the formation of a peptide bond. These free additional groups make possible interaction with nanosilver [23, 47]. Thus, the interaction of silver nanoparticles with threonine, serine, and tyrosine is unlikely. Obtained data are confirmed by the results of quantum chemical calculations (**Table 1**).

Table 1. The Results of Quantum-chemical Calculations.

Amino acid	E, kcal/mol	HOMO, a.u.	LUMO, a.u.	ΔE , a.u.
Proline	-398,553	-0,386	0,148	0,534
Proline+nanosilver	-5573,5	-0,085	-0,039	0,05
Serine	-396,703	-0,213	0,021	0,234
Serine+nanosilver	-5570,76	-0,173	-0,02	0,15

Threonine	-435,811	-0,211	0,017	0,228
Threonine+nanosilver	-5609,86	-0,167	-0,016	0,15
Cysteine	-718,033	-0,24	0,011	0,251
Cysteine+nanosilver	-5890,5	-0,199	-0,036	0,16
Tyrosine	-626,504	-0,214	-0,002	0,212
Tyrosine+nanosilver	-5800,53	-0,165	-0,031	0,13
Tryptophan	-682,279	-0,185	-0,033	0,152
Tryptophan+nanosilver	-5856,83	-0,115	-0,058	0,06
Aspartic Acid	-509,426	-0,223	-0,011	0,212
Aspartic Acid+ nanosilver	-5683,43	-0,196	-0,065	0,13
Glutamic Acid	-548,276	-0,2	-0,072	0,128
Glutamic Acid+nanosilver	-5722,28	-0,16	-0,095	0,07
Histidine	-545,675	-0,212	0,004	0,216
Histidine+nanosilver	-5720,21	-0,12	-0,03	0,09
Asparagine	-489,683	-0,229	0,004	0,233
Asparagine+nanosilver	-5664,23	-0,12	-0,037	0,08
Arginine	-603,07	-0,202	-0,004	0,198
Arginine+nanosilver	-5777,64	-0,09	-0,024	0,07
Lysine	-493,995	-0,168	-0,051	0,117
Lysine+nanosilver	-5668,58	-0,088	-0,054	0,03
Glutamine	-528,788	-0,227	0,006	0,233
Glutamine+nanosilver	-5703,33	-0,115	-0,038	0,08

Table 1 shows a decrease in the system when adding a nanosilver. i.e., the interaction of nanosilver with amino acids is an energetically beneficial process. The most energy-efficient interaction is the formation of the "tryptophan–nanosilver" complex ($E = -5856.83$ kcal/mol), but the difference in the energy of interaction of nanosilver with other amino acids is not very significant. Based on the findings of quantum chemical calculations, the most stable complex is the "cysteine–nanosilver" ($\Delta E = 0.16$ a. u.).

The obtained models of spike protein amino acids with nanosilver, electron density distributions, electron density distribution gradients, and molecular orbitals for "tryptophan–nanosilver" and "cysteine–nanosilver" are shown in **Figures 5 and 6**.




Figure 5. a) Model of "Tryptophan-Nanosilver" Molecular Complex, b) Electron Density Distribution, c) Highest Occupied Molecular Orbital HOMO, d) Lowest Unoccupied Molecular Orbital LUMO, and e) Electron Density Distribution Gradient.

The formation of the "tryptophan–nanosilver" complex has a high probability, which is due to the presence of an additional NH group in the tryptophan structure.




Figure 6. a) Model of “Cysteine-Nanosilver” Molecular Complex, b) Electron Density Distribution, c) Highest Occupied Molecular Orbital HOMO, d) Lowest Unoccupied Molecular Orbital LUMO, and e) Electron Density Distribution Gradient.

Silver nanoparticles can interact with cysteine, which is due to the presence of the SH group in the cysteine structure.

Conclusion

As a result of quantum chemical modeling, it was found that silver nanoparticles can interact with the following amino acids: Proline, glutamine, lysine, arginine, asparagine, histidine, glutamic and aspartic acids, tryptophan, and cysteine, which is due to the presence of additional $-NH_2$, $-NH$, $-SH$ and $-COOH$ groups in these amino acids that are not involved in the formation of a peptide bond. The freedom of additional groups makes it possible to interact with nanosilver. Analysis of the obtained data showed that the most energy-efficient interaction is the formation of the "tryptophan–nanosilver" complex ($E = -5856.83$ kcal/mol). Based on the findings of quantum chemical calculations, the most stable complex is the "cysteine–nanosilver" ($\Delta E = 0.16$ a. u).

Acknowledgments: The authors express their gratitude to Dr. Andrey Vladimirovich Blinov for supervising the project and reviewing and editing the article.

Conflict of interest: None

Financial support: None

Ethics statement: None

References

1. Blinov AV, Kravtsov AA, Krandievskii SO, Timchenko V, Gvozdenko AA, Blinova A. Synthesis of MnO₂ nanoparticles stabilized by methionine. *Russ J Gen Chem.* 2020;90(2):283-6.
2. Bledzhyants GA, Mishvelov AE, Nuzhnaya KV, Anfinogenova OI, Isakova JA, Melkonyan RS, et al. The effectiveness of the medical decision-making support system "electronic clinical pharmacologist" in the management of patients therapeutic profile. *Pharmacophore.* 2019;10(2):76-81.
3. Nuzhnaya KV, Mishvelov AE, Osadchiy SS, Tsoma MV, AM RS KK, Rodin IA, et al. Computer simulation and navigation in surgical operations. *Pharmacophore.* 2019;10(4):43-8.
4. Hight GY, Mishvelov AE, Melchenko EA, Nuzhnaya CV, Epanov VA, Dolgalev AA. New image modeling features for planning surgical interventions. *Res J Pharm Biol Chem Sci.* 2019;10(1):140-3.
5. Demchenkov EL, Nagdalian AA, Budkevich RO, Oboturova NP, Okolelova AI. Usage of atomic force microscopy for detection of the damaging effect of CdCl₂ on red blood cells membrane. *Ecotoxicol Environ Saf.* 2021;208:111683.
6. Luneva, A, Koshchayev, A, Nesterenko, A, Volobueva, E, Boyko, A. Probiotic potential of microorganisms obtained from the intestines of wild birds. *Int Trans J Eng Manag Appl Sci Technol.* 2020;11(12):11A12E.
7. Alshammari E. Implementing eOSCE During COVID-19 Lockdown. *J Adv Pharm Educ Res.* 2020;10(1):174-80.
8. Lestari K, Sitorus T, Instiaty SM, Levita J. Molecular docking of Quinine, Chloroquine and Hydroxychloroquine to Angiotensin Converting Enzyme 2 (ACE2) receptor for discovering new potential COVID-19 antidote. *J Adv Pharm Educ Res.* 2020;10(2):1-4.

9. Meconcelli G, Bazzoni G, Casu C. Auriculotherapy for stress management as self-help in isolation situations (COVID 19). *Int J Pharm Phytopharmacol Res.* 2020;10(3):1-2.
10. Albureikan MO. COVID-19 outbreak in terms of viral transmission and disease biocontrol by healthy microbiome. *Int J Pharm Phytopharmacol Res.* 2020;10(3):139-46.
11. World Health Organization (WHO), Corona Virus Disease 2019 (COVID-19) Situation Report-51, Available from: (March 11 2020): Last accessed May 12 2020 https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200311-sitrep-51-covid-19.pdf?sfvrsn=1ba62e57_10.
12. Centers for Medicare & Medicaid Services (CMS), Adult Elective Surgery and Procedures Recommendations: Limit All Non-essential Planned Surgeries and Procedures, Including Dental, until Further Notice, (April 7, 2020) Available from: <https://www.cms.gov/files/document/covid-elective-surgery-recommendations.pdf>, Accessed date: 13 May 2020
13. Tang B, Wang X, Li Q, Bragazzi NL, Tang SY, Xiao YN, et al. Estimation of the transmission risk of the 2019-nCoV and its implication for public health interventions. *J Clin Med.* 2020;9(2):462. doi:10.3390/jcm9020462
14. Yang Z, Zeng Z, Wang K, Wong SS, Liang W, Zanin M, et al. Modified SEIR and AI prediction of the epidemics trend of COVID-19 in China under public health interventions. *J Thorac Dis.* 2020;12(3):165-74.
15. Oboturova NP, Nagdalian AA, Povetkin SN. Adaptogens instead restricted drugs research for an alternative Items to doping in sport. *Res J Pharm Biol Chem Sci.* 2018;9(2):1111-6.
16. Nesterenko A, Koshchaev A, Kenijz N, Akopyan K, Rebezov M, Okuskhanova E. Biomodification of meat for improving functional-technological properties of minced meat. *Res J Pharm Biol Chem Sci.* 2018;9(6):95-105.
17. Omarov RS, Shlykov SN, Nesterenko AA. Obtaining a biologically active food additive based on formed elements blood of farm animals. *Res J Pharm Biol Chem Sci.* 2018;9(6):1832-9.
18. Nagdalian AA, Pushkin SV, Povetkin S, Nikolaevich K, Egorovna M, Marinicheva MP, et al. Migalomorphic spiders venom: extraction and investigation of biological activity. *Entomol Appl Sci Lett.* 2018;5(3):60-70.
19. Lopteva MS, Povetkin SN, Pushkin SV, Nagdalian AA. 5% Suspension of albendazole echinacea magenta (echinacea purpurea) toxicometric evaluation. *Entomol Appl Sci Lett.* 2018;5(4):30-4.
20. Sizonenko MN, Timchenko LD, Rzhepakovskiy IV, Piskov SI, Areshidze DA, Mikhailenko VV, et al. The new efficiency of the «Srmp» – listerias growth-promoting factor during factory cultivation. *Pharmacophore.* 2019;10(2):85-8.
21. Kenijz NV, Koshchaev AG, Nesterenko AA, Omarov RS, Shlykov SN. Study the effect of cryoprotectants on the activity of yeast cells and the moisture state in dough. *Res J Pharm Biol Chem Sci.* 2018;9(6):1789-96.
22. Nagdalian AA, Oboturova NP, Povetkin SN, Ahmadov VT, Karatunov VA, Gubachikov AZ, et al. Insect's biomass as a livestock feed. study of the impact of insectoprotein on the livestock vitals. *Pharmacophore.* 2020;11(1):27-34.
23. Blinov AV, Gvozdenco AA, Yasnaya MA, Golik AB, Blinova AA, Shevchenko IM, et al. Effect of synthesis parameters on dimensional characteristics of fe3o4 nanoparticles: neural-network research. *Phys Chem Aspects Study Clust Nanostruct Nanomater.* 2019;11:298-306.
24. Barabanov PV, Gerasimov AV, Blinov AV, Kravtsov AA, Kravtsov VA. Influence of nanosilver on the efficiency of *Pisum sativum* crops germination. *Ecotoxicol Environ Saf.* 2018;147:715-9. doi:10.1016/j.ecoenv.2017.09.024
25. Selimov MA, Nagdalian AA, Povetkin SN, Statsenko EN, Kulumbekova IR, Kulumbekov GR, et al. Investigation of CdCl₂ Influence on red blood cell morphology. *Int J Pharm Phytopharmacol Res.* 2019;9(5):8-13.
26. Hite GJ, Mishvelov AE, Melchenko EA, Vlasov AA, Anfinogenova OI, Nuzhnaya CV, et al. Holodoctor planning software real-time surgical intervention. *Pharmacophore.* 2019;10(2):1-12.
27. Morozov VY, Kolesnikov RO, Chernikov AN. Effect from aerosol readjustment air environment on productivity and biochemical blood parameters of young sheep. *Res J Pharm Biol Chem Sci.* 2017;8(6):509-14.
28. Saleeva IP, Morozov VYu, Kolesnikov RO. Disinfectants effect on microbial cell. *Res J Pharm Biol Chem Sci.* 2018;9(4):676-81.
29. Blinova A, Blinov A, Baklanova O, Yasnaya M, Baklanov I, Siddiqui A, et al. Study of wound-healing ointment composition based on highly dispersed zinc oxide modified with nanoscale silver. *Int J Pharm Phytopharmacol Res.* 2020;10(6):237-45. doi:10.51847/Cu6KX0JDE3
30. Hamre D, Procknow JJ. A new virus isolated from the human respiratory tract. *Exp Biol Med.* 1966;21(1):190-3.
31. Muradyan AY, Osidak LV, Rumel NB, Korenyako IE. Significance of coronavirus infection in acute respiratory pathology in children. *Child Infect.* 2003;3:22-5.
32. Profeta A, Siddiqui S, Smetana S, Hossaini M, Hieke S, Enneking U, et al. The Impact of Corona Pandemic on Consumer's Food Consumption–Vulnerability of Households with Children and Income Losses. 2021. doi:10.20944/preprints202101.0153.v2
33. Chen B, Tian EK, He B, Tian L, Han R, Wang S, et al. Overview of lethal human coronaviruses. *Signal Transduct Target Ther.* 2020;5(1):89. doi:10.1038/s41392-020-0190-2
34. Fung TS, Liu DX. Human Coronavirus: Host-Pathogen interaction. *Annu Rev Microbiol.* 2019;73(1):529-57. doi:10.1146/annurev-micro-020518-115759
35. de Groot RJ. Structure, function, and evolution of the hemagglutinin-esterase proteins of corona and toroviruses. *Glycoconj J.* 2006;23(1-2):59-72.
36. Masters PS. The molecular biology of coronaviruses. *Adv. Virus Res.* 2006;66:193-92.

37. Lim Y, Ng Y, Tam J, Liu D. Human coronaviruses: a review of virus-host interactions. *Diseases*. 2016;4(4):26.
38. Belouzard S, Chu VC, Whittaker GR. Activation of the SARS coronavirus spike protein via sequential proteolytic cleavage at two distinct sites. *PNAS*. 2009;106(14):5871-6.
39. Zimmerman T, Siddiqui SA, Bischoff W, Ibrahim SA. Tackling airborne virus threats in the food industry: a proactive approach. *Int J Environ Res Public Health*. 2021;18(8):4335.
40. Zhao X, Sehgal M, Hou Z, Cheng J, Shu S, Wu S, et al. Identification of residues controlling restriction versus enhancing activities of IFITM proteins on entry of human coronaviruses. *J Virol*. 2018;92(6):e01535-17.
41. Blinov AV, Siddiqui SA, Nagdalian AA, Blinova AA, Gvozdenko AA, Raffa VV, et al. Investigation of the influence of Zinc-containing compounds on the components of the colloidal phase of milk. *Arab J Chem*. 2021;14(7):103229.
42. Lunin, LS, Lunina, ML, Kravtsov AA, Blinov AV. Effect of the Ag nanoparticle concentration in TiO₂-Ag functional coatings on the characteristics of GaInP/GaAs/Ge photoconverters. *Semiconductors*. 2018;52(8):993-6. doi:10.1134/S1063782618080122
43. Guo Z, Zeng G, Cui K. Toxicity of environmental nanosilver: mechanism and assessment. *Environ Chem Lett*. 2019;17(1):319-33. doi:10.1007/s10311-018-0800-1
44. Qurat-Ul-Ain, Sarfraz RA, Qayyum A. Mechanism of action of bio-inspired nanosilver particles. *Bioinspired, Biomim Nanobiomaterials*. 2018;7(3):174-86. doi:10.1680/jbibn.17.00026
45. Blinov AV, Gvozdenko AA, Kravtsov AA, Krandievsky SO, Blinova AA, Maglakelidze DG, et al. Synthesis of nanosized manganese methahydroxide stabilized by cysteine. *Mater Chem Phys*. 2021;2651:124510.
46. Nagdalian AA, Rzhepakovsky IV, Siddiqui SA, Piskov SI, Oboturova NP, Timchenko LD, et al. Analysis of the content of mechanically separated poultry meat in sausage using computing microtomography. *J Food Compos Anal*. 2021;100:103918. doi:10.1016/j.jfca.2021.103918
47. Blinov AV, Yasnaya MA, Blinova AA, Shevchenko IM, Momot EV, Gvozdenko AA, et al. Computer quantum-chemical simulation of polymeric stabilization of silver nanoparticles. *Phys Chem Aspects Study Clust Nanostruct Nanomater*. 2019;11:414-21.
48. Gvozdenko AA, Blinov AV, Yasnaya MA, Golik AB, Raffa VV, Kramarenko VN, et al. Computer quantum-chemical simulation of multicomponent SiO₂-MexO_y systems. *Phys Chem Aspects Study Clust Nanostruct Nanomater*. 2020;12:394-404.
49. Yasnaya MA, Blinov AV, Blinova AA, Shevchenko IM, Maglakelidze DG, Senkova AO. Determination of optimal modes for measuring the size of colloidal particles by photon-correlation spectroscopy and acoustic spectroscopy. *Phys Chem Aspects Study Clust Nanostruct Nanomater*. 2020;12:232-42.
50. Tok TT, Tatar G. structures and functions of coronavirus proteins: molecular modeling of Viral Nucleoprotein. *Int J Virol Infect Dis*. 2017;2(1):001-7.