

Pharmacophore

(An International Research Journal)

Available online at <http://www.pharmacophorejournal.com/>

Original Research Paper

ANTIBACTERIAL POTENTIAL OF BISEHRI BOOTI (*AREVA LANATA* LINN. JUSS.)

Najmuddin Ahmad Siddiqui, Ghufraan Ahmad, **Sumbul Rehman*** and Asma Abid

Department of Ilmul Advia (Unani Pharmacology & Pharmaceutical Sciences),
Ajmal Khan Tibbiya College & Hospital, Aligarh Muslim University, Aligarh, India

ABSTRACT

Bisehri booti (*Aerva lanata* (Linn.) Juss.; Family Amaranthaceae) has been used in traditional systems of medicine since long time to treat many diseases including infectious diseases. However, information to revalidate its use on scientific parameters is limited. Therefore, present study has been done to screen antibacterial activity of ethanolic extracts of Bisehri booti against pathological bacterial strains using disk diffusion method and its efficacy was compared with the standard drug ciprofloxacin for Gram-positive and gentamycin for Gram-negative bacterial strains. Qualitative analysis of the extracts was also done to confirm the presence of various phytoactive constituents. It was found that Bisehri booti contains alkaloids, glycosides, flavonoids, carbohydrates, starch, saponin, resins and terpenes. This study scientifically validates the antibacterial potential of Bisehri booti and reconfirms the claims of Unani physicians for its use in various infectious diseases as urinary tract infections, leprosy, dysentery, typhoid, wounds etc. The findings can usefully be applied by researchers elsewhere for comparison of its bioactive constituents and this might be a helpful lead in the rational use of Bisehri booti.

Keywords: Infectious diseases, Bisehri booti (*Aerva lanata* (Linn.) Juss.), Antibacterial screening.

INTRODUCTION

Unani system of medicine provides a large number of drugs of natural origin which can be used today to combat many dreadful diseases which are difficult to be treated with the western medicine along with the emerging health hazards nowadays. Ethno-botanical, medical and phytochemical literature available in our classical books has abundant remedies from natural resources as mentioned by eminent Unani physicians that need to be re-explored now for different health hazards. Unani medicine claims to possess a large number of effective and safe drugs to combat infectious diseases that are in use since centuries (Rehman and Latif, 2015). Bacterial strains have developed resistance to almost all the antibiotics, further some antibiotics have serious undesirable effects (Rehman *et al.*, 2011). This is an alarming situation and calls for serious consideration including drugs available in

indigenous system of medicines. There appears no effort made to investigate the antimicrobial potential of 'Bisehri booti' through in-vitro testing. Present study is based on this rationale and presents results of phyto-chemical and antibacterial screening of this important Unani drug in an effort to revalidate its medical efficacy to combat infectious diseases.

Bisehri Booti

Bisehri booti (*Aerva lanata* (Linn.) Juss.; Family Amaranthaceae) commonly known as mountain knotgrass is a woody, prostrate or succulent perennial herb of 30-60 cm in height. It is a native of Asia, Africa, Australia, Sri Lanka, South Asia, Saudi Arabia, Egypt, Java and Philippines (Lakshmi and Lethi, 2014) grows throughout the hotter plain parts of India as common weed. The plant is villous or pubescent, prostrate dioecious herb 1-4 ft. height, having a tap root which is

cylindrical, branches arising from the stem or root stock in old plants 7-12 cm long, 2-8 mm thick, straight or slightly twisted with many slender, fibrous lateral roots, pale yellowish brown externally, whitish internally, camphoraceous odorous, it has many branches, branched from the root base; pubescent/wooly-tomentose, striate (Thiselton, 1963; Trimen, 1974; Nagaratna *et al.*, 2014). It is used to treat a number of diseases as haematuria, albuminuria, kidney disorders, jaundice, biliousness, in cases of snake bite, used to treat dyspepsia, pneumonia, typhoid and in cases of Herpes in Orissa (Singh and Pandey, 1998; Apai, 2009; Indukuri *et al.*, 2013), for arresting haemorrhage during pregnancy (Yoganarasimhan *et al.*, 1979), burn healing (Upadhaya *et al.*, 1998), anthelmintic and diuretic (Tripathi *et al.*, 1996), antidysentric (Sahoo and Mudgal, 1993); anti-inflammatory, in headache, skin disease (Singh and Pandey, 1980), to dissolve kidney and gallbladder stones (Sudhaker and Chetty, 1998), for uterus clearance after delivery (John, 1984). The flowers are used in dysentery, diarrhea (Sudhaker and Chetty, 1998) and bronchitis (Shah and Gopal, 1985). The leaves are used as anti-malarial and antipyretic (Ahmad, 1995; Dager and Dager, 1991) and expel stone from kidney (Kumar and Pullaiah, 1998), roots are used in headache, scabies, cough (Bedi, 1978); as a demulcent and diuretic (Tripathi *et al.*, 1996); to cure diarrhoea (Mohanty *et al.*, 1996); to treat nasal bleeding, cough, as an antidote for scorpion stings, fractures, spermatorrhoea, to clear uterus after delivery and also to prevent the lactation (Sikarwar and Kaushik, 1993; Rajeswari *et al.*, 2012); in urinary troubles (Hemadri *et al.*, 1980) and as antirheumatic (Kakrani and Saluja, 1994).

MATERIAL AND METHODS

The study was undertaken in the department of Ilmul Advia, A. K. Tibbiya College, Aligarh during 2013-2014.

Collection of Plant Material

The whole plant of Bisehri booti (*Aerva lanata* Linn.) was collected from the campus of Ajmal Khan Tibbiya College (AKTC), Aligarh Muslim

University (AMU) Aligarh and was properly identified by the available literature and in the Pharmacognosy section of the department of Ilmul Advia. A Voucher specimen (SC-0140/13S) of the plant material has also been submitted to the Department of Ilmul Advia, Faculty of Unani Medicine, AMU, Aligarh for the purpose of record and future reference (Figure 1).

Preparation of Ethanolic Extracts

The test drug was dried at room temperature in a ventilated room, milled to fine powder and stored in a close air tight container in dark until use. Extraction was done according to the method described by Afaq *et al.* (1994) and Peach and Tracey (1955) with some minor modifications, keeping in mind that the thermo labile elements present in the drug are destroyed when exposed to a higher temperature beyond 55°C, so the heat wherever needed was kept as low as possible to prevent the loss of thermo-labile substances present in the drugs from destruction. The coarse drug material was extracted with 95% ethanol as a solvent at 50°C for 6 hours and dried under reduced pressure in the Lypholizer. Strict aseptic precautions were followed throughout the process. The stock solutions for extract were prepared from the dried extract so obtained in the Dimethyl Sulphoxide (DMSO) as a solvent for use. The respective stock solutions so prepared were refrigerated till further use.

Phytochemical Analysis

Phytochemical studies of the plant preparations are necessary for standardization, which helps in understanding the significance of phyto-constituents in terms of their observed activities. Phytochemistry also helps in standardizing the herbal preparations so as to get the optimal concentrations of known active constituents, and in preserving their activities (Rehman and Latif, 2015). Qualitative analysis of the chemical constituents present in the drug sample (Bhattacharjee and Das, 2005 and Afaq *et al.*, 1994). Standard tests and methodology has been followed to confirm the presence of chemical constituents (Table-1).

Antibacterial Susceptibility Testing

Antimicrobial susceptibility testing was done by Kirby Bauer's disk diffusion method (Bauer *et al.* 1996) as was performed on biofilm isolates by Kirby-Bauer on Muller Hinton Agar (Barry *et al.*, 1999). Bacterial strains used for the study are listed in Table-2. The standard medium Mueller Hinton Agar, was poured to a depth of 4 mm in a 90 mm petridish (PW008, Himedia Labs Pvt. Ltd., Mumbai, India). The plate was inoculated by streaking the entire surface in three planes with a sterile cotton swab (PW041, Himedia Labs Pvt. Ltd., Mumbai, India) dipped into standardized inoculums, spreaded evenly with the help of L-Spreader (PW1085, Himedia Labs Pvt. Ltd., Mumbai, India). The bacterial inoculum was prepared from an 18 hour broth culture of the microbe to be tested and was standardized with sterile physiologic saline to contain 10^6 cfu/ml. Standardized commercial paper disk containing amounts of the antimicrobial agents to be tested were placed on the surface of the agar. The plate was incubated in an inverted position at 37°C for 18 hours. The diameter of zone of inhibition produced by the drug was measured with the help of Antibiotic zone scale (PW297, Himedia Labs Pvt. Ltd., Mumbai, India) from each disk (Kingsbury and Wagner, 1990).

Statistical Analysis

The statistical analysis was done using gpaidd software, One way ANOVA and the post test named Bonferroni: Selected pairs of column with multiple comparison was performed with p-value <0.05.

RESULTS AND DISCUSSION

Qualitative analysis of the phyto-chemicals of Bisehri booti was carried out for the determination of presence of alkaloids, amino acids, proteins, carbohydrates, flavonoids, glycosides, tannins, sterols, phenols, resins and volatile oil. As the therapeutic properties of the crude drugs are mainly due to physiologically active chemical constituents present in the drugs and the lower percentage of chemical constituents may cause lesser therapeutic values of the drugs and therefore, they are considered as low standard drugs. Moreover the chemical constituents present

in the drug have documented antimicrobial activities. Tannins have the property of precipitating proteins. By an analogous process they prevent the development of bacteria and since the proteins are necessary for their growth and development so, they not only stop their nutrition but also precipitate their own proteins (Said, 1996). The phytochemical analysis of the chemical constituents present in the test drug revealed that it contains alkaloids, phenol, resins, saponins, sugars and tannins. These were the encouraging results for our study as such types of chemical compounds are usually responsible for the therapeutic efficacy of that drug. As can be realized from the fact that alkaloids possess antimicrobial and anti-inflammatory activity, this effect has been confirmed by us in our in-vitro study of antimicrobial screening that the drugs were found to be sensitive to many microbes.

This study also confirms the presence of saponin by qualitative test, as they are generally considered as the soapy substances that are general cleansers, having antiseptic properties (Hirat and Suga, 1983). Sterols either decrease the activity of *S. aureus*, *E. coli*, *P. vulgaris* and *Pseudomonas pyocyanea* or have no effect in case of *Klebsiella* and *S. dysenterica* (Anuradha and Goyal, 1995). Flavonoids possess anti-inflammatory, anti-spasmodic, antibacterial, antifungal and antitumor activity, they act as an anti-oxidant and are also thought to protect plants from UV radiations and micro-organisms (Cushnie and Lamb, 2005). These are few evidences in support of the therapeutic activity of the Unani drugs. Most of these findings were found to be helpful in showing its biological activity.

In an effort to validate the antibacterial efficacy of selected test drug 'Bisehri booti' all ethno-pharmacological knowledge was found to be in favor of our selection of the drug as per the guidelines by Cos *et al.* (2006) in which he described that how we can develop a stronger in vitro proof-of-concept in determining anti-infective potential of natural products. The literature survey revealed that an ample data is available regarding the authentication of Unani

drugs to be used in infectious diseases, however, very little assay has been done on Unani drugs regarding their antimicrobial activity.

Present study is a step ahead in this regard to scientifically validate the antimicrobial activity of selected test drug viz. Bisehri booti using Kirby Bauer's disk diffusion and Agar well diffusion method against different Gram positive and Gram Negative bacterial strains. A total volume of 40 µl of test drugs from concentrations viz. 40.0 µg/ml was used and compared with the standard drug Ciprofloxacin (30µg) for Gram positive and Gentamicin (30 µg) for Gram Negative bacteria and Plane control i.e. DMSO- (Dimethyl Sulphoxide) the solvent used (to exclude out any activity due to the solvent used to dissolve the plant extract). There was an increased inhibitory activity against most of the strains. Among Gram positive strains *B.cereus* (18.8 ± 0.58) > *S.epidermidis* (18.0 ± 0.89) > *S.aureus* (16.2 ± 1.20) > *S.mutans* (16.0 ± 0.54) > *S.pyrogenes* (7.8 ± 0.37) while it was completely resistant to *C.xerosis* ATCC 373 at all concentration. For Gram negative strain used it showed sensitivity to all strains and there was significant inhibitory activity in the order of *E.coli* (22.6 ± 0.24) > *P.aeruginosa* (13.8 ± 1.31) > *P.vulgaris* (9.4 ± 0.67) > *K.pneumoniae* (9.2 ± 0.96). All

showed a significant inhibition as compared to Gentamicin (ZOI- 14.0-14.8 mm).

CONCLUSION

The present study provides the first report to ameliorate the efficacy of 'Bisehri booti' in infectious diseases on scientific parameters. A large number of active phytoconstituents present in the drug have been found obvious to check the growth of bacteria and treat the disease. The study concludes bisehri booti has a potent antibacterial activity against many pathogenic bacterial strains and thus could be used to derive antibacterial agents to fight against number of infectious diseases. Among the gram positive strains it was found to be sensitive to all the tested bacterial strains viz. *S. aureus*, *S. mutans*, *S. epidermidis*, *S. pyogenes*, *B. cereus*, *E. coli*, *K. pneumoniae*, *P. aeruginosa* and *P. vulgaris* while it showed a complete resistance to *C. xerosis*. Most significant sensitivity was shown towards *E.coli*, *B.cereus*, *S. epidermidis* and *S.aureus*.

ACKNOWLEDGEMENT

Authors are thankful to DRS-I (SAP-II) UGC, Department of Ilmul Advia, Ajmal Khan Tibbiya College Hospital, A.M.U., Aligarh, for providing all the financial assistance for this study.

Table 1: Qualitative Analysis of the Phytochemicals

S. No.	Chemical Constituents	Test Reagents	Inference
			Bakain (<i>M.azedarach</i> Linn.)
1.	Alkaloids	Dragendorff's reagent	+
		Wagner's reagent	+
		Mayer's reagent	+
2.	Carbohydrates	Molish Test	+
		Fehling Test	+
		Benedict Test	+
3.	Flavonoids	Mg Ribbon and dil. HCl	+
4.	Glycosides	NaOH Test	+
5.	Phenols	Ferric Chloride Test	-
		Liebermann's test	-
		Lead Acetate test	-
6.	Proteins	Xanthoproteic test	+
		Biuret test	+

7.	Starch	Iodine Test	+
8.	Saponins	Frothing with NaHCO ₃	+
9.	Steroids/ Terpenes	Salkowski Reaction	+
10.	Resins	Acetic anhydride test	-

Indications: '-' Absence and '+' Presence of constituents

Table 2: Microbial Strains Used

S. No.	Bacterial Strains	Type
1.	<i>Streptococcus mutans</i>	Gram Positive
2.	<i>Staphylococcus epidermidis</i>	"
3.	<i>Streptococcus pyogenes</i>	"
4.	<i>Bacillus cereus</i>	"
5.	<i>Staphylococcus aureus</i>	"
6.	<i>Corynebacterium xerosis</i>	"
7.	<i>Escherichia coli</i>	Gram Negative
8.	<i>Proteus vulgaris</i>	"
9.	<i>Pseudomonas aeruginosa</i>	"
10.	<i>Klebsiella pneumoniae</i>	"



Figure 1: Bisehri booti (*Areva lanata* (Linn.) Juss.)

Table 2 (a): Antibacterial Activity of Bisehri booti against Gram positive bacterial strains

S. No.	Test strains	Zone of Inhibition (in mm) expressed as Mean \pm S.E.M ^{Probability of error}		
		Drug Extract (40 μ g/ml)	Control (DMSO- 50 μ l)	Standard (Ciprofloxacin 30 μ g)
1.	<i>S.mutans</i> (ATCC 25175)	16.0 \pm 0.54* (S)	6.6 \pm 0.24(0.54) (R)	21.2 \pm 0.37(0.83) (S)
2.	<i>S.epidermidis</i> (ATCC 155)	18.0 \pm 0.89* (S)	6.4 \pm 0.24 (R)	21.6 \pm 0.24(0.54) (S)
3.	<i>S.pyrogenes</i> (ATCC 14289)	7.8 \pm 0.37*** (S)	6.4 \pm 0.24 (R)	21.2 \pm 0.37 (S)
4.	<i>B.cereus</i> (ATCC 11778)	18.8 \pm 0.58* (S)	6.6 \pm 0.24 (R)	21.4 \pm 0.24 (S)
5.	<i>S.aureus</i> (ATCC 29213)	16.2 \pm 1.20* (S)	6.6 \pm 0.24 (R)	26.8 \pm 0.20 (S)
6.	<i>C.xerosis</i> (ATCC 373)	6.6 \pm 0.24 (R)	6.6 \pm 0.24 (R)	21.2 \pm 0.37 (S)

(S) –Sensitive; (R)– Resistant

Table 2 (b): Antibacterial Activity of Bisehri booti against Gram Negative Bacterial Strains

S. No.	Test strains	Zone of Inhibition (in mm) expressed as Mean \pm S.E.M ^{Probability of error}		
		Drug Extract (40 μ g/ml)	Control (DMSO- 50 μ l)	Standard (Gentamicin 30 μ g)
1.	<i>E.coli</i> (ATCC 25922)	22.6 \pm 0.24 *** (S)	6.4 \pm 0.24 (R)	14.8 \pm 0.20 (S)
2.	<i>P.vulgaris</i> (ATCC 6380)	9.4 \pm 0.67* (S)	6.4 \pm 0.24 (R)	14.0 \pm 0.54 (S)
3.	<i>P.aeruginosa</i> (ATCC 25619)	13.8 \pm 1.31* (S)	6.4 \pm 0.24 (R)	14.8 \pm 0.20 (S)
4.	<i>K.pneumoniae</i> (ATCC 15380)	9.2 \pm 0.96*** (S)	6.4 \pm 0.24 (R)	14.8 \pm 0.20 (S)

(S) –Sensitive; (R)– Resistant

REFERENCES

1. Afaq, SH and Tajuddin, Siddiqui MMH (1994), “*Standardization of Herbal Drugs*”, AMU Publication Division, Aligarh Muslim University Press, Aligarh.
2. Anuradha, V and Goyal, MM (1995), “Phytochemical study on the leaves of *Alstonia scholaris* and their effects on pathogenic organisms”, *Ancient Science of Life*, 15 (1), 30-34.
3. Apai, KG; Rai, VK; Nandy, BC; Meena, KC; Dey, S; Tyagi, PK and Tyagi, LK, (2009), “Hyperglycaemic and anti-hyperlipidemic effect of Ethanolic extract of Aerial parts of *Aerva lanata* Linn.in

- normal and alloxan induced Diabetic rats', *Int J Pharmaceut Sci Drug Res*, Vol. 1, (3), 191-194.
4. Ahmad, A, (1995), "Drugs of plant origin as used by certain tribes of Eastern U.P. (Part-IV)", *Int J Mendel*, Vol.12, 12-14.
5. Barry, LA; Craig, AW; Nadler, H; Reller, BL; Sanders, CC and Swensor, JM (1999), "Methods for determining bacteriicidal activity of antimicrobial agents. Approved Guidelines", *Clinical and Laboratory Standard Institute (CLSI)*, Vol. 19 (18), M26-A,1-19.
6. Bauer, K; Sherris and Turck (1996), "Performance standards for antimicrobial disk susceptibility tests. CLSI (formerly NCCLS), *Am. J. Cl. Path.*, 45,493.
7. Bedi, SJ (1978), "Ethnobotany of Ratan Mahal Hills, Gujrat, India", *Eco.Bot.* 32 (3), 278-284.
8. Cos, P; Vlietinck, AJ; Berghe, DV and Maes, L (2006), "Anti-infective potential of products: How to develop a stronger in vitro 'proof of concept", *Journal of Ethnopharmacology*, 106, 290-302.
9. Cushnie, TP and Lamb, AJ (2005), "Antimicrobial activity of flavonoids", *Int. J. Antimicrob. Agents*, 26 (5), 343-56.
10. Dager, HS and Dager, JC (1991), "Plant folk medicines among the Nicobarese of Katchal islands, India", *Econ Bot.*, Vol. 45, 114-119.
11. Hemadri, K; Raj, PV; Rao, SS and Sharma, CRR (1980), "Folklore claims from Andhra Pradesh-I", *J Sci Res Plant Med.*, Vol. 1, 37-49.
12. Hirat, T and Suga, T (1983), "The efficiency of aloe plants, chemical constituents and biological activities", *Cosmetics and Toiletries*, 98, 105-108.
13. Indukuri, R; Prakash, B; Priyadarshani, RL; Vattipalli, M and Rajkumar, BP (2013), "Evaluation of Anti-ulcer activity of *Aerva lanata* stem extract in rats", *Indo Am. J Pharmaceut Res.*, Vol. 3, (12), 1702-1708.
14. John, D (1984), "One hundred useful raw drugs of the kani tribes of Trivendrum forest division, Kerala, India", *Int J Crude Drug Res*, Vol. 22, 17-39.
15. Kingsbury, DT and Wagner, GE (1990), "*Microbiology*", 2nd Edition, Harvard Publishers, U.S.A, 29-42.
16. Kakrani, HKN and Saluja, AK (1994), "Traditional treatment through herbs in Kutch district, Gujrat state, India. Part II. Analgesic, anti-inflammatory, antiarthritic plants", *Fitoterapia*, Vol.65, 427-430.
17. Kumar, VR and Pullaiah, T (1998), "Medicinal Plants used by the tribals, of Prakasam, Andhra Pradesh", *Ethnobotany*, Vol.10, 97-102.
18. Lakshmi, PP and Lethi, CD (2014), "Effect of *Scoparia dulcis* (Linn.) and *Aerva lanata* (Linn.) whole plant and fruit part extract on urine volume of ethylene glycol induced urolithoasis in male albino rats", *Int. J. Curr. Microbiol.App.Sci*, Vol. 3, (4), 1218-1223.
19. Mohanty, RB; Padhy, SN and Dash, SK (1996), "Traditional phytotherapy for diarrhoeal diseases in Gunjan and Phulbani district of South Orissa, India", *Ethnobotany*, 8, 60-65.
20. Nagaratna, A; Prakash, L; Hegde and Harini, A (2014), "A pharmacological review on Gorkha ganja (*Aerva lanata* (Linn) Juss.Ex Schult). Journal of Pharmacognosy and Phytochemistry", Vol.3 (4), 253-257.
21. Peach, K and Tracey, MV (1955), "Modern methods of Plant analysis", *Springer-Verlag*, 03, 626-627.
22. Rajeswari, DV; Vijayalakshami, S and Gajalakshami, S (2012), "Pharmacological activities of *Aerva lanata*: A prespective rewiew", *IJRP*, Vol.3, (1), 28-30.
23. Rehman, S; Latif, A; Ahmad, S and Khan, AU (2011), "In-vitro Antibacterial screening of *Swertia chirayita* Linn. against MRSA (Methicillin Resistant *Staphylococcus aureus*)", *International*

- Journal of Current Research and Review* 03 (6), 98-104.
24. Rehman, S and Latif, A (2015), "Antibacterial Screening of Karanjwa Seeds (*Caesalpinia bonducella* Roxb.): An effective Unani Medicine for Infectious diseases", *Hippocratic Journal of Unani Medicine* 10 (2), 101-109.
 25. Said, HM (1996), "Medicinal Herbs", *Hamdard Foundation Pakistan*, Pakistan, 144.
 26. Singh, V and Pandey, RP (1998), "*Ethnobotany of Rajasthan*", Scientific Publishers, Jodhpur, 38.
 27. Sahoo, AK and Mugdal, V (1993), "Ethnobotany of South Chotanagpur (Bihar)", *Bull Bot Surv India*, 35, 40-59.
 28. Singh, V and Pandey, RP (1980), "Medicinal plant-lore of the tribals of eastern Rajasthan (India)", *J Econ Tax Bot.*, Vol. 1, 137-147.
 29. Sudhakar, A and Chetty, KM (1998), "Medicinal importance of some angiospermic weeds used by the rural people of Chittor district of Andhra Pradesh, India", *Fitoterapia*, Vol. 69, 390-400.
 30. Shah, GL and Gopal, GV (1985), "Ethnomedical notes from the tribal inhabitants of North Gujrat (India)", *J Econ Tax Bot.*, Vol. 6, 193-201.
 31. Sikarwer, RLS and Kaushik, JP (1993), 'Folk Medicines of Morena district, Madhya Pradesh, India", *Int J Pharmacog.*, Vol. 31, 283-287
 32. Thiselton-Dyre, WT (1963), "*Flora of Tropical Africa*", Vol. VI., Pub. under the Authority of the State for the Colonies, L. REEVE & Co. LTD. The Oast House, Brook Ashfort, Kent, England, 39-40.
 33. Tripathi, YC; Prabhu, VV; Pal, RS and Mishra, RN (1996), "Medicinal plant of Rajasthan in Indian System of Medicine", *Ancient Sci Life.*, Vol. 15, 190-212.
 34. Trimen, H (1974), "*A Hand Book to the Flora of Cylon.*", Part III. M/S Bishen Singh Mahendra Pal Singh, New Cannaught Palace, Dehra Dun, 402-403.
 35. Upadhaya, OP; Kumar, K and Tiwari, RK (1998), "Ethnobotanical study of skin treatment uses of medicinal plants from India", *J Ethnopharmacol.*, Vol. 58, 75-83.
 36. Yoganarsimhan, SN; Bhat, AV and Togunashi, VS (1979), "Medicinal plants from Mysore district Karnataka", *Ind Drug Pharmaceut Ind.*, 14, 7-22.

Correspondence Author:

Sumbul Rehman

Department of Ilmul Advia (Unani Pharmacology & Pharmaceutical Sciences), Ajmal Khan Tibbiya College & Hospital, Aligarh Muslim University, Aligarh, India



Cite This Article: Najmuddin Ahmad, Siddiqui, Ghufraan, Ahmad, Sumbul, Rehman and Asma, Abid (2016), "Antibacterial potential of Bisehri booti (*Areva lanata* Linn. Juss.)", *Pharmacophore*, Vol. 7 (2), 74-81.

