



PRELIMINARY FRACTIONATION AND QUANTIFICATION OF HYOSCYAMINE IN THE CRUDE ETHANOL EXTRACT OF *DATURA STRAMONIUM* LINN SEEDS

Idris A.T^{1*}, Sunday A.M², Ibrahim A.I³, James O.N⁴, Salisu A⁵

1. M.Sc., Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Science, Bayero University Kano, Nigeria.
2. Ph.D., Department of Human Anatomy, Faculty of Basic Medical Sciences, College of Medical Science, Ahmadu Bello University Zaria, Kaduna State, Nigeria.
3. Ph.D., Department of Human Anatomy, Faculty of Basic Medical Sciences, College of Medical Science, Federal University Dutse, Jigawa State, Nigeria.
4. Ph.D., Department of Veterinary Anatomy, Faculty of Veterinary Medicine, Ahmadu Bello University Zaria, Kaduna State, Nigeria.
5. Ph.D., Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Ahmadu Bello University Zaria, Kaduna State, Nigeria.

ARTICLE INFO

Received:
22 May 2020
Received in revised form:
17 Oct 2020
Accepted:
19 Oct 2020
Available online:
28 Oct 2020

Keywords: Chromatography, fraction, hyoscyamine, seeds, rf-value

ABSTRACT

Introduction: *Datura stramonium* Linn (*D. stramonium* L.) is an annual plant that belongs to the Solanaceae, a plant family that is poisonous and emits an unpleasant odor due to the presence of tropane alkaloids. **Aim:** The study was designed to determine the amount of hyoscyamine in the crude ethanol extract of *Datura stramonium* L. seeds. **Materials and Methods:** Fresh *D. stramonium* L. seeds were procured, identified, extracted, and fractioned in chloroform. The percentage yield was determined. The presence of alkaloids was qualitatively confirmed by phytochemical screening and thin-layer chromatography. The retention factor was determined. The high-performance liquid chromatography (HPLC) of the samples was carried out and the calibration curve was plotted to determine the amount of the hyoscyamine fraction. **Results:** An 11.04 µg/ml amount of hyoscyamine was obtained at the limit of detection (LOD) and quantification (LOQ) of 0.75 and 2.26 µg/ml respectively. **Conclusion:** In conclusion, the fraction of hyoscyamine in the *D. stramonium* L. seeds was little compared to the quantity of the crude seeds. **Key Messages:** The extraction and fractionation of hyoscyamine from the crude ethanol extract of *Datura stramonium* Linn seeds reveal the presence of several phytoconstituents including alkaloids. The hyoscyamine yield was little when compared to the crude sample of the seed used.

Copyright © 2013 - All Rights Reserved - Pharmacophore

To Cite This Article: Idris A.T, Sunday A.M, Ibrahim A.I, James O.N, Salisu A, (2020), "Preliminary Fractionation and Quantification of Hyoscyamine In the Crude Ethanol Extract of *Datura Stramonium* Linn Seeds", *Pharmacophore*, 11(5), 32-40.

Introduction

Datura stramonium (*D. stramonium*) is an annual plant that belongs to the Solanaceae, a plant family that emits an unpleasant smell due to the presence of tropane alkaloids [1]. All parts of the plant are toxic, especially the ripe seeds which contain the highest concentration of the active compound [2]. It contains several alkaloids, such as atropine, scopolamine [3, 4], and hyoscyamine. Other phytoconstituents include; megastigmane, sesquiterpenes, tannins, phlobatannins, cardiac glycosides, carbohydrates, and flavonoids, which are present primarily in the seeds and flowers [5]. The phytoconstituents vary with the species, developmental stage, environment, and part of the plant. The seeds are believed to be responsible for the anticholinergic activity of the plant [6]. *Datura* intoxication is characterized by anticholinergic symptoms [7] in both the central and peripheral nervous systems. The herb plays a significant role traditionally in the disease treatment due to the mind-altering property [8] it possesses. The fruit juice is used for the treatment of dandruff and falling hair when applied to the scalp [9]. It is applied to soothe painful wounds and sores [10]. Hyoscyamine is one of the predominant alkaloids of *D. stramonium* [10] that is used as anti-peristaltic/spasmodic [11] in the treatment of gastrointestinal-related disorders. The

Corresponding Author: Idris A.T, M.Sc., Department of Anatomy, Faculty of Basic Medical Sciences, College of Health Science, Bayero University Kano, Nigeria. Email: gwtel4u@gmail.com

seeds of this herb are used to calm hysteric, psychotic, and insomnia patients [12]. It also uses to relax smooth muscles in the bronchial tube during asthmatic bronchial spasm [13].

In Nigeria, *D. stramonium* L. seeds are consumed in a mixture of a locally made juice, *Zobo* by the youths especially at festivities to induce hallucinogenic sensation. This results in several cases of poisoning with fatal consequences. The fact that the plant constituents vary with location, soil region, pH, and time of harvest, calls for an evaluation of its poisonous phytoconstituents such as hyoscyamine. However, a few pieces of literature are available on the amount of hyoscyamine content in the crude extract of *D. stramonium* L. The present study was designed to quantify the amount of hyoscyamine in the crude ethanol extract of *D. stramonium* L. seeds. Plants are considered as a source of novel pharmaceutical products and inexpensive raw material for the synthesis of some known drugs, as they have thousands of active secondary metabolites [14-17].

Aims: to determine the amount of hyoscyamine in the crude ethanol extract of *Datura stramonium* Linn seeds

Settings and Design: study was quantitative carried out at the Human Anatomy Department, Faculty of Basic Medical Sciences, College of Medical Sciences, Ahmadu Bello University Zaria, Kaduna State, Nigeria.

Methods and Material

Ethical approval with registration ABUCAUC/2018/042 was obtained from the Ahmadu Bello University Committee on Animal Use and Care.

Plant material

Fresh *Datura stramonium* seeds were procured from Sharada residential area of Nassarawa Local Government, Kano State, Nigeria. The seeds were identified and the voucher number (VN108) was issued at the herbarium of Botany Department, Faculty of Life Sciences, Ahmadu Bello University, Zaria, Kaduna state, Nigeria. The seeds were separated from the pods, washed thoroughly with clean tap water, and air-dried under shade. Two thousand grams of the dried seeds were weighed using a digital weighing machine (LB-1000, China), grounded to a pulp using an electronic blender. The powdered sample was collected into a sterile cellophane bag and kept in a cool dry place for extraction.



Figure 1: *D. stramonium* L. (A): plant, (B): ripe fruits, (C): seeds, (D): powder

Extraction of *D. stramonium* L. Seeds

Extraction was carried out using cold maceration according to Djilani *et al.* [18]. The 200g of pulverized seeds was extracted in 1, 500 ml of 70% (v/v) ethanol at room temperature for 72 hrs (figure 2a). The extract was filtered and the solvent was evaporated in a water-bath at 40 °C (figure 2b). The residue, dissolved in H₂O and acidified with H₂SO₄ to pH 3-4, was extracted with petroleum ether and diethyl ether to remove lipophilic, acidic, and neutral material. After basifying the aqueous solution to pH 9-10 with NH₄OH (25%, m/m), it was extracted with chloroform. The extract washed with distilled water to neutral pH, dried with Na₂SO₄, and concentrated to dryness under reduced pressure to obtain crude alkaloids. Preliminary phytochemical analysis was carried out and the percentage yield of the crude extract was determined as;

$$\% \text{ Yield} = \frac{\text{Weight of the extract}}{\text{Initial weight of the sample}} \times 100$$




Figure 2: Ethanol extraction of crude *D. stramonium L.* seeds (A): cold maceration, (B): Ethanol extract on water-bath.

Fractionation of Hyoscyamine

The fractionation was carried out according to Salamah and Ningsih [19]. Five grams of the viscous extract was dissolved in 10 ml of water (figure 3a). Then, the solution was poured into a separating funnel, 10 ml of chloroform was added, and shaken to solve with two phases, namely water, and chloroform. These two phases were separated and collected (figure 3b). This was repeated until the chloroform phase had the same color as the chloroform solvent (figure 3d). The chloroform was then evaporated and recrystallized to obtain the hyoscyamine fraction (figure 3e). The alkaloid was analyzed with UV-Vis spectrophotometric method. The extraction and fractionation were carried out at the Department of Pharmacognosy, Faculty of Pharmaceutical Sciences, Ahmadu Bello University Zaria.

$$\% \text{ Yield} = \frac{\text{Weight of the fraction}}{\text{Initial weight of the sample}} \times 100$$



Figure 3: Fractionation of hyoscyamine fraction of *D. stramonium L.* seeds

A: Crude ethanol extract of the seeds dissolved in hot distilled water, **B:** Basification, partitioning, and dissolution of the crude extract of the seeds with ammonia, hydrochloric acid, and chloroform, **C:** Evaporation of the fraction in water-bath, **D:** Re-purification of the acid layer with ammonia, **E:** Evaporation of the chloroform layer, **F:** Hyoscyamine fraction of *D. stramonium L.* seeds in Eppendorf's tube.

The thin-layer chromatography (TLC)

With a pencil, a thin mark was made at the bottom of the pre-coated silica plate about 0.5 cm from the bottom to serve as the origin of the applied sample spots. Then, samples solutions (standard *l*-hyoscyamine powder and fractionate aliquot) were then applied separately and parallel to each other on the spots marked at the baseline at equal distances respectively. The mobile phase Butanol – Acetic acid – Water (BAW) in ratios of 4:1:1 was poured into the TLC chamber to a level of few centimeters above the chamber bottom. For the side of the plate with the sample line to face the mobile phase, the plate prepared with the sample spot was placed in the TLC chamber. The chamber was closed with a lid. This was allowed for spots to sufficiently develop to the solvent front of about 1 cm before the plate apex. After 1 hour, the plate was removed, sprayed Dragendorff's reagent, and heated in an oven for 5 minutes. The plate was visualized in a visible light chamber for the presence of the spots and calculation of R_f value [20].

$$R_f = \frac{\text{Distance travelled by the spot}}{\text{Distance travelled by the solvent front}}$$

Where: R_f = Retention factor.




Figure 4: Sprayed chromatographic plate showing orange-colored spots of hyoscyamine in the standard powder (St) and the *D. stramonium* L. sample (Ds)

The high-performance liquid chromatography quantification of the hyoscyamine fraction

A reversed-phase Techsphere 50DS C_{18} HPLC column (25 cm \times 4.6 mm i.d.) particle size 5 μ m, Supelco, Bellefonte, PA, USA) with oven temperature, 40°C in conjunction with a UV adsorption detector operating at 230 nm was employed. The mobile phase was a mixture of 20% acetonitrile, and 45 % methanol, 35 % water (H_2O), and 0.1 mol/L phosphoric acid which adjusted the pH to 7.0 and was used at a flow rate of 1 ml/min. A 1-mg aliquot of standard *l*-hyoscyamine (Sigma, St. Louis, MO, USA) was weighed using an electronic chemical balance (Shimadzu, USA) and was dissolved in 10 ml of standard methanol to make a stock solution of 100 mg/ μ L. A 6.4 ml of the stock solution was added to 10 ml of the methanol to make a 64 mg/ μ L solution. Two more solutions of concentration, 32 and 16 mg/ μ L respectively of the standard hyoscyamine were prepared by serial dilution to allow for a total of three graded solutions for the quantifying the hyoscyamine concentration amount in the sample fraction.




Figure 5: a&b, HPLC chromatogram of *l*-hyoscyamine standard powder (A) and hyoscyamine fraction of *D. stramonium* L. seeds (B).

The determination of the standard curve

The calibration curve for hyoscyamine was constructed by plotting the peak area of samples versus their concentrations (figure 6). The equation of a straight line was established, where *x* represents the concentration [$\mu\text{g/ml}$], while *y* = the area of the peak from the sample. The quantity of hyoscyamine compound in the *D. stramonium* L. ethanol extract was calculated at the moment that the area of the sample peak was reached using the equation. The area of the peak for hyoscyamine was taken from the chromatogram of *D. stramonium* L. fraction. The Department of Pharmaceutical Chemistry, Faculty of Pharmaceutical Sciences, Ahmadu Bello University Zaria provided the location for all the analyses that were carried out.




Figure 6: Standard curve for the quantification of hyoscyamine fraction using High-Performance Liquid Chromatography (HPLC) Analysis

Statistical analysis used: The percentage yields were expressed as the simple fraction of the samples and total crude samples. LINEST function test was used to develop the regression equation using the Microsoft Excel program.

Results and Discussion

Hyoscyamine and scopolamine are the major tropane alkaloids aside from several minor alkaloids identified in *Datura* species. Some of the minor alkaloids in *D. stramonium* include; tigloidin, aposcopolamine, apotropin, [21-24]. Table 1, shows phytoconstituents present in the crude ethanol extract of *d. stramonium* L. seeds. The present phytochemical analysis of the crude *D. stramonium* L. seeds showed that alkaloids, carbohydrates, saponins, cardiac glycosides, steroids, triterpenes, flavonoids, and tannins were all detected. This was in keeping with [2, 25, 26], which revealed the presence of similar phytoconstituents present in higher concentrations [27].

Table 1: Phytoconstituents Present in the Crude Ethanol Extract of *D. stramonium* L. Seeds

Test	Result	Confirmation
Alkaloids		
a. <i>Wagner's test:</i>	A reddish-brown precipitate was formed	+
b. <i>Dragendorff's test:</i>	A reddish-brown precipitate was formed	+
c. <i>Mayer's test:</i>	A whitish creamy precipitate was formed	+
		+
Carbohydrates		
a. <i>Molisch's test:</i>	A violet ring was formed	+
b. <i>Fehling's test:</i>	A brick-red precipitate was formed	+
		+
Saponins		
a. <i>Frothing test:</i>	A thin layer foam was formed	+
		+
Cardiac glycosides		
a. <i>Keller – Killiani test:</i>	A blue coloration was formed	+
		+
Tannins		
a. <i>Ferric Chloride test:</i>	A dark green coloration was formed	+
		+
Flavonoids		
a. <i>Alkaline reagent test</i>	A yellow fluorescence was formed	+
		+
Steroids & Triterpenes		
a. <i>Liebermann – Burchard</i>	A green coloration was formed	+

In *D. stramonium* L. seeds, hyoscyamine, and scopolamine account for 66% and 20% of the total tropane alkaloid content, respectively [28]. Our study observed that the percentage yield of crude ethanol extract of the seeds was 8.04%, while that of the hyoscyamine fraction from the crude extract was 3.60%. This suggested a relatively low yield of the fraction compared to the total crude extract used. The relatively low yield observed could be explained by the fact that hyoscyamine, which exists in a racemic mixture (1:1) of (-) -hyoscyamine and (+)-hyoscyamine, might have undergone some enantiomerisation thereby converting of (-)-hyoscyamine to (+)-hyoscyamine thus enabling the two enantiomers of hyoscyamine to coexist in a sample [29], thereby reducing the percentage abundance of the (-) -hyoscyamine. The enantiomerisation of hyoscyamine from (-) to (+)-hyoscyamine is possible under aqueous alkaline solution (pH > 9) combined with elevated temperatures (> 80 °C) conditions. Although cold maceration during crude ethanol extraction, the fractionation process involved heat application and basification of the sample at different stages. This has altered the optimal acid-base equilibrium in favor of the enantiomerisation. 63-70% of hyoscyamine and 42-80% of scopolamine were the decreases reported. Although part of the plant from which the extraction was made differed, the variation in the yield might also result due to intraspecies variation in genetic makeup, stage of development, duration of plant collection, and the difference in geographical location. Figure 4 shows two orange spots of the hyoscyamine in the extract and the standard powder was qualitatively identified using thin-layer chromatography. The chromatogram of the standard powder shows an enlarged orange spot of l-hyoscyamine sulphate (St) on the right-hand side, compared to a relatively small spot (Ds) of the extract on the right. Comparatively, the chromatogram of the sample resembled each other with regard to the position, size, and color. This proved that the hyoscyamine was the compound in the sample. The retention factor (Rf) value determined were 0.53 and 0.48 for the standard powder and the sample hyoscyamine fraction respectively, which suggested that both the sample and standard powder were alkaloids. This has confirmed the presence of alkaloid while the similar rf-value has confirmed alkaloid was hyoscyamine. Orange coloration was a typical alkaloid qualitative chromatographic test. In a similar study [30], orange coloration in the chromatogram plate of the tropane atropine and scopolamine alkaloids for the standard atropine sulfate and scopolamine hydrobromide with position and size that resembled the chromatogram of their standard powders observed.

Table 2. shows the graded doses of standard l-hyoscyamine powder, peak areas, and retention time produced by the HPLC for the quantification of hyoscyamine in the *D. stramonium* L. seed. The peak areas were dose-dependent. One calibration curve was obtained for hyoscyamine representing the peak area dependent on concentration based on the areas obtained. Figure 5a shows l-hyoscyamine (above) and figure 5b hyoscyamine fraction (below) chromatograms respectively. In both cases, the retention time was 1.660 and 1.683 min respectively. A good correlation of linearity has been achieved (n = 3; R² = 0.9995) and, in the range of 1–10 µg/ml for hyoscyamine. A limit of detection (LOD) and limit of quantification (LOQ) of the fraction was also found as 0.75 and 2.26 µg/ml respectively. Based on the HPLC chromatogram, the hyoscyamine came out from the sample at 1.683 minutes. In table 4, the quantity of the hyoscyamine in the seeds was 11.04 µg/ml obtained from the standard curve using 8250 µg/ml of the hyoscyamine fraction at peak area was 55.31 mUA*S. In a similar study [26], the alkaloids quantity in the ethanol extract of *D. stramonium* L. seeds was reported as 1.7 mg/mL, out of which 1.13 mg/mL for atropine and 0.57 mg/mL for scopolamine. High-Performance Liquid Chromatography (HPLC) was reported to

be one of the most commonly used techniques in alkaloids analysis [31]. We observed a relatively small quantity of the hyoscyamine, the lower limit of detection and quantification in the sample fraction. This was not a surprise as commercial production of alkaloids was generally considered to yield low, especially for hyoscyamine [32]. A study that quantified hyoscyamines in *Datura innoxia* and found a relatively even lower amount of the compound [33]. It was generally however reported that the main alkaloid in seeds of *D. stramonium* L. was the atropine although scopolamine, an enantiomer of hyoscyamine also present in significant amounts [34, 35], and that transformation of hyoscyamine-scopolamine occurs in *Datura* species. This probably could have attributed to the lower yield of the hyoscyamine in the fractioned sample.

Table 2: The standard quantity of hyoscyamine samples and areas

Quantity ($\mu\text{g/ml}$)	Peak Areas (mUA*S)	Time/min
16	09.4576	1.664
32	10.1167	1.667
64	11.1797	1.660

Table 3. shows the quantity of the hyoscyamine fraction of the ethanol extract of *D. stramonium* L. The absorption peak for hyoscyamine was at 1.683 minutes. Comparing the result with the absorption spectrums of standards hyoscyamine, this shows that the sample was correctly identified. Using the equations from the standard curves of standards solutions and that of the peak area from the extract, the alkaloids quantity in μL of the sample was calculated. Based on this, the concentration of the hyoscyamine alkaloid was calculated in the fraction of the *D. stramonium* L. seeds.

Table 3: Standard Quantity of Hyoscyamine Samples and Areas

Quantity ($\mu\text{g/ml}$)	Peak Areas (mUA*S)	Time/min
16	09.4576	1.664
32	10.1167	1.667
64	11.1797	1.660

Table 4: Hyoscyamine Quantity In the Fraction

Conc. ($\mu\text{g/ml}$)	Peak Areas (mUA*S)	Time/min	Qty ($\mu\text{g/ml}$)
8250	55.31	1.68	11.04

Conclusions

D. stramonium L. seeds contain several phytoconstituents including alkaloids. Although the fraction hyoscyamine was little when compared to the crude seed extract, this may suggest the high toxicity potential of the compound.

Acknowledgment

We sincerely appreciate Mallam Kabiru, Mallam Saidu Bala-riti, Abubakar Chawai Sulaiman of the Departments of Pharmacognosy, Pharmacology, and Pharmaceutical Chemistry for the assistance rendered in the course of this research.

References

1. Kumar A. A textbook of practical botany II. 7th ed. India: Rastogi Publications; 2009.
2. Shagal M, Modibbo U, Liman A. The pharmacological justification for the ethnomedical use of *Datura stramonium* stem-bark extract in the treatment of diseases caused by some pathogenic bacteria. *International Research Journal of Pharmacy and Pharmacology*. 2012;2:16-19.
3. Friedman M, Levin C. Composition of jimson weed (*Datura stramonium*) seeds. *Journal of Agricultural and Food Chemistry*. 1989;37(4):998-1005.
4. Brunton L, Hilal-Dandan R, Knollmann B. Goodman & Gilman's the pharmacological basis of therapeutics. New York: McGraw-Hill Med; 2006.
5. Godofredo S. Talong punay / Talampunay / *Datura metel* / Thorn Apple / Devil's trumpet: Medicinal Herbs / Philippine Alternative Medicine [Internet]. Stuartxchange.org. 2011 [cited 18 September 2020]. Available from: <http://www.stuartxchange.org/Talampunay.html>
6. Joshi P, Wicks A, Munshi S. Recurrent autumnal psychosis. *Postgraduate Medical Journal*. 2003;79(930):239-240.
7. Freye E. Toxicity of *Datura Stramonium*. *Pharmacology and Abuse of Cocaine, Amphetamines, Ecstasy, and Related Designer Drugs*. 2009;217-218.
8. Maheshwari N, Khan A, Chopade B. Rediscovering the medicinal properties of *Datura* sp.: A review. *Journal of Medicinal Plants Research*. 2013;7:2885-2889.

9. Hunter R, Anwar s, sembe S, Aliugo P. *Datura stramonium* Medicinal Uses, Side Effects and Benefits [Internet]. Gyanunlimited - A Hub of Alternative Medicine and Holistic Health. 2017 [cited 18 September 2020]. Available from: <http://www.gyanunlimited.com/health/datura-stramonium-medicinal-uses-side-effects-and-benefits/9184>
10. Soni P, Siddiqui A, Dwivedi J, Soni V. Pharmacological properties of *Datura stramonium* L. as a potential medicinal tree: An overview. *Asian Pacific Journal of Tropical Biomedicine*. 2012;2(12):1002-1008.
11. Bova J, Jurdi R, Bennett W. Antispasmodic drugs to reduce discomfort and colonic spasm during barium enemas: comparison of oral hyoscyamine, i.v. glucagon, and no drug. *American Journal of Roentgenology*. 1993;161(5):965-968.
12. Stafford G, Pedersen P, Jäger A, Van Staden J. Monoamine oxidase inhibition by southern African traditional medicinal plants. *South African Journal of Botany*. 2007;73(3):384-390.
13. Human bronchial smooth: Topics by Science.gov [Internet]. Science.gov. 2020 [cited 18 September 2020]. Available from: <https://www.science.gov/topicpages/h/human+bronchial+smooth>
14. Nagaram Choudhary and Mallya R. Phytochemical investigation and antibacterial activity of a medicinal plant. *International Journal of Pharmaceutical and Phytopharmacological Research*. 2019;9(4):53-8.
15. Yasin G, Anwer I, Majeed I, Sabir M, Mumtaz S, Mehmood A. "Pharmacodynamics of Secondary Metabolites Extracts of Some Plants from Cholistan Desert in Altering in Vitro Human Hematological Indices", *International Journal of Pharmaceutical and Phytopharmacological Research*. 2020;10(2):138-147.
16. Ahmad MS, Shawky A, Ghobashy MO, Felifel RH. Effect of Some medicinal plants on life cycle of Citrus Brown Mites (*Eutetranychusorientalis*). *International Journal of Pharmaceutical Research and Allied Sciences*. 2018 Oct 1;7(4):13-7.
17. Ayah B, Sara MN, Shima R. Decontamination of Eggshell Contaminated with *Salmonella Typhimurium* Using Natural Plant Extracts. I Some medicinal plants on life cycle of Citrus Brown Mites (*Eutetranychusorientalis*). *International Journal of Pharmaceutical Research and Allied Sciences*. 2018 Jan 1;7(3):10-9.
18. Djilani A, Legseir B, Soulimani R, Dicko A, Younos C. New extraction technique for alkaloids. *Journal of the Brazilian Chemical Society*. 2006;17(3):518-520.
19. Salamah N, Ningsih D. Total alkaloid content in various fractions of *Tabernaemontana sphaerocarpa* Bl. (Jembirit) leaves. *IOP Conference Series: Materials Science and Engineering*. 2017;259:012017.
20. Yubin J, Miao Y, Bing W, Yao Z. The extraction, separation, and purification of alkaloids in natural medicine. *Journal of Chemical and Pharmaceutical Research*. 2014;4:338-345.
21. Shaikh A, Sablay Z. Hallucinogenic plant poisoning in children. *Saudi Medical Journal*. 2005;26:118-121.
22. Romeike A. Beit räge zur chemischen Physiologie der mydriatisch wirkenden Solanaceen-Alkaloide. *Pharmazie*. 1953;8:668-674 and 729-747.
23. Phillipson J, Handa S. N-oxides of hyoscyamine and hyoscyne in the Solanaceae. *Phytochemistry*. 1975;14(4):999-1003.
24. Bucher E, Meszaros L. Stechapfel Samen (*Datura spec.*) und deren Alkaloide in Futtermitteln. *Kraftfutter*. 1989;3(89):76-82.
25. Ivancheva S, Nikolova M, Tsvetkova R. Pharmacological activities and biologically active compounds of Bulgarian medicinal plants. *Phytochemistry: Advances in Research*. Kerala: Signpost; 2006.
26. Steenkamp P, Harding N, van Heerden F, van Wyk B. Fatal *Datura* poisoning: identification of atropine and scopolamine by high-performance liquid chromatography/photodiode array/mass spectrometry. *Forensic Science International*. 2004;145(1):31-39.
27. Oseni O, Olarinoye C, Amoo I. Studies on chemical compositions and functional properties of thorn apple (*Datura stramonium* L) Solanaceae. *African Journal of Food Science*. 2011;5(2):40-44.
28. El Bazaoui A, Bellimam M, Soulaymani A. Nine new tropane alkaloids from *Datura stramonium* L. identified by GC/MS. *Fitoterapia*. 2011;82(2):193-197.
29. Eich E. Solanaceae and Convolvulaceae: secondary metabolites. Biosynthesis, chemotaxonomy, biological and economic significance. A handbook. Germany: Springer; 2008.
30. Matsuura H, Malik S, de Costa F, Yousefzadi M, Mirjalili M, Arroo R et al. Specialized Plant Metabolism Characteristics and Impact on Target Molecule Biotechnological Production. *Molecular Biotechnology*. 2017;60(2):169-183.
31. Tayoub G, Sulaiman H, Alorfi M. Quantitative identification of Atropine and Scopolamine in wild *Datura* (*Datura metel*) using HPLC. *Journal of Natural Products*. 2016;9:21-28.
32. List G, Spencer G, Hunt W. Toxic weed seed contaminants in soybean processing. *Journal of the American Oil Chemists' Society*. 1979;56(8):706-710.
33. Klein-Schwartz W, Oderda G. Jimsonweed Intoxication in Adolescents and Young Adults. *Archives of Pediatrics & Adolescent Medicine*. 1984;138(8):737.
34. Dugan G, Gumbmann M, Friedman M. Toxicological evaluation of jimson weed (*Datura stramonium*) seed. *Food and Chemical Toxicology*. 1989;27(8):501-510.
35. Miraldi E, Masti A, Ferri S, Barni Comparini I. Distribution of hyoscyamine and scopolamine in *Datura stramonium*. *Fitoterapia*. 2001;72(6):644-648.

Appendix-1



AHMADU BELLO UNIVERSITY, ZARIA
DIRECTORATE OF ACADEMIC PLANNING & MONITORING

Vice Chancellor Prof. Ibrahim Garba, B.Sc. (Hons) Geology, M.Sc. (Mineral Exploration) A.B.U., Ph.D Geology (London), D.L.C., FNMGS
Director: Prof. M.F. Ishiyaku, B.Sc. (Hons) Botany (ABU), M.Sc. Plant Breeding (UnJos), Ph.D Agriculture (University of Reading, U.K.), MASN, MBSN

Tel. Zaria + 234-069-551912 Fax. 234 (069) 550022 Telegram: UNIBELLO Zaria e-mail: dapm@abu.edu.ng

Appl No.: ABUCAUC/2018/Human Anatomy/042
Approval No: ABUCAUC/2018/042

3rd October, 2018

Dr. S. A. Musa
Department of Human Anatomy,
Faculty of Basic Medical Sciences,
College of Health Sciences,
Ahmadu Bello University,
Zaria.

Sir,

APPROVAL OF RESEARCH STUDY 'EVALUATION OF INGESTION OF HYOSCYAMINE FRACTIONATE OF DATURA STRAMONIUM SEEDS ON CEREBRUM DEVELOPMENT IN WISTAR RATS (*RATTUS NORVEGICUS*)'

This is to convey the approval of the ABUCAUC to you for the aforesated study domiciled in the Department of Human Anatomy. The approval is predicated on the assumption that you shall maintain and care for the Experimental Animals as approved after the visitation of the Committee.

Monitoring of the Research by spot checks, invitations or any other means the Committee deems fit shall be undertaken at the convenience of the Committee.

This approval can and shall be revoked should a significant breach in the terms and condition of the approval occur. It is hence your responsibility to ensure that the agreed terms are maintained to the end of the Study.

The said approval shall be posted on the ABUCAUC Page on the University's website. Note upon completion of the research, ethical clearance certificate will be issued.

U.D. Abdullahi

For: Chairman, ABUCAUC.

Cc. Director, DAPM
Director, IC & ICT
Provost, College of Health Sciences
Dean, Faculty of Basic Medical Sciences,
HOD, Human Anatomy,
Prof. Aliyu Mohammed