



## ANTIBACTERIAL ACTIVITY AND GC-MASS ANALYSIS OF ORGANIC EXTRACT FROM PERSIAN GULF *HALICLONA SP*

Mohammad Reza Shushizadeh<sup>1\*</sup>, Sahar Behroozi<sup>2</sup>, Abdol-Azim Behfar<sup>1,4</sup>, Melika Nazemi<sup>3</sup>

1. Marine Pharmaceutical Science Research Center, Ahvaz Jundishapur University of Medical Sciences, IR Iran
2. Islamic Azad University of Ahvaz, Science & Research branch, IR Iran
3. Persian Gulf and Oman Sea Ecological Center, Iranian Fisheries Science Research Institute, Agricultural Research, Education and Extension Organization (AREEO), Bandar Abbas, Iran
4. Food and Drug Laboratory Research Center, Food and Drug Organization, MOH&ME, Tehran, Iran

### ARTICLE INFO

#### Received:

29<sup>th</sup> Sep 2017

#### Received in revised form:

04<sup>th</sup> Mar 2018

#### Accepted:

16<sup>th</sup> Mar 2018

#### Available online:

28<sup>th</sup> Apr 2018

**Keywords:** Marine Sponges, Extract, Diethyl Ether, Cholestanol, Antibacterial Activity

### ABSTRACT

Marine organisms given the opportunity to develop more than terrestrial organisms, potentially producing new molecules in the properties of interest therefore marine organisms give us more molecules in which the result would be more information for research. As marine organisms, sponges are a variety of chemicals, including terpenes, polysaccharides and sterols have medicinal properties, antioxidant, antiviral, antibacterial, antifungal, anticancer and AIDS inhibition. Persian Gulf has many species of sponges and *Haliclona sp.* was collected from Kharg Island, then rinsed and finally extracted with ether as solvent. Five combinations of cholestanol, 2-(Acetoxymethyl)-3-(methoxycarbonyl)bi-phenylene carboxylate, silicic acid diethyl bis(trimethylsilyl) ester, 2,4-dimethyl-Benzo[H]Quinoline and 1-(3,5-dimethylpyrazolyl),2-(2-benzothiazolylthio) ethanone were analyzed using gas chromatography and mass spectrometry methods. Finally antimicrobial capacity was determined.

Copyright © 2013 - All Rights Reserved - Pharmacophore

**To Cite This Article:** Mohammad Reza Shushizadeh, Sahar Behroozi, Abdol-Azim Behfar, Melika Nazemi, (2018), "Antibacterial activity and GC-Mass analysis of organic extract from Persian Gulf *Haliclona SP.*", **Pharmacophore**, 9(2), 19-24.

### Introduction

Marine animals in general and marine invertebrates in particular are promising organisms for preparation of biological active compounds.[1] Marine sponges (Porifera) are considered to be true "chemical factories" producing natural compounds which have been identified and elucidated by physical and spectroscopic properties, but their bioactivities are still unknown.[2] *Haliclona* sponges have produced many natural materials which can be used for preparation of steroids,[3] polyketides,[4] macrolides,[5] alkaloids,[6] terpenes,[7] with antioxidant, antimicrobial and anticancer activities.

In this research work we report the GC-Mass analyzing results of several compounds corresponded to organic extracts from Persian Gulf marine sponge, *Haliclona* species for possible antibacterial activity of them.

### Materials and Methods

#### Collection of sample

Marine sponge was collected from Kharg Island of Persian Gulf at depths of 10-15 m in September 2013. Sample was transported and stored frozen prior to exhaustive extraction in organic solvent. Taxonomic was done and certified by Iranian Fisheries Science Research Institute Bandar Abbas. The sample was identified as *Haliclona sp.* as shown in figure 1.



**Figure 1.** Persian Gulf marine sponge, *Haliclona species*

**Preparation of the extract**

50 g of sponge sample was extracted with diethyl ether (100 ml) at 25 °C overnight on a shaker. The extract was filtered by a Buchner funnel using suction and kept in the refrigerator -20 °C for antibacterial activity and further GC-Mass analysis.

**Microorganisms and culture media**

All microorganisms i.e., *Escherichia coli*, *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Bacillus subtilis* were obtained from the Microbiology Laboratory of Ahwaz Golestan Hospital.

**Antimicrobial assay**

Well agar diffusion method was used for microbial test of extract. Nutrient agar was utilized as culturing media for microorganisms. For antibacterial activity, 20ml of agar was poured in each petri plate, allowed to set at 37 °C for 24 hr. Cultures were swabbed in nutrient agar plates using sterile cotton.

The organic extracts were prepared by transferring 10, 20, 30, 40 and 50 mg of each extract into 1ml of DMSO (Merck, Germany), swab them and incubated at desired condition. The inhibitory zone (mm) was applied for antibacterial activity. Antibiotics such as vancomycin, tetracycline, penicillin, gentamycin, ceftriaxone, amoxicillin were used as positive controls. DMSO and Et<sub>2</sub>O were also tested as negative control to ensure that they do not interfere with the tests.

*Haliclona sp.* extract demonstrate considerable antibacterial activity, was selected for determination of MIC.[8]

**Minimum inhibitory concentration (MIC)**

Small volume microtitre plates and a stock of 50 mg/ml of extract into DMSO were used in this method.[8] To determine MIC of extract, 100µL of Mueller Hinton broth was transferred into microtitre plate. From this reason, 100 µL was placed into wells of column 1. The contents of well were mixed thoroughly, 100 µL from column 1 and added to column 2, two-fold serial dilution. Then, each well is inoculated with 50 µL of strain of microorganism pre-adjusted at 0.5 McFarland Standard.

**Minimum bactericidal concentration (MBC)**

For determination of MBC, samples with negative microbial growth were subjected for determination of living cells after 24 h of incubation.

**Gas chromatography-mass spectrum (GC-MS)**

GC-MS Agilent GC 7890, Mass 5975 fitted with a fused HP-5ms (5% phenyl methylpolysiloxane) capillary column (0.25 mm × 30 m, 0.25 µm film thickness) with helium gas at 1 ml/minute are used. The sample was injected (sampling time, 5 minutes). The respective temperatures and ionization chamber were 290°C and 280°C. Temperature programs for the column oven were as follows: program, 60°C for 1 minute, elevated to 130°C at 20°C/minute, then to 210°C at 10°C/minute, then to 260°C at 10°C/minute, then to 300°C at 10°C/minute; it was finally maintained at 300°C.

**Identification of compounds**

Interpretation of mass spectra was done according to National Institute Standard and Technology (NIST 08s), WILEY 8 and FAME.

**Results**

The *Haliclona sp.* extract was tested for antimicrobial activity against 4 bacterial species namely *E.coli*, *P. aeruginosa*, *S. aureus* and *B. subtilis* by the well agar diffusion method. All extracts in table1 with 50 mg/ml concentration exhibited activity. All test microorganisms was present mainly in test extract. Moreover, the data in this table indicate that this extract was more effective as compared to other marine sponges.[8]

**Table 1.** Antimicrobial activity of Persian Gulf sponge

Microorganisms	Diethyl ether	Tetracycline	Penicillin	Amoxicillin	Ceftriaxone	Gentamicin	Vancomycin
<i>E. coli</i>	21±2.31	13±1.2	17±1.01	17.33±1.5	25.32±1.7	28.33±1.33	19.67±1.35
<i>P. Aeruginosa</i>	6.33±2.5	18.6±1.25	8.41±1.04	5.31±1.21	13.67±1.33	18.31±1.21	18.7±1.25
<i>S. aureus</i>	28.67±2.5	18.6±1.33	7.37±1.43	6.45±1	8.33±1.22	18.42±1.04	10.33±1.23
<i>B. subtilis</i>	26.67±1.35	25.33±1.11	19.12±1.21	21.39±1.02	35.51±1.13	23.12±1.35	16.46±1.35

**Table 2.** MIC and MBC of Persian Gulf sponge

Extract	Microorganisms							
	E.Coli		P.Aeruginosa		S.Aureus		B.Subtilis	
	MIC	MBC	MIC	MBC	MIC	MBC	MIC	MBC
Diethyl ether	12.5	50	12.5	25	12.5	50	3.12	50

The extract was effective against pathogenic bacteria. The MIC of extract at 3.12 mg/ml show inhibition the growth of *B. subtilis*.

Finally desired extract was tested to TLC for the preliminary compound separation and then to GC-MS analysis to characterize the compound. GC-MS of the *Haliclona* sp. (Figure 2 and Table 3) revealed compounds namely Cholestanol, 2-(Acetoxymethyl)-3-(methoxycarbonyl)bi-phenylene carboxylate, silicic acid diethyl bis(trimethylsilyl)ester, 2,4-dimethyl-Benzo[H]Quinoline, and 1-(3,5-dimethylpyrazolyl) 2-(2-benzothiazolylthio) ethanone. Cholestanol was identified as sterol compound with high abundant (82.75%). MS spectra of five mentioned compounds are given in Figures 3,4.




Figure 2. GC Spectrum of *Haliclona* sp. extract

Table 3. Bioactive of components identified in the extract

Entry	RT	Name of compound	Peak area %	Nature of compound
1	25.168	Cholestanol	82.75	Sterol
2	25.340	2-(Acetoxymethyl)-3-(methoxycarbonyl) biphenylene	2.12	Aromatic compounds
3	25.603	Silicic acid, diethyl bis(trimethylsilyl) ester	1.93	Silicon ester
4	26.284	2,4-dimethyl-Benzo[H]quinoline	7.65	Aromatic compounds
5	27.217	1-(3,5-dimethylpyrazolyl), 2-(2-benzothiazolylthio) Ethanone	5.55	Aromatic compounds




Figure 3. Mass spectrum of *Haliclona sp.* extract with RT=25.168 (right) and cholestanol reference (left)




Figure 4. Mass spectrum of extract with RT=26.284 (right) and 2,4-dimethyl-Benzo[H]quinoline reference (left)

## Discussion

Antibacterial activity has previously been described in marine sponge species.[9] A variety of antimicrobial compounds, including terpenes[10-12], sterols[13] and alkaloids[14] have been isolated from them. The antibacterial activity of extract in Table 1 against gram-negative bacteria proved more effective than gram-positive. Also this extract was effective against *S. aureus* compared with penicillin, amoxicillin, tetracycline, vancomycin, gentamycin and ceftriaxone.

Marine supplies are a potent source for unexplored chemical moieties that may have vital biological and economic properties. Marine diaspora has proven its importance as several of compounds of economic significance are isolated and mass produced across the globe. Very little of these living organisms was studied and therefore provides ample opportunities to hunt for vital substances.

In this study, GC-MS as a good method can be analyze the unknown organic materials from Persian Gulf sponge *Haliclona sp.*. We obtained mixtures of sterols with other compounds which investigated by chromatography on silica gel columns. GC investigation of isolated mixture showed at least 5 different compounds. So this mixture was subjected to GC-MS. For example

the reference mass spectrum and fragmentation pattern corresponded to cholestanol were displayed in figure 5 and figure 6.[15]




Figure 5. Mass spectrum the reference of cholestanol




Figure6. Fragmentation pattern of cholestanol

### Conclusion

Persian Gulf sponge *Haliclona sp.* produces numerous unique biomaterial of potential medicinal value. This species has several components. From it, five different compounds were identified by GC-MS. Most of these compounds are widely can be used in pharmaceutical, medicinal, cosmetic and other fields. The results clearly showed that high antimicrobial activity of extract against various gram-positive and negative bacteria. The MIC of extract inhibited the growth of *B. subtilis* at 3.12 mg/ml.

## References

1. Richelle-Maurer E., Gomez R., Braekman J.C, Van de Vyver G., Van Soest R.W.M., and C. Devijver, Primary cultures from the marine sponge *Xestospongia muta* (Petrosiidae, Haplosclerida). *J Biotech*, 100, 2003, 169-176.
2. Bacus, G.J; Green, G. Toxicity in sponges and holothurians: a geographic pattern. *Science* 1974, 13, 951–953.
3. Findlay, J. A.; Patil, A. D. Novel sterols from the finger sponge *Haliclona oculata*. *Can. J. Chem.* 1985, 63, 2406–2410.
4. Williams DH, Faulkner DJ: Three new acetylenes from the palauan sponge *Halaciona* sp. *Journal of Natural Product*. 1996, 59: 1099-1101.
5. Erickson, K. L.; Beutler, J. A.; Cardellina, J. H., II; Boyd, M. R. Salicylhalamides A and B, Novel Cytotoxic Macrolides from the Marine Sponge *Haliclona* sp., *J. Org. Chem.* 1997, 62, 8188–8192.
6. Koren-Goldshlager, G.; Kashman, Y.; Schleyer, M. Haliclorensin, a Novel Diamino Alkaloid from the Marine Sponge *Haliclona tulearensis*. *J. Nat. Prod.* 1998, 61, 282–284.
7. Crews, P.; Harrison, B. New triterpene-ketides (merotriterpenes), haliclotriol A and B, from an Indo-Pacific *Haliclona* sponge. *Tetrahedron* 2000, 56, 9039–9046.
8. Safaeian S., Hosseini H., Abbas Pour Asadolah A., Farmohamadi S. Antimicrobial activity of marine sponge extracts of offshore zone from Nay Band Bay, Iran. *Journal of Medical Mycology* (2009) 19, 11-16.
9. Rifai S., Fassouane A., El-Abbouyi A., Wardani A., Kijjoa A. and Van Soest R. Screening of antimicrobial activity of marine sponge extracts. *Journal of Medical Mycology* (2005) 15, 33–38.
10. Musman M, Ohtani I, Nagaoka D, Tanaka J, Higa T. Hipposulfates A and B, new sesterterpene sulfates from an Okinawan sponge, *Hippospongia* cf. *metachromia*. *J Nat Prod* 2001;64:350–2.
11. Motti, C.A., Bourguet-Kondracki, M.L., Longeon, A., Doyle, J.R., Llewellyn, L.E., Tapiolas, D.M. and Yin, P., 2007. Comparison of the biological properties of several marine sponge-derived sesquiterpenoid quinones. *Molecules*, 12, 1376-1388.
12. Blunt, J.W., Copp, B.R., Hu, W.-P., Munro, M.H., Northcote, P.T. and Prinsep, M.R., 2007. Marine natural products. *Natural Product Reports*, 24, 31-86.
13. Wei-Hua, J., Jing, L., Qian, L., TingTing, X., Guo-Hua, S., Hao-Bing, Y., Fan, Y., Bing-Nan, H., Min, L. and Hou-Wen, L., 2014. Dysidinoid A, an unusual meroterpenoid with anti-mrsa activity from the South China Sea sponge *Dysidea* sp. *Molecules*, 19, 18025-18032.
14. Laport M.S., Santos O.C.S. and Muricy G. Marine sponges: potential sources of new antimicrobial drugs. *Current Pharmaceutical Biotechnology*, 2009, 10, 86-105.
15. S. Ramos S., Ramalho L., Costa M., Teixeira A., Almeida P., Costa M. 2007 Mass spectrometric analysis of epimeric 3 $\alpha$ /3 $\beta$  functionalized steroids: ARKIVOC (xiv): 163-171.