

## KERMES DYE EXTRACT FROM *COCCUS ILICIS* INSECT AS AN ALTERNATIVE COUNTER STAIN INSTEAD OF EOSIN IN VARIOUS TISSUE CONSTITUENTS: AN EXPERIMENTAL STUDY

Nosiba Elsafi Mustafa<sup>1</sup>, Osman Mohammed Elmahi<sup>2</sup>, Hisham Ali Waggiallah<sup>3\*</sup>  
Lienda Bashier Eltayeb<sup>3</sup>

1. *Department of Histopathology and Cytology, Faculty of Medical Laboratory Sciences, University of Khartoum. Khartoum, Sudan.*
2. *Department of Histopathology and Cytology, Faculty of Medical Laboratory Sciences, University of Karary, Omdurman, Sudan*
3. *Department of Medical Laboratory Sciences, College of Applied Medical Sciences, Prince Sattam Bin Abdulaziz University, Alkharj, Saudia Arabia.*

### ARTICLE INFO

#### Received:

30 Apr 2020

#### Received in revised form:

27 Aug 2020

#### Accepted:

02 Sep 2020

#### Available online:

28 Oct 2020

**Keywords:** Counterstain, kermes dye, Crimson, eosin, hematoxylin, K.vermilio, K.ilices

### ABSTRACT

Kermes is one of the known oldest dyes. It is obtained from the bodies of an insect, Kermes ilices (formerly known as Coccus ilicis), and Kermes vermilio. It is chemically very similar to carmine and, as the older name indicates, the insects are related to those from which carmine is obtained. This study was conducted to evaluate the efficacy of Kermes dye as a counterstain after hematoxylin, and to compare its staining ability with that of routinely used Eosin dye. 0.5g and 1gm of kermes dye were respectively dissolved in 100 ml of D.W. The solution was used to stain different tissue structures and the result was compared with 0.5% and 1% eosin. It was revealed that kermes dye could be used as a counterstain after hematoxylin, its staining ability was also good and comparable to that of eosin dye with a special affinity for collagen and muscle fibers. Kermes dye could be used as a histological stain, which stains similar to eosin dye and its specific affinity for collagen and muscle fibers authenticates its role in the treatment of collagen and muscle disorder.

Copyright © 2013 - All Rights Reserved - Pharmacophore

**To Cite This Article:** Nosiba Elsafi Mustafa, Osman Mohammed Elmahi, Hisham Ali Waggiallah, Lienda Bashier Eltayeb, (2020), "Kermes dye extract from *Coccus ilicis* insect as an alternative counter stain instead of Eosin in various tissue constituents: An experimental study", *Pharmacophore*, 10(5), 14-19.

### Introduction

Although there is huge progress in diagnostic facilities in histopathology, simple stains and H&E still remain of an important value [1, 2]. This study was done to find out the staining ability of crimson as a natural dye. Because the natural dyes are cheap, easy to prepare and use, available, eco-friendly it could substitute many different dyes in the histopathology lab [3, 4]. Crimson (C) is a strong reddish - purplish color, originally used to describe the color of the kermes dye produced from a scale insect, Kermes vermilio that feed on the sap of Kermes oak [5] "Family Kermesidae (Hemiptera, Coccoidea) exclusively develops and feeds on Fagaceae trees [6]. This scale insect family contains 100 species distributed among 10 genera and they are currently known in the Palaearctic, Oriental, and Nearctic regions of the world. Kermes Boitard is the major genus in the Mediterranean and European regions. In these regions, 20 species of Kermes have been recorded, all off deciduous and evergreen oak trees (Quercus) [6]. Some scale insects have been known as sources of red dye used in the textile, art, and wine industries in the Mediterranean, Middle East and Central Asia regions for many centuries [7]. Researchers have suggested that the red dye used for both ritual and secular purposes during ancient times in Israel apparently had been imported from neighboring countries where such dye producing scale insects as Kermes vermilio Planchon and Porphyrophora hameli Brandt are known [8]. However, red dye has been extracted from both the eggs and adult females of Kermes echinatus and chemically analyses have suggested that Kermes echinatus might be the "Tolaat Shani" (scarlet worm in Hebrew) [9]

"Kermesic acid, as Dimroth termed the pigment, was first prepared in the pure state by Heisse. According to Dimroth, who followed Heisse's procedure, the crude material is first released from a wax, cerotic ceryl ester/no by extracting with ether;

**Corresponding Author:** Dr. Hisham Ali Waggiallah, Assistant Professor of Hematology and Blood transfusion, College of Applied Medical Sciences, Prince Sattam University, Al- Kharj-Riyadh, Saudi Arabia. E-mail: [hishamwagg30@hotmail.com](mailto:hishamwagg30@hotmail.com)

then the pigment, which is present as a salt, is released by ethereal hydrochloric acid and taken up in ether. It is purified through the sodium salt and again precipitated with acid. 5 kg of crude material yield approximately 50-55 grams of pigment. Kermesic acid, C forms brick-red needles and is dissolved in water to yellow-red, and in sulfuric acid to violet-red solutions. It dyes wool orange-red from an acid bath and gives scarlet-red dyeings, which are less blue than those of carmine on a tin mordant. The constitution of kermesic acid 211 is deduced from the following observations. It contains four hydroxyl groups, which may be acetylated, and oxidation yields nitrococcus acid, also obtained from cochineal. Methylation with dimethyl sulfate gives a trimethyl derivative, by oxidation of which with potassium permanganate the methyl esters of two acids are obtained". Crude kermes contains, in addition to kermesic acid, a second pigment, flavokermesic acid. Carmine could be used as a source of Kermesic acid and flavokermesic acid [10].

## Materials and Methods

The present experimental study was carried out in the Department of Histopathology and Cytology, Faculty of Medical Laboratory Sciences, University of Khartoum.

### Preparation of Kermes dye:

The Kermes dye powder or crystals are available commercially; 0.5g and 1 g of this powdered material were weighted using electronic/digital weighing machine, then dissolved in 100 ml of D.W in a beaker. Two concentrations of this dye were selected to standardize this dye with different times namely; 2min, 4min, and 8 min.

### Rabbit dissection and tissue preparation:

Three successive days well-fed rabbit put in a closed cabinet with a piece of cotton filled with chloroform, rabbit died after about 20 minutes, this was assured after the pulse the heart has been checked for different time intervals. After that, the rabbit was fixed in a dissection board and most of the tissues were taken and washed with normal saline, this was followed by fixing them in 10% formalin (each organ was put in a closed urine container containing about 50 ml of 10% formalin) at the room temperature for 48 hours.

The following are the size of the tissue taken from the rabbit (sometimes just part of the tissue was taken in case of the large organs):

- Liver: ½ one lobe 5cm × 3cm.
- Tongue: 4cm×1.2cm×1cm.
- Colon: We took just 3Cm × 0.5Cm long.

Small pieces from all the organs mentioned above (about 3-5mm) were taken in a separate cassette, processed with an automatic tissue processor following the standard routine tissue processing programs protocol:

### Staining various tissue structures:

Two sets of 3 μ thick sections were obtained. Each set comprised of 180 slides (60 liver, 60 tongues, and 60 colons). Sections were first dewaxed in xylene for 6 min and hydrated through descending grades of alcohol, then were stained with Mayer's hematoxylin for 10 minutes, followed by washing in running tap water for 10 minutes, then one set (½ of the sections / 30 of each organ) was counterstained with 0.5% eosin, and the remnants (30 of each organ) were stained with 0.5% Crimson for 3min for the two counterstains. The slides were rinsed then dehydrated by air and mounted.

Sections were then dehydrated by air, cleared in xylene, and finally mounted with the synthetic (DPX) mounting media. Sections were observed using binocular Olympus microscope model no. CH20i at ×10 and ×40 magnifications.

## Results

The slides were evaluated by experts subjectively for the quality of cytoplasm and connective tissue staining, for nuclei staining, and finally for nuclear-cytoplasmic contrast.


Table 1 shows the frequencies and compare between the two counterstain staining qualities.

**Table 1: Comparison of Crimson and Eosin dyes used in different types of tissues**


Parameters	Type of stain	Liver	Tongue	Colon	P-value
Score of nucleus stain	Eosin	Good 2(2.2%) Excellent 28(31.1%)	Good 2(2.2%) Excellent 28(31.1%)	Good 2 (2.2%) Excellent 28(31.1%)	0.058
	Kermes (Crimson)	Good 10(11.1%) Excellent 20(22.2%)	Good 0(0.0%) Excellent 30(33.3%)	Good 4(4.4%) Excellent 26(28.9%)	
	Eosin	Good 0(0.0%) Excellent 30(33.3%)	Good 15(16.7%) Excellent 15(16.7%)	Good 15(16.7%) Excellent 15(16.7%)	

Score of cytoplasm stain	Kermes (Crimson)	Good 4(4.4%) Excellent 26(28.9%)	Good 3(3.3%) Excellent 27(30.0%)	Good 9(10.0%) Excellent 21(23.3%)	0.043*
Nucleus cytoplasm contrast	Eosin	Excellent 30 (33.3%)	Poor 1 (1.1%) Good 4 (4.4%) Excellent 25 (27.8%)	Good 1(1.1%) Excellent 29 (32.2%)	0.013*
	Kermes (Crimson)	Excellent 30 (33.3%)	Excellent 30(33.3%)	Excellent 30(33.3%)	


\*P ≥ 0.05 significant




**Figure 1:** Colon stained with Mayer's hematoxylin and 0.5% Crimson for 3minutes in the right, and with H&E in the left. RBCs are sharply stained.




**Figures 2:** Tongue Stained with HK (Hematoxylin in the left and Kermes dye/Crimson in the right) for 3miutes




**Figure 3:** stained with HE for 3min ( $\times 40$ ). Notice increasing the light source with the photo on the upper right (Kermes dye), which shows the epithelium of the tongue



**Figure 4:** shows liver stained with Mayer's counter stained with HE



**Figure 5:** The liver stained with Mayer's counter stained with HK (Crimson).



**Figure 6:** Muscle stained with 0.05% HK for 2 min. Notice the obvious muscle striations

## Discussion

Kermes dye is an old dye that is produced from the dried body of the female gender of some species of the family Kermesidae, and it has been used for dyeing textiles and for cosmetic issues [11, 12].

To our knowledge, this trial may be the first attempt to use kermes dye in histology. Although we found no paper discussing this issue, there are a lot of references that mentioned the chemical structure of the kermes dye. Generally, the reaction of the Crimson seems to be similar to that of acidic dyes (where acidic dyes have a special affinity to basic tissue components) (Figure 1)

As some synthetic dyes have an inherent tendency to be toxic and carcinogenic, it is recommended to be substituted by other more eco-friendly dyes. Crimson is safe to the environment in terms of toxicity and can be accounted for simplicity of preparation as a green chemical and can be used in research as a cost-effective reagent. Crimson and green chemical are prepared without any additives; therefore, the quality of staining may be enhanced with some additives such as glacial acetic acid.

In this research, several trials were accomplished with 72 slides (36 with Harris hematoxylin and the rest with Mayer's hematoxylin), to standardize the concentration and determine the suitable time of staining, which is between 1-4 min, and the perfect time is 2 or 3min. The stain is very sharp and crisp especially with Harris hematoxylin when the light source was increased (Figure 2). It is better to use the stain within two weeks of preparation because its color starts to fade gradually, a fresh solution does not need filtration but after one week it is better to be filtered. In this study, the best concentration of Crimson that could be used was 0.5%, for about 1-4 minutes. There may be some variations for different tissue types because of the sample's nature, poor staining was developed due to the increase of both, concentration and time of staining. No slide was poorly stained with 0.5% crimson, which means in tissues containing a high amount of collagen fibers or muscles (such as tongue) (Figures 2,3), satisfactory cytoplasmic staining may be obtained in just two minutes.

The Kermes stain (Crimson) can stain a wide range of tissue structures with varying shades of colors, these could be observed on RBCs, keratin, muscle, and fibers all these structures could be easily differentiated (Figure 5).

Statistical analysis was performed with SPSS version 22. As shown in Table 1 Pearson correlation (2-tailed) led to the following results: comparison between the type of cytoplasmic stains (Eosin and Crimson) to identify the quality of stain (P-value) = 0.043, which was considered significant at the 0.05 level. This indicated the effect of cytoplasmic stain type on the quality of cytoplasmic staining. Correlation between the type of cytoplasmic stain and the score of nuclear stain was not significant (correlation coefficient = 0.141, P-value = 0.058), which indicates no effect of the type of cytoplasmic stain on the quality of nuclear stain. On the other hand, the type of cytoplasmic stain would surely affect the score of nuclear-cytoplasmic contrast, because the correlation coefficient between the two variables was -0.186 and the p-value were 0.013, which is highly significant. All data are summarized in Table 1.

The mean variation of nuclear stain between Crimson and eosin was -0.089 and 2-tailed significance (P-value = 0.058), which indicates no significant difference between the nuclear staining quality of the two cytoplasmic stains. Figure 2 indicates that the excellent nuclear stain is the dominant category in both types of cytoplasmic stains.

The mean difference of cytoplasmic staining score between eosin and crimson was 0.122 and the P-value was 0.043, indicating a significant difference between the two stains. Figure 1 shows that the excellent scores of cytoplasmic stain are more predominant than good scores in all tissue types, in contrast with that of eosin, the excellent staining quality of cytoplasm is predominant in the colon and liver while in the tongue good observations are equal to the excellent observations.

From this point of view, we recommend that adding some additives to the staining solutions may enhance the staining quality. Moreover, using different nuclear stains and increasing the number of slides from varying tissue types will increase

the sensitivity of the results and give more insights about the differences and similarities between the new stain and eosin, leading to more satisfactory results. In addition, trying to better understand the dye tissue reaction and this chemical structure should be confirmed through advanced tools for more details such as NMR, HPLC, and XRF.

### Conclusion

Kermes dye could be used as a histological stain, which stains the tissue samples similar to eosin dye and its specific affinity to collagen and muscle fibers authenticates its role in the treatment of collagen and muscle disorders.

### Disclosure of potential conflicts of interest

The authors declare that they have no conflict of interest.

### Acknowledgments

This study was supported by the Deanship of Scientific Research at Prince Sattam Bin Abdul Aziz University.

### References

1. Elias W. The Anti-Diabetic Effect of Hibiscus Cannabinus Extract on the Submandibular Salivary Gland of Alloxan-Induced Diabetic Albino Rats. *Int. J. Pharm. Res. Allied Sci.* 2020; 9(2):195-202.
2. Heshmat S, W Haroun, Olfat G. Shaker, MahaKh. Abd El-wahed, Tarek I. Abd El-Galil, Eman A. Abdein, Bone marrow-derived mesenchymal stem cells attenuate the subchronic adverse effect of lead acetate on the kidney of adult albino rat. *J Adv Pharm Edu Res.* 2018; 8(1):45-52.
3. Fahdil A, Dawood AN, Olaiwy A. Adsorption of Orange G Dye from Aqueous Solutions Using Magnesium Oxide Nanoparticles. *J. Biochem. Technol.* 2018; 9(3):31-6.
4. Al-Obaidi NS, Sattar OD, Hadi FF, Ali AS, Zaki BT. Synthesis and Characterization of Some Azo Dyes Derived from 4-Aminoacetophenone, 1, 4 Phenylene Diamine and Studying its Dyeing Performance and Antibacterial Activity. *J. Biochem. Technol.* 2018; 9(4):33-42.
5. Satyanarayana, DV, Ramesh CK. Dyeing of cotton cloth with natural dye extracted from pomegranate peel and its fastness. *International Journal of Engineering Sciences & Research Technology*, 2013; 2(10): 2664-69.
6. Spodek M, Ben-Dov Y. Morphology of the first-instar nymph and adult female of *Kermes echinatus* Balachowsky, with a comparison to *K. vermilio* Planchon (Hemiptera, Coccoidea, Kermesidae). *Zookeys.* 2012; (246):11-26.
7. Cardon, D. Analyse des colorants des textiles du Taklamakan in: Debaine6Francfort, Corinne and Abduressul Idriss (eds). *Keriyā, mémoires d'un fleuve.* Paris: Editions Findakly, 2001, p. 215.
8. Sandberg G. *The red dyes: cochineal, madder and murex purple.* Lark Books Press, New York, 1997; 215 pp.
9. Amar, Z., Gottlieb, H., Varshavsky, L., Iluz, D. The Scarlet Dye of the Holy Land. *BioScience*, 2005; 55(12): 1080–1083.
10. Stathopoulou K, Valianou L, Skaltsounis A, Karapanagiotis I, Magiatis P. Structure elucidation and chromatographic identification of anthraquinone components of cochineal (*Dactylopius coccus*) detected in historical objects. *Analytica chimica acta*, 2013; 804: 264-272.
11. Ferreira ES, Hulme AN, McNab H, Quye A. The natural constituents of historical textile dyes. *Chem Soc Rev.* 2004; 33(6):329-36.
12. Meloan SN, Valentine LS, Puchtler H. On the structure of carminic acid and carmine. *Histochemie*, 1971; 27(2), 87-95.