

EXAMINATION OF WORK-RELATED MUSCULOSKELETAL DISORDERS AND ITS RELATED FACTORS AMONG FARMERS OF ASADABAD CITY IN 2015

Fariba Amiri¹, Yusef-Ali Karimi², Majid Motamedzadeh³, Manouchehr Karami⁴,
Rashid Heidari Moghadam⁵, Vida Samiei⁶

1. *MA of Health Management Services, Health Deputy, Hamadan University of Medical Sciences, Hamadan, Iran*
2. *Professional Health Expert, Health Deputy, Asad Abad Faculty of Medical Sciences, Asad Abad, Hamadan, Iran*
3. *PHD, Department of Ergonomics, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran*
4. *Professor, Social Determinants of Health Research Center, Hamadan University of Medical Sciences, Hamadan, Iran*
5. *Associate Professor, Department of Ergonomics, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran*
6. *PHD, Health Deputy, Hamadan University of Medical Sciences, Hamadan, Iran*

ARTICLE INFO

Received:

03th Jun 2017

Accepted:

29th Nov 2017

Available online:

14th Dec 2017

Keywords: *Farmers, musculoskeletal disorders, Nordic questionnaire.*

ABSTRACT

Introduction: farming has been rated one of the most widespread and dangerous occupational activities that requires tolerating inappropriate physical condition. This study has been carried out with the aim of investigating the prevalence of musculoskeletal disorders and its related factors among farmers of Asadabad city in 2015.

Materials and Methods: this study was a cross-sectional, descriptive-analytic study that has been conducted on 378 male farmers in Asadabad city of Hamadan. Nordic standard questionnaire was used for data collection in this research which was completed by trained personnel and was analyzed statistically.

Findings: The prevalence rate of musculoskeletal disorders in farmers was 6.9% in the neck area, 6.6% in shoulders, 1.1% in the elbow area, 4.5% in the wrist area, 6.1% in the back, 19% in the waist, 10.1% in the hip, 2.1% in the knee and 5% in the foot. Based on the obtained results, there was a significant relationship between the developments of musculoskeletal disorders in the foot area with age and between the development of musculoskeletal disorders in the back area and farmers' awareness level. There was also significant relationship between the development of musculoskeletal disorders in the neck, shoulder, back and waist areas with the history of disease ($P < 0.05$). There is also a significant relationship between the development of these disorders in the neck, shoulder, back, waist, knee and legs with a history of occupational accidents ($P < 0.05$).

Conclusion: according to the significant prevalence of musculoskeletal disorders among farmers, it is necessary to pursue ergonomic interventions, designing tools and performing exercise movements at defined time intervals and promoting farmers' level of knowledge in the form of educational programs of professional health.

Copyright © 2013 - All Rights Reserved - Pharmacophore

To Cite This Article: Fariba Amiri, Yusef-Ali Karimi, Majid Motamedzadeh, Manouchehr Karami, Rashid Heidari Moghadam, Vida Samiei, (2017), "Examination of Work-related Musculoskeletal Disorders and its Related Factors among farmers of Asadabad city in 2015", *Pharmacophore*, 8(6S), e-1173399.

Introduction

Work-related Muscular Musculoskeletal Disorders (WMSDs) is one of the most important occupational health issues in today's world and has almost a high prevalence in all occupations (1). Musculoskeletal disorders are defined as injuries and

Corresponding Author: Majid Motamedzadeh, PHD, Department of Ergonomics, School of Public Health, Hamadan University of Medical Sciences, Hamadan, Iran. Email: motamedzade@yahoo.com

complications in the muscles, nerves, tendons, ligaments, joints, cartilage and spinal cord (2). The World Labor Organization has estimated that the annual losses from work-related injuries and illnesses account for 10 to 15 percent of gross national product (5). Occupational injuries account for over 10 million and 8 percent of the total global damage. (6). Based on the World Labor Organization, WMSDs alone, impose the greatest economic losses (40%) on countries among other work-related injuries and illnesses (7). In Iran, musculoskeletal disorders are also ranked fourth in total permanent disability. This type of damages is more frequent in occupations with manual carrying of loads and is one of the major occupational problems; so that 19.1% of occupational injuries were due to heavy tasks and lifting loads by hand (3) Work-related musculoskeletal disorders are known to be the major factor of losing the working time, increasing the costs and human labor injuries (4). The feeling of pain and discomfort in various parts of the musculoskeletal system is the main reason of workers' absences. Studies have shown that musculoskeletal disorders are the reason of more than half of the absences in the workplace (8). The prevalence and incidence of WMSDs in industrial developing countries is more intense because the mechanization process in developed countries has reduced the level of physical activity pressure on individuals and eliminated or controlled the risk factors of WMSDs, but in the industrial developing countries where many activities are still carried out manually with the aid of using traditional workforce, workers are exposed to Biomechanical risk factors and other factors that contribute to the occurrence of WMSDs (9). Musculoskeletal disorders are a multifactorial phenomenon. In general, all risk factors can be classified into four groups: genetic factors, anatomical factors, psychosocial factors and biomechanical factors. Genetic and morphological factors (as non-interventional factors) and psychosocial and biomechanical factors (as modifiable factors) can be used to prevent the occurrence of damage (10). Cumulative trauma lesions occur when the work is repeatedly carried out by applying force and in this case, a member of the body is involved. this diseases are common in tools using. A large part of muscular and bone complications ,abnormalities and joint pain in farmers as the largest part of their discomfort, are caused by harmful ergonomic factors. which defined as, the lack of two-way coordination of the farmer and his work (11). Fleger and Wolf defined WMSDs as the main cause of living with disabilities in all countries and continents (12). Also, studying the workers' compensation in California showed that 43 percent of the total injuries reported about the farmers of the country were related to WMSDs and 25 Percentage of injuries in farmers was caused by excessive activity (13). In a study conducted by Nasrin Sadeghi et al on the saffron farmers in Gonabad city, it was found that the physical condition of the majority of saffron pickers was ergonomically at a very high level of danger and the necessity of applying methods to promptly modify their body status is suggested. (14) Agriculture is one of the most widespread and dangerous occupational activities that about 63% of the population of developing countries are engaged in it (4).

Since agriculture requires tolerating inappropriate physical condition such as bending, kneeling, crawling for harvesting, repetitive and stressful tasks , musculoskeletal disorders will be an integral part of this occupation. In the present study, the risk level of the farmers' physical condition is determined from the ergonomic view point by evaluating the physical condition of these individuals during work, and the appropriate corrective action to prevent musculoskeletal disorders was proposed. The city of Asadabad is one of the agricultural poles in Hamadan province. Agriculture is the main occupation of people who live in this city and according to the statistics of the health center, Asadabad city has covered 6500 farmers within 100 villages with resident. The above study has been carried out to investigate the prevalence of musculoskeletal disorders and related occupational risk factors among the farmers of Asadabad city in 2015.

Method of investigation:

In this cross-sectional study, 378 farmers of the agricultural area of Asadabad city of Hamadan were entered into the study and were examined. Nordic and Body map were the data gathering tools in this research which were completed by trained personnel and were analyzed statistically. The prevalence rate was used to report musculoskeletal disorders. Chi-Square test was used to examine the relationship between group variables with musculoskeletal disorder. The significance level in all tests was determined to be less than 0.05. Data were analyzed using SPSS software.

Findings:

The number of farmers participating in this study was 378 people that 84.7% (320 individuals) were married and 15.3% (58) were single. 16.7% of the participants in this study had an occupational accident and 83.3% had no experience of occupational accident. 17.5% of the participants had a history of disease other than musculoskeletal disorders influencing these disorders and 82.5% of the participants did not have such diseases. The level of awareness of the participants about professional health in agriculture was 21.7% low level of awareness, 59% with a moderate level and 19.3% had a high level of awareness. 7.9% of the participants in this study were left-handed and 92.1% were right-handed (Table 1).

Table 1: Characteristics of subjects (qualitative)

Row	variable	Variable levels	Frequency (%)
1	Marital status	Single	58(15.3)
		Married	320(84.7)
2	Occupational accident history	have	63(16.7)
		does not have	315(83.3)
3	History of disease	Have	66(17.5)
		Does not have	312(82.5)
4	Awareness level	Low	82(21.7)
		Mean	223(59)
		High	73(19.3)
5	Using hands	Left handed	30(7.9)
		Right handed	348(92.1)

The mean age range of the participants in this study was around 44.4 years old and the mean BMI of the farmers participating in the study was 17.24%. The mean weight of the participants was 70.9 kg and the mean height of them was 1.75cm, the mean work experience of the participants was 24.8 years old and the mean weight of daily loads carried by the participants was 12.3 kg (Table 2).

Table 2: Characteristics of subjects under study (quantitative):

Row	Variable	Mean	Standard deviation
1	Age	44.4	10.5
2	BMI	24.17	3.3
3	Weight	70.9	1.02
4	Height	1.75	8.41
5	work experience	24.8	11.8
6	Load carrying capability	12.3	0.8

The rate of musculoskeletal disorders in the neck area was 6.9%, 6.6% in the shoulders, 1.1% in elbow, 4.5% in the wrist area, 6.6% in the back area, 19% in the waist area, 10.1% in the hip ,1.2% in the knee and 5% in the foot area (Table 3).

Table 3: The prevalence of musculoskeletal disorders in farmers:

Row	Area with musculoskeletal disorders	Abundance (percent)
1	Neck	26(6.9)
2	Shoulder	25(6.6)
3	Elbow	4(1.1)
4	Wrist	17(4.5)
5	Back	23(6.1)
6	Waist	72(19)
7	Hips and thighs	38(10.1)
8	Knee	8(2.1)
9	Leg	19(5)

- 1- The most common musculoskeletal disorders in the waist area (65%) have happened among those who have a normal body mass index. Afterwards, the second most common musculoskeletal disorders (47%) have occurred among those with musculoskeletal disorders in the shoulder region and with a body mass higher than normal. There is no significant relationship between the developments of musculoskeletal disorders and body mass index in farmers.
- 2- The most common musculoskeletal disorders were observed among farmers who were aged between 59-50 years old and in the waist area at a rate of 21%. There is no significant relationship between the development of musculoskeletal disorders and the age group of farmers.
- 3- The highest rate of musculoskeletal disorders was among married farmers in the waist area (19.4%) and was also in the waist area for single farmers (17.2%). According to the results obtained, there was no significant relationship between the prevalence of musculoskeletal disorders and marital status in all parts under study (Pearson Chi-Square >0.05).
- 4- The results of the relationship between the prevalence of musculoskeletal disorders with an experience of occupational accidents in farmers indicated that there was a significant relationship between the prevalence of musculoskeletal disorders in various organs of neck, shoulder, back, waist, knee and legs with an experience of occupational accidents in farmers (Pearson Chi-Square <0.05) and most of the farmers with musculoskeletal disorders had an experience of occupational accidents in their organs. The highest proportion of farmers (72%) had musculoskeletal disorders in their waist area and had an experience of occupational accidents.
- 5- According to the results, the highest rates of musculoskeletal disorders have been created in the waist area and in the range of 20-29 years of work experience. There is no significant relationship between the development of musculoskeletal disorders and the level of work experience among farmers.
- 6- Among the musculoskeletal disorders in different organs of the body with a history of disease in farmers, the highest rate of disorders was related to farmers with musculoskeletal disorders in the waist area with a history of disease (46%). The results indicated that there was a significant relationship between the developments of musculoskeletal disorders in the neck, shoulder, back and the waist area with the history of disease in farmers.
- 7- The lowest level of knowledge about agricultural health was observed among farmers who had musculoskeletal disorders in the back area, which was 40% of people suffering from musculoskeletal disorders in the back area who had a low level of knowledge about agricultural health. 56% of people suffering from musculoskeletal disorders in the back area had a moderate level of knowledge about agricultural health. Afterwards, the highest statistics was related to farmers with musculoskeletal disorders in the waist (63%) with a moderate level of knowledge. Based on the results, there was a relationship between the developments of musculoskeletal disorders in the back area of farmers with their level of knowledge (Pearson Chi-Square < 0.05).
- 8- According to the results, the highest rates of musculoskeletal disorders have been observed in the waist area and among those farmers who carry a load of about 10-19 kilograms each day. There was no significant relationship between the developments of musculoskeletal disorders and the average daily load carried by farmers.

Discussion

Results of this research project, indicated that the highest rate of musculoskeletal disorders was in the lumbar area of the farmers. Since most of the farm works such as reaping or squat are done under bent condition, the pressure on the lumbar spine, especially the fifth lumbar vertebra was the leading cause of the prevalence of musculoskeletal complications. Therefore, these findings are consistent with the results of Razavi et al. (15). Also, Osborne et al, conducted a study on Irish farmers formed and he got the same results as the present study (16). The findings of this study also showed no significant relationship between musculoskeletal disorders in farmers and demographic characteristics such as body mass index. The result of this study is consistent with the research results of Rogayeh Abedini et al. (17), Hoda Rahimi Fard et al. (18), Alireza Chobineh et al. (19), Taleb Askaripour et al. (20) and Sayyed Nager Asaadi (21).

Based on the results of this study, it was found a significant relationship between age and musculoskeletal disorders in the foot area. Since the musculoskeletal problems in the foot area increases as people get older, agricultural activities will intensify this matter and these disorders will become more obvious in the leg. Therefore, the results of this study is consistent with the research results of Gibrail Nasl Seraji (22), Hoda Rahimi Fard et al. (18) Alireza Choatineh et al. (19), Somayeh Belqan Abadi et al. Davoud Nasiri Zarrin Ghobadi et al. Javad Tayife Rahimian et al.

The results showed no significant relationship between musculoskeletal disorders and marital status. This result reflects the fact that marriage and bachelorhood cannot affect musculoskeletal disorders in the agricultural sector, which this result is consistent with the research results of Seyyed Hamid Sharifnia et al (23).

It was also found a significant relationship between musculoskeletal disorders in the neck, shoulder, back, waist, knee and legs in farmers with an experience of occupational accidents. This result shows the impact of occupational accidents on the development of musculoskeletal disorders in farmers. Since a work accident can cause damage to any organs of the body, the injured member, in the case of recovery, is also prone to musculoskeletal disorders. This result is consistent with the results of Farideh Hajizadeh et al. (24) and Talib Askaripour et al. (20).

There is a significant relationship between the development of musculoskeletal disorders in the neck, shoulder, back and waist areas in farmers and the history of disease (other than diseases related to the development of musculoskeletal

disorders). Farmers with a history of disease are somewhat prone to musculoskeletal disorders which the reason can be attributed to the negative impact of disease on the farmer's body. The results of this study are consistent with the research results of Farideh Hajizadeh et al (24), Rajabali Hakamabadi et al (25), Mahnaz Solhi et al (26), Somayeh Balqanabadi et al. There is a significant relationship between the developments of musculoskeletal disorders in the back area of the farmers and their level of knowledge about ergonomics and occupational health issues which shows the direct impact of education on the incidence rate of musculoskeletal disorders. Based on the results obtained, 40% of farmers suffering from musculoskeletal disorders in their back area had a low level of knowledge about occupational health, 56% of them had a moderate level of knowledge and only 4% of farmers suffering from musculoskeletal disorders in their back area had a high level of knowledge about occupational health. Therefore, farmers who had a high level of knowledge or were somehow involved in professional health educations, were less likely to suffer musculoskeletal disorders by applying professional health principles in their work, compared to others who had a lower level of knowledge about occupational health. The results are consistent with the research results of Sayyid Abolfazl Zakrian et al. (27) and Mohammad Ronkni et al. (28). According to the results obtained in this study, there was no significant relationship between musculoskeletal disorders which were induced by farm work and the average daily load carried by farmers. The results of this research are consistent with the research results of Ehsanollah Habibi (29) and Dawood Eskandari (30). Since agriculture jobs have high repetitive motions, maintaining the sitting position with bent knees and carrying heavy loads due to their working nature, therefore knees, legs, ankles and lower parts of farmers' back and waist are prone to develop musculoskeletal disorders. Maybe it can be said that the prevalence of musculoskeletal disorders in developing countries such as Iran is more intensive, because in advanced countries, the process of automation and mechanization has partly reduced the pressures of physical activities on individual and has removed or controlled the risk of effective factors on these disorders.

But in developing countries where still many activities are carried out manually and with the aid of farmer's physical strength in the traditional manner, farmers are exposed to the risks of biomechanical factors and other factors that contribute to the development of musculoskeletal disorders. Hence, in such a condition, it is normal for these disorders to be more frequent and severe.

Conclusion

Considering that the farmers in the field of agriculture do most of their activities in a sitting position and bending at their waist, high rates of work related musculoskeletal disorders, especially in the lumbar area, is justifiable compared to other body areas. Working with a bending at the waist or sitting on the knee or standing for a long time will also cause pain in the joints of the knees, varicose veins on the legs and back pain. Therefore, in order to prevent these complications, they shall occasionally rest between shifts and do various types of movements at the workplace such as waist swing exercise, rubbing the calf muscles, sitting and getting up continuously. Farmers who have to stand for a long time should put one of their legs on a short staircase with a height of at least 35 centimeters.

It is also necessary to carry out ergonomic interventions to correct the work-related physical conditions, designing tools and performing exercise movements at certain time intervals, promoting the farmers' level of knowledge through healthcare professional education programs. Considering that training has a significant role in reducing musculoskeletal disorders among farmers, providing necessary educations on how to carry out the farm work properly, lifting and carrying loads and also, using the work and rest periods, the correct design of work tools and agricultural machinery standardization can reduce the prevalence rate of musculoskeletal disorders. As the agricultural health program in the health sector of the universities of the country is currently running with the aim of promoting farmers' health level, giving more attention to farmers' training about occupational health and safety issues in the working environment, especially ergonomic topics is recommended.

References

1. Smith DR, Leggat PA, Speare R. Musculoskeletal disorders and psychosocial risk factors among veterinarians in Queensland, Australia. *Aust Vet J.* 2009;87(7):260-5.
2. Gangopadhyay S, Ghosh T, DAS T, Ghoshal G, DAS BB. Prevalence of upper limb musculo skeletal disorders among brass metal workers in West Bengal, India. *Ind Health.* 2009;45(2):365-70. occupational ergonomics Tehran, Iran: Fanavaran
3. Chubineh A. Posture analysis methods in Press; 2004. [In Persian]
4. Bruno R. da Costa PT. Risk factors for work-related musculoskeletal disorders: a systematic review of recent longitudinal studies. *American Journal of Industrial Medicine,* March 2010;53(3): 285–323.
5. Niu S. Ergonomics and occupational safety and health: An ILO perspective. *Applied Ergonomics.* 2010;41:744-753
6. Choobineh A, Sani GP, Rohani MS, Pour MG, Neghab M. Perceived demands and musculoskeletal symptoms among employees of an Iranian petrochemical industry. *International Journal of Industrial Ergonomics.* 2009;39:766-770
7. Palmer KT, Harris EC, Linaker C, Cooper C, Coggon D. Optimising case definitions of upper limb disorder for aetiological research and prevention: a review. *Occupational and environmental medicine.* 2012;69:71-78
8. Abdoli AM. Body Mechanic and principle of work station design. Omid: 1st edition 2009: 46-50.
9. Descatha A, Roquelaure Y, Chastang J-F. Work, a prognosis factor for upper extremity musculoskeletal disorders. *Journal of Occup Environ Med* 2009; 66:351-352.

- 10-Razavi S.M. Prevalence of Musculoskeletal disorders and its risk factors among, mothers' home working. Quarterly Journal of Sabzevar University of Medical Sciences 2012-2013; 19(4)
- 11-Sadeghi S. REBA Method Posture Analysis in Saffron Pickers in 2010; 16 (1).
12. Woolf AD, Pfleger B. Burden of major musculoskeletal conditions. Bulletin of the World Health Organization. 2003;81:646-656.
- 13.N Sadeghi, Merry Ali Fanny M. Evaluation of body condition were harvested flowers and provide corrective procedures 87 years. Journal of Medical Sciences and Health Services GONABAD 1388;5(4). [Persian]
- 14 . National Safety council. Accident facts. Chicago: National Safety Council, 3rd ed. 2005.
15. Razavi SM, Bashtani A, Zarghani Sh, Tabarraie Y A survey on Prevalence of Musculoskeletal disorders and associated risk factors among Sabzevarian farmers in 2011. Quarterly Journal of Sabzevar University of Medical Sciences 2014;20(5):766-772.
- 16.Aoife Osborne, Catherine Blake, David Meredith, Anne Kinsella, James Phelan, John McNamara, and Cairiona Cunningham. AMERICAN JOURNAL OF INDUSTRIAL MEDICINE 2012; 8: 1-8.
17. Abedini R, Choobineh A. Musculoskeletal Disorders Related to Patient Transfer in Hospital Nursing Personnel 2012 ;8(3):385-396.
18. Rahimifard H, Heidari HR. Study of Prevalence of Musculoskeletal Disorders and Identification of Factors affecting them among Furniture Industry workers with Rapid Entire Body Assessment (REBA) Method .7(6)2011:916-925.
19. Choobineh AR, Mokhtarzadeh A, Salehi M, Tabatabaei SH . Ergonomic evaluation of exposure to musculoskeletal disorders risk factors by QEC technique in a rubber factory.
20. Askaripoor T, Kermani A, Jandaghi J, Farivar F. Survey of Musculoskeletal Disorders and Ergonomic Risk Factors among Dentists and Providing Control Measures in Semnan. Journal of Health 2013.4(3):241248
21. Assadi, SN. Assessment of occupational risk factors for upper limbs musculoskeletal disorders in educational employees. Iranian Journal of Research in Environmental Health. 2015;1 (2): 119-124.
- 22.Nasl Seraji J, Hossini MH, Shahtaheri SJ, Golbabai F. Evaluation of ergonomic postures of dental professions by Rapid Entire Body Assessment(REBA),in Birjand ,Iran. Journal of Dentistry of Medical Sciences 2005;18(1):61-67
23. Bolghanabadi S, Dehghan H , Mehdi Pour. The relationship between musculoskeletal disorders, stress and fatigue in workers of a food industry. Journal of Ergonomics,2014 ;2(1): 54-63.
24. Nasiry Zarrin Ghabae D, Haresabadi M , Bagheri nesami M. Work-Related Musculoskeletal Disorders and Their Relationships with the Quality of Life in nurses. Journal of Ergonomics, 2016;4(1): 39-46.
25. Tayefe Rahimian J, Choobineh A , Dehghan N. Ergonomic Evaluation of Exposure to Risk Factors of Musculoskeletal Disorders in Welders. Journal of Ergonomics, 2014;5(3): 18-26.
25. Tayefe Rahimian1, Alireza Choobineh2, Naser Dehghan
26. Sharif nia SH, Haghdoost AA, Hajihosseini F, Hojjati H. Relationship between the musculoskeletal disorders with the ergonomic factors in nurses.2012;12(4):372-379
27. Hajizadeh F, Motamedzade M, Golmohammadi R, Soltanian AR. Work ability assessment and its relationship with severity of musculoskeletal . disorders among workers in a cement plant. Journal of Occupational Hygiene Engineering 2015; 2(2):15-22.
28. Solhi M, Khalili M, Eshragian A, Zakerian A. Prevalence of symptoms of musculoskeletal disorders and predictors of proper posture among computer users based on stages of change model in computer users in central Headquarter, Tehran University of Medical Sciences. Iran Occupational Health 2015; 11(5):43-52
29. Rokni M , Abadi MH , Saremi M, Mir Mohammadi MT . Prevalence of musculoskeletal disorders in nurses and its relationship with the knowledge of ergonomic and environmental factors. Journal of Gorgan University of Medical Sciences 2016;18(1):128-132.
30. Habibi E , Kazemi M , Safari SH, Hassanzadeh A. The Relationship between Lifting Capacity with the NIOSH Equation and the Risk of Musculoskeletal Disorders with the RULA Method in Health Service Personal of Isfahan, Iran. Journal of Health System 2012; 8(1):131-137.